

Sr.No.	LIST OF FIGURES	Pg.No.
I.1	Generalised geological map of Rajasthan and North Gujarat, the inset open square represents the study area.	3
II.1	Geological map of Phalasia and Jharol area, Southern Rajasthan.	37
III.1	Structural map showing various axial traces around Phalasia and Jharol area, southern Rajasthan.	58
III.2	Tectonic map of Phalasia area.	59
III.3	Geotectonic map of Kotra-Jharol area.	61
III.4a&b	Fracture controlled pattern of Kaliawa Nenbara shear zone	63-64
III.5	Geological cross-section along the latitude $24^{\circ}15'$ showing co-axial $F_1$ and $F_2$ folds. $S_1$ -axial trace of $F_1$ and $S_2$ -axial trace of $F_2$ .	69
III.6(A)	Schematic illustration of veins deforming during superimposed simple shear, determining simple shear.	83
(B)&(C)	Schematic illustration showing the development of sigmoidal tension gashes.	83
(D)	Development of asymmetrical folds during simple shear.	83
III.7	Veins developed on account of the flexural slip mechanism.	84

III.8	The movement of the border grains are governed by the main shear which is sinistral, and the inner grains have been rotated on account of the movement of the adjacent grains.	120
III.9	Analogy between the use of microfolds adjacent to porphyroblasts in metamorphic rocks and phenocrysts in volcanic rocks to determine the sense of shear.	122
III.10	Microscopic sketch showing sinistral shear sense locally developed due to pure shear deformation which later on changed into simple shear. This is few of such features survived after the area has undergone simple shear deformation.	124
III.11	Microscopic sketch illustrating two generations of garnet grains, pre-Kinematic and Synkinematic growths.	125
III.12	Rotation of $S_e$ with respect to stationary $S_i$ due to coaxial deformation.	132
III.13	Microscopic sketch showing stretched garnet porphyroblast splitting the mica flakes in garnet-mica-schist.	132
IV.1	Map of the Aravalli-Delhi belt showing distribution of types and zones of metamorphism.	155

IV.2	Pressure-temperature diagram showing the fields of the various metamorphic facies.	160
IV.3	Hypothetical curves of Univariant equilibrium for reactions leading to appearance of sillimanite in almandine-amphibolite facies of Turner and Verhoogen, 1962.	164
IV.4	Sketch map showing polymetamorphism of Jharol group of rocks.	176
V.1	Basin evolution of the Jharol group and the main Aravalli sequence.	183
V.2	Sketch showing the sense of movement of the Rakhabdev-Dungarpur transform fault and the Modi-Dikaliya strike-slip fault.	187
V.3	Tectonic model illustrating shear deformation in the southern Rajasthan-North Gujarat region.	195
V.4A	Sinistral shear movement due to subduction tectonics involving mantle viscous drag resulted in the development of decollement fold stack ( $AF_2$ folding).	198
V.4B	$AF_3$ folding Superimposed on $AF_2$ folds, controlled mainly by simple shear mechanism accompanied by flattening.	199