LIST OF PLATE

A. Topographical relief of Jaisalmer basin, Locality : NW of Jaisalmer.
B. Sand dunes covering western part of Jaisalmer basin Locality : Sam'village.
Plate No. I.2
A. Architectural Panorama of Golden city of Jaisalmer showing economic importance of yellow, calcareous sandstone and sandy limestone of Jaisalmer Formation; (Salimsingh Ji-Ki-Haweli, Jaisalmer).

B. Architectural panorama of Golden city of Jaisalmer showing detail of arches and carvings; (Gaj palace, Jaisalmer).

Plate No. IV. 1

- A. Ferruginous, gritty, poorly sorted sandstone with silicified fossilwood (not seen in photograph). Odania Member, Lathi Formation; Locality: Fossilwood park, Akal.
- B. Fine grained, ferruginous, well sorted sandstone showing NNE-SSW trend of joints, Odania Member, Lathi Formation; Locality: Fossil-Wood park, Akal.

L

ي.

PLATE

Plate No. I.1

.

5

5

Page No.

9

9

46

46

PLATE

Plate No. IV. 2

A. Cross bedded fine grained sandstone with thin intercalation of marl and hard siltstone, Odania Member, Lathi Formation; Locality: 1 Km NE of Akal, road section.

48

 B. Horizontal, thinly bedded, ferruginous siltstone and light brown hard compact, fine grained sandstone. At places sandstone shows lensoid nature. Odania Member Lathi Formation; Locality: 1 Km NE of Akal, road section. 48

Plate No. IV. 3

- A. Ferruginous marl bed on the top of Lathi Formation mark the gradational contact between overlying Hamira Member (Jaisalmer Formation; lower most part) and underlying Lathi Formation; Locality: North of Thaiyat village.
- B. Yellowish brown, calcareous claystone showing ptygmatic and micro-folds within the ferruginous claystone near Fatehgarh fault at the base of Lathi Formation; Locality: NW of Fatehgarh village.

Plate No. IV. 4

- A. Lower contact of Jaisalmer Formation with underlying Lathi Formation, marked by reddish brown marl bed. Locality: Near Thaiyat.
- 51

50

50

B. Thinly bedded, light yellow, calcareous sandstone and marl, Hamira Member, Jaisalmer Formation: Locality: Near Thaiyat, road section.

51

Plate No. IV. 5

A. Yellowish brown, moderately hard, marl and light

۰ ـ

brown cross-laminated, fine grained, sandstone, Hamira Member, Jaisalmer Formation; Locality: 7 Km SE of Jaisalmer town.

B. Cross bedded fine grained, fossiliferous, sandstone, Joyan Member, Jaisalmer Formation; Locality: 7 Km SE of Jaisalmer town.

Plate No. IV. 6

- A. Light yellow, fine grained, sandstone (Basal part of Fort Member) showing faint cross lamination, Fort Member, Jaisalmer Formation.
- Light yellow, hard sandy colitic limestone showing herring bone cross lamination, indicating tidal effect during deposition, Fort Member, Jaisalmer Formation: Locality : Fort Escarpment around Jaisalmer.

<u>ج</u>د

Plate No. IV. 7

- A. Intraformational conglomerate bed exposed in Badabag section, Locality: Badabag. 59
- B. Cross bedded, calcareous sandstone, exposed in type section of Badabag Member; Locality: Badabag village. 59

Plate No. IV. 8

- A. Light yellow, fine grained moderately sorted crossbedded, sandstone (lower) overlain by flaggy oolitic limestone, Badabag Member, Jaisalmer Formation. 61
- B. Ferruginous fossiliferous oolitic claystone, overlying hard, compact micritic limestone, suggesting unstable shelf condition of deposotion, Kuldhar Member, Jaisalmer Formation; Locality; Kuldhar Nala section Near Kuldhar.

61

56

56

58

Page No.

Plate No. IV. 9

- Ferruginous, argillaceous, limestone with band of shale; Α. 62 exposed in Kuldhar Nala section near Kuldhar village.
- в. Gypsiferous shale representing topmost unit of Kuldhar Member, exposed in Kuldhar Nala section, suggesting shallowing of sea during deposition of upper part of Jaisalmer Formation; Locality: Near Kuldhar village.

62

Plate No. IV. 10

Light brown, thinly bedded, very fine grained Α. sandstone, Kuldhar Member, Jaisalmer Formation; Locality: Near Biramsar River section.

63

в. Conglomerate bed overlying highly calcareous light brown, sandstone, upper part of Jaisalmer Formation (Kuldhar Member), marking unconformable contact between Jaisalmer and Baisakhi Formation; Locality : 1/2 Km NW of Biramsar village in Biramsar river section.

63

Plate No. IV. 11

Α. Yellow, fine to very fine grained, mod. hard. well sorted, argillaceous sandstone intercalated with light yellow claystone, Ludharwa Member, Baisakhi Formation; Locality: NW of Ludharwa village.

67

Thinly bedded, light brown, fine grained, compact, sandstone; Rupsi Member, Baisakhi Formation; Locality : Β. North of Baisakhi village.

67

Plate No. IV. 12

Contact of thinly laminated very fine sandstone of Baisakhi Formation and gritty ferr.sandstone of Bhadasar Α. Formation (Kolar Dungar Member), Locality: Bhadasar Escarpment; NW of Bhadasar village. . 70

. .

B. Alternation of fine grained, light brown, sandstone and siltstone, showing thin lamination of ferruginous clays; Kolar Dungar Member, Bhadasar Formation; Locality : Bhadasar Village, Bhadasar Escarpment.

70

Plate No. IV. 13

A. Light brown, fine grained, hard and compact sandstone showing Cross lamination, Mokal Member, Bhadasar Formation; Locality : Near Mokal village.

71

71

B. Ferruginous sandstone, hard, compact, highly jointed, showing slump like structure, Mokal Member, Bhadasar Formation; Locality : NNE of Mokal village.

Plate No. VI. 1

A. Photomicrograph of quartz wacke showing moderately sorted fine grained aggregates of quartz, mica flakes set in clayey matrix, tiny crystals of calcite observed within the matrix, Thaiyat Member, Lathi Formation; X55; XN.

129

129

B. Photomicrography of Calcareous quartz arenite showing bimodal populations of quartz grains suggesting existance of more than one provenance, odania Member, Lathi formation, 45;XN.

Plate No. VI. 2

....

- A. Photomicrograph of calcareous quartz arenite showing quartz overgrowth and sparry calcite cement, Odania Member, Lathi Formation; X45; XN.
- 131

131

B. Photomicrograph of quartz arenite showing bent mica, strained and fractured quartz suggesting effect of tectonic episode, lower part of Odania Member, Lathi Formation; X45; XN. Plate No. VI. 3

A. Photomicrograph of argillaceous siltstone microfacies of Odania Member, Lathi Formation showing aggregate of fine to very fine sand grade quartz set in clay matrix, at places influx of ferruginous matter and tiny particles of dolomite within the matrix is also observed; X55; XN.

133

B. Photomicrograph of calcareous quartz arenite showing aggregate of quartz, plagioclase feldspars and chert fragments, Hamira Member, Jaisalmer Formation; X55; XN.

133

Plate No. VI. 4

A. Photomicrograph showing subrounded occasionally subangular and rounded quartz bound by ferruginous matrix. Partially quartz grains are cemented by zoned ferroan dolomite and calcite, Hamira Member, Jaisalmer Formation; X55; XN.

135

B. Photomicrograph showing dominantly angular to subangular quartz set in ferruginous clay matrix, polycrystalline quartz are also seen, Kuldhar Member, Jaisalmer Formation; X55; XN.

135

Plate No. VI. 5

A. Photomicrograph showing fine to medium grained, ill sorted quartz and occasional feldspar in quartz arenite microfacies. The cements are mainly ferruginous. At places calcite cements are also observed. The quartz boundaries are corroded by calcite, Ludharwa Member, Baisakhi Formation; X45; XN.

139

1

B. Photomicrograph of calcareous quartz wacke comprising dominantly fine grained quartz set in calcareous clay

PLATE

matrix, occasional feldspar and ferruginised shell fragments are also observed. The quartz boundaries are generally corroded by calcite, Baisakhi Member Baisakhi Formation; X45;XN.

139

Plate No. VI. 6

A. Photomicrograph of quartz wacke comprising fine to medium grained, mainly monocrystalline occasionally polycrystalline quartz and feldspar (microcline) set in ferruginous clay matrix. At places patches of calcite are also been observed. Mokal Member, Bhadasar Formation; X45; XN.

142

142

B. Photomicrograph of quartz arenite showing aggregates of moderately sorted grains of quartz, feldspar set in admixed matrix of ferrugnious clays and calcareous matter. Corroded quartz boundaries, overgrowth of quartz and ferrugnious rim around the quartz grains are observed, Kolar Dungar Member, Bhadasar Formation; X55; XN.

Plate No. VI. 7

A. Photomicrograph of mudstone microfacies comprising micrite and scattered small skeletal shell fragments, Jaisalmer Formation, Vikran Nai Structure; X55; XN.

148

B. Photomicrograph showing mud supported microfacies of pelletal wackestone comprising of fine grained micritic pellets along with smaller forams. At places micritic matrix shows sparitisation, Kuldhar Member, Jaisalmer Formation; X55; XN.

148

۱

Plate No. VI. 8

A. Photomicrograph showing mud-supported, bioclastic wackestone microfacies. The bioclasts are generally

. •

algal fragments and crinoids. The matrix is microspar at places showing selective dolomitisation, Badabag Member Jaisalmer Formation; X 25; XN. 150 Β. Photomicrograph showing grain supported pelloidal bioclastic packstone microfacies. The bioclasts are Jaisalmer completely sparitised, Fort Member, Formation; X65; XN. 150 Plate No. VI. 9 showing grain Α. Photomicrograph supported pelletal The pellets are well sorted. grainstone microfacies. very fine grained and micritic in nature. Occasional milliolids are also observed alongwith pellets. The interspaces are occupied by calcite, Jaisalmer Formation;

152

B. Photomicrograph showing grain supported colitic grainstone microfacies composed of spheroidal toellipsoidal shape of colites showing concentric as well as radial structures, Jaisalmer Formation, Sadewala Structure; X55; XN.

Vikran Nai Structure; X55; XN.

152

Plate No. VI. 10

- A. Photomicrograph of coralline-oolitic grainstone microfacies observed in upper part of Jaisalmer Formation in Vikran Nai Structure X55; XN.
- B. Photomicrograph showing grain supported Oncolite-oolitic grainstone microfacies comprising algal coated oncolite grains. The nuclei of the grains are micrite and fossil fragments. At places microstylolitic junction has been noticed along the grains boundaries. Fracture and solution porosities have also been observed in this microfacies. Jaisalmer Formation, Sumarwali Talai Structure; X20; PP.

154

Plate No. VIII. 1

A. Photomicrograph showing corroded quartz boundaries by calcite cement in the early diagenetic phase, later on complete replacement of calcite by iron oxide (hematite). Hematite cement causes the complete destruction of porosity, Mokal Member, Bhadasar Formation; X50; PP.

222

222

B. SEM photomicrograph of lower sandstone facies in Fort Member, showing grain packing predominantly tangential, point and line contacts which create floating appearance in thin section.

Plate No. VIII. 2

A. Photomicrograph showing clay rims around the detrital quartz grains in the sandstones section, lower part of Fort Member, Jaisalmer Formation. X50, PP.

231

B. Photomicrograph showing replacement of feldspar by calcite and selective replacement of calcite by iron oxide, Kolar Dungar Member, Bhadasar Formation; X50; XN.

231

Plate No. VIII. 3

A. Photomicrograph showing quartz overgrowth formed as the first generation of cement and followed by calcite which generally filled the pore spaces, Thaiyat Member, Lathi Formation; X50; XN.

232

B. Photomicrograph showing microstylolite and fracture porosity in Oncolite-Oolite grainstone facies of subsurface section in Upper part of Jaisalmer Formation, Sumarwali Talai Structure; X20; PP.

232

Plate No. VIII. 4

- Calcareous quartz arenite showing crystalline replacement Α. of dolomite with considerable iron. Dolomite probably formed in late stage of diagenesis, Joyan Member, Jaisalmer Formation, X55; XN. Β. Photomicrograph showing neomorphic development of calcite in micritic mass of mudstone facies of subsurface section of Jaisalmer Formation, Vikran Nai Structure; X55, XN. 235 Plate No. VIII. 5 Α. Photomicrograph showing neomorphic development of calcite from ferruginous micrite matrix in oolitic
 - wackestone facies, Jaisalmer Formation. Kuldhar Member; X25 ; XN.
 - в. Photomicrograph showing selective dolomitisation resulting in scattered well development rhombs of dolomite in rare association with anhydrite, middle part of Vikran Nai Structure; Jaisalmer Formation; X55; XN.

Plate No. VIII. 6

- Α. Herring bone cross bedded structure observed in sandy oolitic limestone of Fort Member, Jaisalmer Formation.
- 238
- в. Photomicrograph showing fine fringing fibrous calcite, oomoldic and dissolution perosities in colitic grainstone facies of upper part of Jaisalmer Formation (1855m), Bhuana Structure; X85; XN.

Plate No. VIII. 7

Α. Photomicrograph showing microstylolitic junction and development of fracture porosity in oncolite-oolite

235

237

237

238

ł

		grainstone microfacies at the top of Jaisalmer Formation in Sumarwali Talai Structure; X 50; PP.	240
	в.	Photomicrograph showing microstylolite development at the junction of the grains in pelloidal-wackestone facies indicative of deep burial ^{dj} agenesis, Lower part of Jaisalmer Formation, Bhuana Structure; X70 ; XN.	240
Plate	No. V	/III. 8	
	Α.	Photomicrograph showing equant blocky calcite acting as pore occluding cement in oolitic grainstone facies of Sadewala Structure; Jaisalmer Formation; X70 ; XN.	242
	в.	Photomicrograph showing development of blocky calcite occluding the intergranular spaces in coralline-oolitic grainstone facies, middle part of Jaisalmer Formation, Vikran Nai Structure; X60; XN.	242
Plate	No. V	/111. 9	
	Α.	Photomicrograph showing syntaxial overgrowth rim cement on echinoderm plate in oolitic grainstone microfacies observed in the upper part of Jaisalmer Formation, Vikran Nai Structure; X75 ; XN.	244
	в.	SEM photomicrograph showing intergranular porosity in Pelletal grainstone facies (Left corner of Photograph). In the lower part, note calcite spar.	244
Plate	No. V	/III. 10	
	Α.	SEM photomicrograph showing infilling of intergranular spaces by Kaolinite clay, Lathi Formation.	246
	Β.	Photomicrograph of calcareous quartz arenite showing well sorted quartz grains, chert and rare oolite with calcite cement and selective dolomitisation, basal part of Jaisalmer Formation, Sumarwall; Talai Structure; X70; XN.	246

,

ı

- -

Plate No. VIII. 11

.....

- SEM photomicrograph showing microporosity development Α. due to dissolution phenomenon. 247
- Β. SEM photomicrograph of varible dissolution texture in 247 Lower sandstone facies in Badabag Member.

Plate No. VIII. 12

- Α. SEM photomicrograph showing microstylolite in the calcite spar, developed due to dissolution. 249
- в. SEM photomicrograph showing calcite, occluding the intergranular spaces in quartz arenite microfacies, though minor amount of porosity is visible in this section. 249

Plate No. VIII. 13

- Α. Photomicrograph showing ferruginous cement replaced by calcite as evidenced by, presence of remnant iron oxide, Thaiyat Member, Lathi Formation; X50; XN.
 - 251
- Β. SEM photomicrograph showing textural relationship, clay coating on quartz grains and lining pores; Sandstone facies of upper Lathi Formation.