

CONTENTS

<u>Title</u>	<u>Page no.</u>
CHAPTER 1 INTRODUCTION	1
PURPOSE AND SCOPE	2
OBJECTIVES	3
STUDY AREA	4
Location	4
Communication	4
Physiography and drainage	5
Climate	5
Flora	6
Fauna	6
People and occupation	7
APPROACH AND METHODOLOGY	7
CHAPTER 2 REGIONAL GEOLOGY	9
STRUCTURE AND TECTONICS	10
MESOZOIC STRATIGRAPHY	11
Jhurio Formation	12
Jumara Formation	13
Jhuran Formation	14
Bhuj Formation	14
TERTIARY STRATIGRAPHY	15
Matanomadh Formation	16
Naredi Formation	16
Harudi Formation	17
Fulra Limestone	18
Maniyara Fort Formation	18
Khari Nadi formation	19
Chhasra Formation	20
Sandhan Formation	20
GREAT RANN OF KACHCHH BASIN	21
CHAPTER 3 MATERIALS AND METHODOLOGY	25
PARTICLE SIZE ANALYSIS	26
SEDIMENTOLOGY	27
ENVIRONMENTAL MAGNETICS	27
PALYNOLOGY	33
RADIOCARBON DATING/ CHRONOLOGY	34
CHAPTER 4 CHRONOLOGY AND AGE MODEL OF THE CORES	36
THE DHORDO CORE AND THE BERADA CORE	36
AMS RADIOCARBON DATING OF THE CORES	37

Organic carbon dating of the cores	38
Inorganic dating of the cores	42
AGE MODEL OF THE DHORDO CORE	45
Greenlandian stage - Early and Late-Holocene	47
Northgrippian stage -Middle Holocene	47
Meghalayan stage - Late Holocene	48
AGE MODEL OF THE BERADA CORE	49
Greenlandian stage /Lower and Early Holocene	49
Northgrippian stage/ Mid- Holocene stage	50
Meghalayan stage/ Late Holocene stage	50

CHAPTER 5 PHYSICAL STRATIGRAPHY AND SEDIMENTOLOGICAL CHARACTERISTICS

RECOGNISING SEDIMENTARY STRUCTURE IN SOFT SEDIMENT CORES	53
SEDIMENTARY CHARACTERISTICS OF THE CORES	53
DHORDO CORE	54
Unit 1 Lower intertidal to subtidal flat deposits	56
Unit 2 Upper tidal flat deposit (Intertidal)	59
Unit 3 Supra tidal flat deposits	60
BERADA CORE	61
Unit 1 Fluvial sediment deposits	62
Unit 2 Estuarine deposits (marshy deposition)	64
Unit 3 Subtidal flat deposits	66
Unit 4 Intertidal flat deposits (mix tidal to mud tidal zone)	67
Unit 5 Supratidal flat deposits	68
STRATIGRAPHY AND DEPOSITIONAL CONDITIONS IN GRK	69

CHAPTER 6 GRAIN SIZE ANALYSIS AND TEXTURAL CLASSIFICATION

GRAIN SIZE PARAMETERS	73
Mean Size (Mz)	73
Standard Deviation (σI)	73
Skewness (SkI)	74
Kurtosis (KG)	74
GRAIN SIZE ANALYSIS IN GREAT RANN OF KACHCHH (GRK)	74
DHORDO CORE	75
Grain size variations	75
Greenlandian Stage	75
Northgrippian Stage	77
Meghalayan Stage	80

Textural classification of Dhordo core	81
BERADA CORE	84
Grain size variations	86
Greenlandian Stage	86
Northgrippian Stage	87
Meghalayan Stage	87
Textural classification of Berada core	90
CHAPTER 7 ENVIRONMENTAL MAGNETIC STUDIES	94
ENVIRONMENTAL MAGNETIC PARAMETERS	94
Concentration parameters	96
Grain size parameters	96
Mineralogy of magnetic minerals	96
DHORDO CORE	97
Concentration parameters	97
Grain size parameters	98
Mineralogy of magnetic minerals	100
PALAEOENVIRONMENTAL IMPLICATIONS IN DHORDO CORE	101
Greenlandian Stage	101
Northgrippian Stage	102
Meghalayan Stage	103
BERADA CORE	104
Concentration parameters	104
Grain size parameters	105
Mineralogy of magnetic minerals	107
PALAEOENVIRONMENTAL IMPLICATIONS IN BERADA CORE	109
Greenlandian Stage	109
Northgrippian Stage	110
Meghalayan Stage	110
CHAPTER 8 PALYNOLOGICAL STUDIES	111
PALYNOLOGICAL STUDIES IN GRK	112
DHORDO CORE	112
Palynological studies on Dhordo core	113
Greenlandian to Northgrippian Stage	113
Northgrippian Stage	116
Meghalayan Stage	117
Palaeoenvironmental conditions of Dhordo core	118
BERADA CORE	123
Palynological studies on Berada core	123
Greenlandian to Northgrippian Stage	124
Northgrippian Stage	127

Meghalayan Stage	128
Palaeoenvironmental conditions in Berada core	130
CHAPTER 9 DISCUSSION AND SYNTHESIS	134
THE GREAT RANN OF KACHCHH	134
SEDIMENTATION PATTERN IN GRK	135
Greenlandian Stage	135
Northgrippian Stage	136
Meghalayan Stage	136
HOLOCENE SEDIMENT ACCUMULATION CURVE	137
Greenlandian Stage	138
Northgrippian Stage	141
Meghalayan Stage	142
PALAEOENVIRONMENTAL CONDITIONS IN GRK	142
Greenlandian Stage	143
Northgrippian Stage	146
Meghalayan Stage	147
CHAPTER 10 CONCLUSIONS	149
REFERENCES	152
LIST OF PUBLICATIONS	172

LIST OF FIGURES

Figure no.	Description	Page no.
Figure 1.1	Location map of Great Rann of Kachchh basin showing geomorphic zone based on modern inundation pattern and present and paleo river channels (modified after Khonde, 2014). Red square (solid) a and b represents the Dhordo core and Berada core location. Pachham Island (Pa), Khadir Island (Kh), Bela Island (Be), Chorar Hill (ch)	2
Figure 2.1	Geological map of the Kachchh basin (after Biswas, 1993). Map redrawn by Patidar (2010).	10
Figure 3.1	Schematic view of the multi-proxy studies carried out during the study	26
Figure 3.2	X-Ray radiographs of all core pipes from Dhordo Core site. Note the excellent core recovery and fine scale laminations in the core pipes.	29
Figure 3.3	X-Ray radiographs of all core pipes from Berada core site. Note the core excellent recovery and fine scale laminations in the core pipes.	30
Figure 3.4	Bartington MS2WFP Susceptibility Meter (noise level $\sim 3 \times 10^{-9} \text{ m}^3 \text{ kg}^{-1}$ for a 10 g sample). The instrument is used to calculate concentration of magnetic mineral	32
Figure 3.5	AGICO JR-6 Spinner Magnetometer (JR-6 Dual Speed Spinner Magnetometer) Measurement over 11 magnitudes ($10^6 - 10^4 \text{ A/m}$).	32
Figure 3.6	ASC Scientific D2000 Alternating Field Demagnetizer	33
Figure 3.7	Demonstration of complete working model of Accelerator mass spectroscopy used for radiocarbon dating.	35
Figure 4.1	Graphical representation of the all the organic carbon ages from Dhordo core processed at IUAC (Inter University Accelerator Centre, New Delhi). Note the reversal and overestimation of the ages obtained.	40
Figure 4.2	Graphical representation of the all the organic carbon ages from Berada core processed at IUAC (Inter University Accelerator Centre, New Delhi). Note the reversal and overestimation of the ages obtained.	41

Figure 4.3	Age model of the Dhordo core reconstruction using the Bayesian age-depth model. Note the high rates of sedimentation throughout the core, especially during the Greenlandian stage.	46
Figure 4.4	Berada age depth model calculated using Bayesian age-depth model. The graphical representation of the inorganic samples dated at various facilities.	49
Figure 5.1a	Photographs of 11 core pipes out of 24 pipes are shown, each pipe is shown in the combination of the split pipe photograph acquired manually (right) along with their respective x ray photographs (left). Scale is attached along with photograph of each core pipe.	55
Figure 5.1b	Photographs of 13 core pipes out of 24 pipes are shown, each pipe is shown in the combination of the split pipe photograph acquired manually (right) along with their respective x ray photographs (left). Scale is attached along with photograph of each core pipe.	55
Figure 5.2	Complete shells of Bivalves and Gastropods recovered from the Dhordo core.	58
Figure 5.3	Broken shell present in the muddy sediment recovered from the Dhordo core.	58
Figure 5.4a	Selected photograph of sedimentary structure a) 0.3 – 1 m, recent soil sediments deposits b) 1.5 – 2 m gradual contact or erosional contact separating undisturbed sediments from well preserved sediment b') 2.1 – 2.2 m, flaser structure (bit core) c) 2.4 – 3.1 m, greyish yellow and greyish dark very fine sand to silt, containing sand–mud couplets, thin rhythmic laminations d) 3.6 – 3.9 m, organic matter dissolution mark e) 4.9 – 6.2 m, colour alteration of grey and yellowish grey sand-clay f) 7 – 7.9 m, laminated sand mud caplets g) 10.4 – 12.3 m highly compacted lenticular bedding h) 13.5 – 13.8 m, cross bedding i-j) 14.2 – 15.1 m; 16.7 - 17.2 m, parallel lamination.	59
Figure 5.4b	Selected photograph of sedimentary structure k) 18.3 – 18.7 m, clayey silt with very fine sand couplets, cross laminations and bi-directional current structures l) 19.1 – 19.7 m, transition from bi-directional cross bedding to high energy cross laminations m) 19.7 – 20.2 m, thickly interlaminated to thinly interbedded sand and mud couplets are common n) 20.2 – 20.4 m, massive brownish lump deposit with broken shells & (Figure. 5.3) o) 25 – 25.3 m, presence of gravel within deformed structure p) 25.5 – 25.9 m, massive deposit with alternate grey and brown colour sediment, presence of complete gastropod shell q-r) 29.7 – 30.1 m; 31.1 – 33.2 m, broken shell layer, scattered organic matter throughout	60

core pipe s) 35.8 – 37.5 m, structure less greenish to brown clay deposit t-u-v) 39.1 - 40 m; 41.6 – 43.2 m; 47.9 – 48.9 m, burrow activity along with track and trail present w) 57.4 – 58.1 m highly compact silt layers and laminations.

- Figure 5.5a Photographs of 8 core pipes out of 15 pipes are shown, each pipe is shown in the combination of the split pipe photograph acquired manually (right) along with their respective x ray photographs (left). Scale is attached along with photograph of each core pipe. 62
- Figure 5.5b Photographs of 8 core pipes out of 15 pipes are shown, each pipe is shown in the combination of the split pipe photograph acquired manually (right) along with their respective x ray photographs (left). Scale is attached along with photograph of each core pipe. 63
- Figure 5.6 Selected photographs and X-ray radiographs of the slab samples obtained from Berada cores. a) 1.35 – 0.3 m, uniform deposit of sand b) 1.75–2.30 m, sand clay brownish colour sediments with poorly laminated sediments c) 4.75–6.50 m, colour alternation of greenish grey to brown clayey sand, dotted line is drawn to represent the lithological colour variation. d) 17.6 – 15.8 m, high mud content, dotted line representing the highly bio turbidite sediment unit overlying homogenous deposition, high mud content. e) 24–22.1 presence of gravel within deformed structure f) 27.25–25.60 m, represents rootlets, note the upward increasing sand mud couplets: lenticular to wavy structure with increase in sand content. g) 35.8 –35.6 m, increasing upwards mud and sand intercalation. h) 39.1–38.1 m, extensive burrowing activity with scattered organic matter i) 39.6 –39.1 m, peat layer within clayey sediments having gradual contact with the underling homogenous undisturbed sediment deposit. j) 44.3 – 43.2 m, hard compact brownish oxidised clay with laminations. 64
- Figure 5.7 Complete shells of Gastropods and Bivalves recovered from the Berada core. 65
- Figure 5.8 Broken shells recovered from the Berada core. 65
- Figure 5.9 Stratigraphic cross-sections with isochrons of the Great Rann of Kachchh basin (X – X'), including detailed information about the lithology, dating, sedimentary facies and depositional conditions. Isochrons are drawn based on radiocarbon ages. Note the termination of fluvial and estuarine depositional condition before Dhordo core. 71

- Figure 6.1 Grain size variation of sand silt clay along with lithological variation of the Dhordo core from GRK basin. 76
- Figure 6.2 a) Diagram of mean grain size versus sorting of sediments from Dhordo core showing different energy and depositional environmental conditions (modified after Tanner (1991). b) Bivariate plot of grain-size distribution parameters of sorting and skewness showing energy of depositional environment. (after Bjørlykke, 2010). c) Line graphical presentation of mean, sorting and skewness along with the depths. 77
- Figure 6.3 a) Lithology of the Dhordo core from GRK basin along with chronological age of the core. b) Triangular plot of grain size distribution in Dhordo core and its textural classification following Blott and Pye, (2011). c) Enlarge view of the textural attributes of GRK sediments in Dhordo core. Blue square box with cross showing sediments deposited during Meghalayan Stage, Yellow triangle reflects Northgrippian Stage and red cross showing Greenlandian Stage. 82
- Figure 6.4 Grain size variation of sand silt clay along with lithological variation of the Berada core from GRK basin. 85
- Figure 6.5. Diagram of mean grain size versus sorting of sediments from Berada core showing different energy and depositional environmental conditions (modified after Tanner (1991). b) Bivariate plot of grain-size distribution parameters of sorting and skewness showing energy of depositional environment. (after Bjørlykke, 2010). c) Line graphical presentation of mean, sorting and skewness along with the depths. 88
- Figure 6.6 a) Lithology of the Berada core from GRK basin along with chronological age of the core. b) Triangular plot of grain size distribution in Berada core and its textural classification following Blott and Pye, (2011). c) Enlarge view of the textural attributes of GRK sediments in Berada core. Blue square box with cross showing sediments deposited during Greenlandian Stage, red square showing Northgrippian Stage and Yellow square reflects Meghalayan Stage. 91
- Figure 7.1 Temporal scale variability in concentration environmental magnetic parameters in Dhordo core, western India. 98

Figure 7.2	Temporal scale variability in grain size environmental magnetic parameters in Dhordo core, western India.	99
Figure 7.3	Temporal scale variability in mineralogical environmental magnetic parameters in Dhordo core, western India.	100
Figure 7.4	Scatter plots between a) χ_{ARM} vs χ_{lf} , b) χ_{lf} vs S-ratio and c) $\chi_{ARM}/SIRM$ vs χ_{lf} showing a distinction based on Dhordo chronological division.	101
Figure 7.5	Temporal scale variability in concentration environmental magnetic parameters in Berada core, western India.	105
Figure 7.6	Temporal scale variability in grain size environmental magnetic parameters in Berada core, western India	106
Figure 7.7	Temporal scale variability in mineralogical environmental magnetic parameters in Berada core, western India.	108
Figure 7.8	Scatter plots between a) χ_{ARM} vs χ_{lf} , b) χ_{lf} vs S-ratio and c) $\chi_{ARM}/SIRM$ vs χ_{lf} showing a distinction based on Berada chronological division.	108
Figure 8.1	The microphotographs show morphology of pollens collected from the Dhordo core. 1&2. <i>Rhizophora</i> spp., 3. <i>Sonneratia</i> sp., 4. <i>Typha</i> ., 5. <i>Avecennia</i> ., 6. <i>Casuarina</i> ., 7. <i>Acacia</i> ., 8. <i>Utricularia</i> ., 9. <i>Emblica</i> ., 10. <i>Terminalia</i> ., 11. <i>Crotolaria</i> ., 12. <i>Holoptelea</i> ., 13. <i>Acantheceae</i> ., 14. <i>Tinospora cordifolia</i> ., 15. <i>Poaceae</i> ., 16. <i>Cereal</i> ia., 17. <i>Amaranthaceae</i> ., 18. <i>Brassicaceae</i> ., 19. <i>Artemisia</i> ., 20. <i>Ranunculaceae</i> ., 21&40. <i>Alternanthera</i> ., 22. <i>Cannabis sativa</i> ., 23. <i>Malvaceae</i> ., 24. <i>Asteroidae/Tubuliflorae</i> ., 25. <i>Cichorioideae/Liguliflorae</i> ., 26. <i>Cyperceae</i> ., 27. <i>Zygnema</i> ., 28. <i>Spirogyra</i> ., 29. <i>Pseudoschizaea</i> ., 30. <i>Potamogeton</i> ., 31. Monlete fern spore I., 32. Monlete fern spore II., 33. Trilete fern Spore I., 34. Trilete fern Spore II., 35. <i>Pinns</i> ., 36. <i>Picea</i> ., 37. Dinoflegellate cysts. 38. Foraminiferal linings. 39. <i>Nigrospora</i> ., 41. Globose misofossil with long protuberances., 42. Croton	121
Figure 8.2	Pollen diagram represented in-depth showing percentages of main pollen taxa of Dhordo core. Pollen zones are based on pollen abundance.	122
Figure 8.3	The microphotographs show morphology of pollens collected from the Berada core. 1. <i>Rhizophora</i> spp., 2&3. <i>Sonneratia</i> sp., 4. <i>Bruguiera</i> sp., 5 <i>Nypa</i> ., 6. <i>Holoptelea</i> ., 7. <i>Casuarina</i> ., 8. <i>Syzygium</i> ., 9. <i>Acanthaceae</i> ., 10. <i>Poaceae</i> ., 11. <i>Cereal</i> ia., 12. <i>Amaranthaceae</i> ., 13. <i>Brassicaceae</i> ., 14. <i>Alternanthera</i> ., 15. <i>Cannabis sativa</i> ., 16. <i>Malvaceae</i> ., 17. <i>Asteroidae/Tubuliflorae</i> .,	129

18. *Xanthium.*, 19. *Cyperceae.*, 20. *Polygonum.*, 21. *Zygnema.*,
 22. *Spirogyra.*, 23. *Pseudoschizaea.*, 24. *Potamogeton.*, 25.
 Monlete fern spore., 26. Trilete fern Spore., 27. *Pinns.*, 28.
Cedrus., 29. *Ephedra.*, 30. Dinoflegellate cysts 31. *Nigrospora.*,
 32. Foraminiferal linings.

- | | | |
|-------------|---|-----|
| Figure 8.4 | Pollen diagram represented in-depth showing percentages of main pollen taxa of Berada core. Pollen zones are based on pollen abundance. | 133 |
| Figure 9.1 | Accumulation curves for the Dhordo and Berada cores and the respected depositional units of both cores. The sea-level curves for the last 11 kyr are illustrated based on Fairbanks et al., 1989; Hashimi et al., 1995; Camoine et al., 1997; Makawana et al., 2018). | 140 |
| Figure 9.2a | Graphical representation of environmental magnetic parameters and grain size variation in Dhordo core during Holocene period. | 145 |
| Figure 9.2b | Graphical representation of environmental magnetic parameters and grain size variation in Berada core during Holocene period. | 145 |

LIST OF TABLES

Table no.	Description	Page no.
Table 2.1	Litho-Stratigraphy of Mesozoic rock of Kachchh Basin, (after Biswas, 1977)	12
Table 2.2	Stratigraphy of Tertiary sediments of Kachchh basin (after Biswas, 1992)	15
Table 4.1	AMS Radiocarbon ages (organic carbon) from both the cores (Dhordo and Berada) processed at the IUAC (Inter University Accelerator Centre, New Delhi).	39
Table 4.2	Previously available AMS radiocarbon ages from both the cores (Dhordo and Berada) published by Khonde et al. (2017a, b). The samples were processed at The University of Arizona, USA.	40
Table 4.3	Radiocarbon ages from both the cores (Dhordo and Berada) processed at The University of Arizona, USA (after Khonde et al., 2017a, b). Sample material used were shell of macrofossils acquired from both the cores.	44
Table 4.4	Radiocarbon ages from both the cores (Dhordo and Berada) obtained in the present study and processed at WHOI (Woods Hole Oceanography Institute, USA) and PRL (Physical Research Laboratory, Ahmedabad). Samples used were foraminifera and broken shells collected at the various depths from the cores.	44
Table 4.5	AMS radiocarbon ages of Dhordo core used for reconstructing the age model. [#-present study; *-after Khonde et al., (2017a b)].	45
Table 4.6	Radiocarbon age of Berada core used for the age model in this study. Note that the ages marked with * are previously published dates by Khonde et al., (2017a b). Ages with # marks are generated during the present study.	48
Table 6.1	The grain size distribution of the Dhordo core.	78
Table 6.2	Textural distribution of the sediments based on Blot and Pye, (2012).	82
Table 6.3	The grain size distribution of the Berada core.	88

Table 6.4	Textural distribution of the sediments based on Blot and Pye, (2012).	91
Table 8.1	Showing percentage of pollen grain abundance in the samples analysed for Dhordo core.	114
Table 8.2	Showing percentage of pollen grain abundance in the samples analysed for Berada core.	125