

List of Figures

Sr. No	Name of the Figure	Pg. No
1	Figure 1.1 pathway by which fats stores in the adipose tissue.	5
2	Figure 1.2 Fundamentals of energy balance and energy regulation	7
3	Figure 2. 1 Achyranthus Aspera Seed	28
4	Figure 2 Murraya Koenigii(L) Leaves	29
5	Figure 2.3 Commiphora mukul gum	30
6	Figure 2.4 Garcinia Indica Fruit	32
7	Figure 4.1 Result of responses	63
8	Figure 4.2 ANOVA for Response surface Linear Model	64
9	Figure 4.3 ANOVA study Results for Disintegration time	65
10	Figure 4.4 Desirability plot contour for Disintegration time.	65
11	Figure 4.5 Desirability plot (3D surface) for Disintegration time	66
12	Figure 4.6 ANOVA for Response surface Linear Model	66
13	Figure 4.7 Desirability plot (Contour plot) for Hardness	67
14	Figure 4.8 Desirability plot (3D surface) for Hardness	68
15	Figure 4.9 Desirability plot (Contour plot) for optimized batch	70
16	Figure 10 Desirability plot (3D surface) for optimized batch	70
17	Figure 11 Overlay plot for design space	71
18	Figure 4.12 Structures of (A) Natural Products NPs and (B) Drug molecules collected from ZINC database	86
19	Figure 5.1 Calibration graph of Gallic acid	96
20	Figure 5.2 Calibration plot of gallic acid for tannin content	97
21	Figure 5.3 Calibration plot of Quercetin	98
22	Figure 5.4 Overlay spectra of Overlay spectra for both markers GA and OA	101
23	Figure 5.5 HPLC Chromatogram of Simultaneous estimation of Gallic acid and Oleanolic acid	101
24	Figure 5.6 Overlay HPLC Chromatogram for different linearity concentration for both	102
25	Figure 5.7 Overlay HPLC Chromatogram for different linearity concentration for Galic Acid	103
26	Figure 5.8 Overlay HPLC Chromatogram for different linearity concentration for Oleanolic acid	103
27	Figure 5.9 Calibration curve between Area of peak GA verses its Concentration	104
28	Figure 5.10 Calibration curve between Area of peak OA verses Concentration	104
29	Figure 5.11 TLC plate at UV scan	109
30	Figure 5.12 TLC Plate after spraying with reagent methanolic sulphuric acid	109
31	Figure 5.13 HPTLC Chromatogram of Gallic acid and E-Guggulosterone at 270 nm	109

32	Figure 5.14HPTLC Chromatogram of Oleanolic acid after spraying with reagent scanning at 368 nm	110
33	Figure 5.15HPTLC graph for linearity of Gallic acid and E-Guggulosterone scanned at 270 nm	111
34	Figure 5.16Overlay HPTLC chromatogram for Linearity of Gallic acid and E-Guggulosterone scanned at 270 nm	111
35	Figure 5.17HPTLC chromatogram for Linearity of Oleanolic acid scanned at 368 nm.	112
36	Figure 5.18Overlay HPTLC chromatogram for linearity of Oleanolic acid scanned at 368 nm	112
37	Figure 5.19Calibration curve of Gallic acid.	113
38	Figure 5.20Calibration Curve of E-Guggulsterone	113
39	Figure 5.21Calibration Curve of Oleanolic acid.	114
40	Figure 5.22HPTLC chromatogram of laboratory prepared polyherbal tablet	117
41	Figure 5.23Docking interactions of (A) Oleanolic acid and (B) Acarbose in the active sites of alpha-amylase (PDB id: 6GXV)	120
42	Figure 5.24Docking interactions of (A) Garcinia (B) Oleanolic acid and (C) Orlistat in the active sites of human gastric lipase (PDB id: 1HLG)	123
43	Figure 5.25Docking interactions of (A) Guggulsterone (B) Oleanolic acid (C) Mahanine (D) Quercetin and (E) Clofibrate in the active sites of human PPAR alpha (PDB id: 3VI8)	126
44	Figure 5.26. Docking interactions of (A) Mahanine (B) Oleanolic acid (C) Rimonabant in the active sites of human obesity protein, leptin (PDB id: 1AX8)	128
45	Figure 5.27Docking interactions of (A) Mahanine (B) Oleanolic acid (C) Rimonabant in the active sites of leptin receptor (PDB id: 3V6O)	128
46	Figure 5.28Docking interactions of (A) Quercetin (B) Oleanolic acid (C) Rimonabant in the active sites of cannabinoid receptor type 1 (PDB id: 7V3Z)	131
47	Figure 5.29Docking interactions of (A) Quercetin (B) Oleanolic acid (C) Atorvastatin in the active sites of human HMG-CoA reductase (PDB id: 1DQA)	133