

LIST OF FIGURES

<u>Fig. No.</u>	<u>Description</u>	<u>Page</u>
3-1	Fusarium wilt	38
3-2	Experimental protocol 1: Varietal screening of <u>N. tabacum</u> against <u>F. oxysporum</u> (wilt pathogen)	43
3-3	Camera-lucid drawings of <u>Fusarium oxysporum</u> Schlechtendahl f. sp. <u>nicotianae</u> (J. Johnson) Snyder & Hansen.	49
3-4	Tobacco inoculation with <u>F. oxysporum</u> .	51
3-5	Wilt symptoms on tobacco	52
3-6	Anatomical changes in the wilted plants due to fungus and CF inoculation.	55
3-7	Varietal screening of tobacco to the wilt pathogen <u>F. oxysporum</u>	58
3-8	Varietal screening of tobacco to the CF of <u>F. oxysporum</u>	61
3-9	Graphical representation of correlation.	64
4-1	Low molecular weight toxins of <u>Fusarium</u> .	75
4-2	Experimental protocol 2: effect of <u>Fusarium</u> CF on the growth of whole plants, excised leaf disks, anthers and cells of tobacco.	78
4-3	Growth inhibition of seedlings, excised leaf disks, anthers and cells of tobacco due to CF of <u>F. oxysporum</u> .	87
4-4	Effect of CF levels on chlorophyll-contents of tobacco leaf disks at different hours of incubation.	89
4-5	Effect of CF levels on dry weight of leaf disks, chlorophyll pigments, pH change of the culture medium and the electrolyte leachate in the bathing solution.	90
4-6	Wilt inducing activity of different fractions of CF on tobacco plants.	92

<u>Fig. No.</u>	<u>Description</u>	<u>Page</u>
4-7	Fusaric acid and wilt induction on tobacco	94
4-8	Fractions obtained from Sephadex G100 by passing concentrated aqueous CF =absorption and bioassay (chlorophyll estimation).	95
4-9	Spectrophotometric scanning for absorption of the column eluted culture fractions (No.11 to 14)	96
4-10	UV absorption of fractions obtained from Sephadex G25 by passing concentrated aqueous CF.	98
4-11	Effect of sucrose conc. in culture medium on the synthesis of fungal toxin(s).	100
4-12	CF from various sucrose conc. media and its wilt induction on tobacco plants.	101
4-13	Effect of purified Fusarium toxin on the growth of excised leaf disks and seedling of tobacco.	105
5-1	Experimental protocol 3: Cell line selection of tobacco resistant to <u>F. oxysporum</u> .	117
5-2	Dose sensitivity of CF of <u>F.oxysporum</u> on the efficiency of tobacco.	122
5-3	In vitro morphogenesis of tobacco in presence of selection pressure.	124
5-4	In vitro culture of excised anthers of tobacco, undergoing androgenesis after 4 weeks.	127
5-5	Stimulatory effect of activated charcoal (upto 1%) on androgenesis in the excised anthers.	129
5-6	Leaf-disk bioassay on LD50 selected and regenerated tobacco plants for disease resistance	133
5-7	Leaf-disk bioassay of haploids for disease resistance.	136
5-8	A) LD50 selected tobacco plants transplanted to pots containing garden soil mixture in greenhouse. B) Infection of LD50 selected tobacco plant in lethal inoculum density of <u>Fusarium</u> .	139

<u>Fig. No.</u>	<u>Description</u>	<u>Page</u>
5-9	Control study to see whether the resistancy obtained is due to somaclonal variation or not.	140
5-10	A) Haploids transplanted to pot B) Anther culture in the presence of fungal CF.	143
5-11	Effect of CF conc. on anther culture (culture medium added with activated charcoal).	144