

B I B L I O G R A P H Y

BIBLIOGRAPHY

- Allsopp, A. (1963a). Morphogenesis in Marsilea. J. Linn. Soc. (Bot.) Lond., 58:417-427.
- Allsopp, A. (1964). The metabolic status and morphogenesis. Phytomorphology, 14:1-27.
- Atkinson, L.R. and Stokey, A.G. (1964). Comparative morphology of the gametophyte of the homosporous ferns. Phytomorphology, 14:51-70.
- Baur, L. (1963). On the physiology of sporogonium differentiation in mosses. J. Linn. Soc. (Bot.) Lond., 58:343-351.
- Bell, P.R. (1958). Variations in germination-rate and development of fern spores in culture. Ann. Bot., 22:503-511.
- Bell, P.R. (1959). The experimental investigation of the Pteridophyte life-cycle. J. Linn. Soc. (Bot.) Lond., 56:188-203.
- Bell, P.R. (1961). Failure of nucleotides to diffuse freely into the embryo of Pteridium aquilinum. Nature, 191:91-92.
- Bell, P.R. (1970). The archegoniate revolution. Sc. Progress, 58:27-45.
- Bell, P.R. and Muhlethaler, K. (1962b). A membrane peculiar to the egg in the gametophyte of Pteridium aquilinum. Nature, 195:198.

- Bell, P.R. and Richards, B.M. (1958). Induced apospory in Polypodiaceous ferns. *Nature*, 182:1748-1749.
- Bergmann, L. (1960). Growth and division of single cells of higher plants in vitro. *J. Gen. Physiol.*, 43:841-851.
- Bopp, M. (1968). Control of differentiation in fern-allies and Bryophytes. *Ann. Rev. Pl. Physiol.*, 19:361-380.
- Bristow, J.M. (1962). The controlled in vitro, differentiation of callus derived from a fern, Pteris cretica L., into gametophytic or sporophytic tissues. *Dev. Biol.*, 4:361-375.
- Brown, E.D.W. (1923). Apogamy in Phegopteris polypodioides. *Bull. Torrey. Bot. Club*, 50:17-34.
- Butcher, D.N. and Street, H.E. (1964). Excised root culture. *Bot. Rev.*, 30:513-586.
- Carlson, P.S., Smith, H.H. and Dearing, R.D. (1972). Parasexual interspecific hybridization. *Proc. Nat. Acad. Sci. U.S.A.*, 69:2292-2294.
- Caponetti, J.D. (1972). Morphogenetic studies on excised leaves of Osmunda cinnamomea: Morphological and histological effects of sucrose in sterile nutrient culture. *Bot. Gaz.*, 133:421-435.
- Caponetti, J.D. and Steeves, T.A. (1970). Morphogenetic studies on excised leaves of Osmunda cinnamomea: histological studies of leaf development in sterile nutrient culture. *Can. J. Bot.*, 48:1005-1016.

- Clutter, M.E. and Sussex, I.M. (1965). Meiosis and sporogenesis in excised fern leaves grown in sterile culture. Bot. Gaz., 126:72-78.
- Conway, E. (1949). The autecology of bracken (Pteridium aquilinum (L.) Kunze): the germination of the spore, and the development of the prothallus and the young sporophyte. Proc. Roy. Soc. Edinburgh, B.63: 325-342.
- DeMaggio, A.E. (1963). Morphogenetic factors influencing the development of fern embryos. J. Linn. Soc.(Bot.) Lond., 58:361-376.
- DeMaggio, A.E. (1964). "Organisation in a gametophyte callus of Lycopodium and its morphogenetic implications". Proc. Nat. Acad. Sci. U.S.A., 52:854-859.
- DeMaggio, A.E. (1966). Biochemical and morphological differentiation in a haploid tissue culture of Lycopodium. Am. J. Bot. 53:613.
- DeMaggio, A.E. (1968). Meiosis in vitro sporogenesis in cultured fern plants. Am. J. Bot., 55:915-922.
- DeMaggio, A.E. and Wetmore, R.H. (1961). Morphogenetic studies in the fern Todea barbara. III. Experimental embryology. Am. J. Bot., 48:551-565.
- Erdtman, G. (1954). "An introduction to Pollen analysis". Waltham, Mass., U.S.A. Chronica Botanica Company Stockholm:Almqvist and Wiksell.

- Freeberg, J.A. (1957). The apogamous development of sporelings of Lycopodium cernuum, L. Complantum var. flabelliforme Fernald and L. selago in vitro. Phytomorphology, 7:217-229.
- Fridborg, G. (1971). Growth and organogenesis in Tissue cultures of Allium cepa. var. proliferum. Physiol. Plant., 25:436-440.
- Gautheret, R.J. (1939). Sur la possibilite' de realiser la culture indefinite des tissus de tubercules de carotte. C.R. Acad. Sci., Paris, 208:118-120.
- Gautheret, R.J. (1959). "La Culture des tissus ve ge taux, techniques et realisations". Masson, Paris.
- * Haberlandt, G. (1902). Kulturversuche mit isolierten Pflanzenzellen. S.B. Akad. Wiss. Wien, Math-Naturw. Kl.III, 69-92.
- Harvey, W.H. and Caponetti, J.D. (1972). In vitro studies on the induction of sporogenous tissue on leaves of cinnamon fern I: Environmental factors. Can. J. Bot., 50:2673-2682.
- * Heilbronn, A. (1910). Apogamie, Bastardierung und Erglichkeitsverhatnisse bei einigen Farne. Flora, 101:1-42.
- * Heim, C. (1896). Untersuchugen uber Farnprothallien. Flora, 82:329-373.
- Heslop-Harrison, J. (1967). Differentiation. Ann. Rev. Pl. Physiol., 18:325-348.

- Hevly, R.H. (1963). Adaptations to Cheilanthoid ferns to desert environment. Jour. Arizona Acad. Sci., 2:164-175.
- Ito, M. (1962). Studies on the differentiation of fern gametophytes I. Regeneration of single cells isolated from cordate gametophytes of Pteris vittata. Bot. Mag. Tokyo, 75:19-27.
- Jayasekera, R.D.E. and Bell, P.R. (1959). The effect of various experimental treatments on the development of the embryo of the fern Thelypteris palustris. Planta, 54: 1-14.
- Jensen, W.A. (1962). Botanical Histochemistry Principles and Practice. W.H. Freeman and Company, San Francisco and London, United States of America.
- Kato, Y. (1963b). Physiological and morphogenetic studies of fern gametophytes in aseptic culture. I. Callus tissue from dark-cultured Pteris vittata L. Bot. Gaz., 124:413-416.
- Kato, Y. (1964). Physiological and morphogenetic studies of fern gametophytes in aseptic culture. III. Growth and differentiation of single cells isolated from callus tissues of Pteris vittata. Cytologia, 29:79-85.
- Kato, Y. (1965a). Physiological and morphogenetic studies of fern gametophytes and sporophytes in aseptic culture. IV. Controlled differentiation in leaf callus tissues. Cytologia, 30:67-74.

- Kato, Y. (1969). Physiological and morphogenetic studies of fern gametophytes and sporophytes in aseptic culture. VII. Experimental modifications of dimensional growth in gametophytes of Pteris vittata. *Phytomorphology*, 19:114-121.
- Kato, Y. (1970). Physiological and morphogenetic studies of fern gametophytes and sporophytes in aseptic culture. XII. Sporophyte formation in dark-cultured gametophyte of Pteris vittata L. *Bot. Gaz.*, 131: 203-209.
- Kotte, W. (1922). Kulturversuche mit isoliertion wurzelspitzen. *Beitr. allg. Bot.*, 2:413-434.
- Laetsch, W.M. and Briggs, W.R. (1961). Kinetin modification of sporeling ontogeny in Marsilea vestita. *Am. J. Bot.*, 48: 369-377.
- Lal, M. (1963). Experimental Induction of Apogamy and the Control of Differentiation in Gametophytic Callus of the Moss Phycomitrium coorgense Broth. In 'Plant Tissue and Organ Culture' - A Symposium. International Society of Plant Morphologists, Delhi (P. Maheshwari and N.S. Ranga Swamy eds.).
- *Lang, W.H. (1898). On apogamy and development of sporangia on fern prothalli. *Phil. Trans. R. Soc.*, 110B:187-236.
- *Lang, W.H. (1909). In 'Discussion on "Alternation of Generations" at the Linnean Society'. *New Phytol.*, 8:104-116.
- Lang, W.H. (1929). On a variety of Scolopendrium that bears sporangia on the prothallus. *Ann. Bot.*, 43:355-374.

- Loyal, D.S. (1963). Some evolutionary trends in family. Thelypteridaceae with particular reference to Himalayan species. The memiors of the Indian Bot. Soc., 4: 22-29.
- Manton, I. (1950). Problems of cytology and evolution in the Pteridophyta. University Press, Cambriadge.
- Mehra, P.N. (1973). Breakage of barriers between sporophytic and gametophytic generations of ferns in vitro. Advance notes on Symposia and Discussion. Sixtieth (Diamond Jubilee) Session of the Indian Science Congress Chandigarh, India.
- Mehra, P.N. and Palta, H.K. (1971). In vitro controlled differentiation of the root callus of Cyclosorus dentatus. Phytomorphology, 21:367-375.
- Mehra, P.N. and Sulklyan, D.S. (1969). In vitro studies on apogamy, apospory and controlled differentiation of rhizome segments of the fern, Ampelopteris prolifera (Retz.) Copel. J. Linn. Soc. (Bot.) Lond., 62:431-443.
- Mehta, A.R. (1973). Suspension Cultures. Advance notes on Symposia and Discussion. Sixtieth (Diamond Jubilee) Session of the Indian Science Congress, Chandigarh, India.
- Miller, J.H. (1968). Fern gametophyte as experimental material. Bot. Rev., 34:361-440.

- Miller, J.H. and Miller, P.M. (1964). Effects of auxin on cell enlargement in fern gametophytes: developmental stage of the cell and response to auxin. *Am. J. Bot.*, 51:431-436.
- Miller, C.O., Skoog, F., Von Saltza, M., and Strong, F. (1955). Kinetin, a cell division factor from deoxyribonucleic acid. *J. Am. Chem. Soc.*, 77:566-576.
- Mohr, H. (1956a). Die Beeinflussung der Keimung von Farnsporen durch Licht und andere Faktoren. *Planta*, 46:534-551.
- Mohr, H. (1962). The influence of visible radiation on the germination of archegoniate spores and the growth of the fern protonema. *J. Linn. Soc. (Bot.) Lond.*, 58:287-296.
- * Morel, G. (1956a). Nouvelles me'thods de culture de tissue. *Rev. gen. botan.* 63:314-324.
- Morel, G. (1963). Leaf regeneration in Adiantum pedatum. *J. Linn. Soc. (Bot.) Lond.*, 58:381-384.
- Morel, G. and Wetmore, R.H. (1951). Fern callus tissue culture. *Am. J. Bot.*, 38:140-143.
- Munore, M.H. and Bell, P.R. (1970). The fine structure of fern root cells showing apospory. *Dev. Biol.*, 23:550-562.
- Munore, M.H. and Sussex, I.M. (1969). Gametophyte formation in bracken fern root cultures. *Can. J. Bot.*, 47:617-621.

- Murashige, T. and Skoog, F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.*, 15:473-497.
- Naf, U. (1962). Developmental physiology of lower archegoniates. *Ann. Rev. Pl. Physiol.*, 13:507-532.
- Nayar, B.K. (1962). Studies in Pteridaceae-V: Contributions to the morphology of some species of the maiden hair ferns. *J. Linn. Soc. (Bot.) Lond.*, 58:185-199.
- Nayar, B.K. and Kaur, S. (1969). Types of prothallial development in homosporous ferns. *Phytomorphology*, 19:179-188.
- Nayar, B.K. and Kaur, S. (1971). Gametophytes of homosporous ferns. *Bot. Rev.*, 37:295-396.
- Nitsch, J.P. (1951). Growth and development in vitro of excised ovaries. *Am. J. Bot.* 38:566-577.
- Nitsch, J.P., and Nitsch, C. (1969). Haploid plants from pollen grains. *Science*, 163:85-87.
- Nobecourt, P. (1939). Sur les radicules naissant des cultures de tissus du tubercule de carotte. *C.R. Soc. Biol. Paris*, 130:1270-1271.
- Osborne, D. (1962). Effect of kinetin on protein and nucleic acid metabolism in Xanthium leaves during senescence. *Plant Physiol.*, 37:595-602.
- Partanen, C.R. (1972). Comparison of gametophytic callus and tumor tissues of Pteridium aquilinum. *Bot. Gaz.*, 133:287-292.

- Partanen, J.N. and Partanen, C.R. (1963). Observations on the culture of roots of the bracken fern. Can. J. Bot., 41:1657-1661.
- Peterson, R.L. (1967). Callus induction in Ophioglossum petiolatum Hook. Can. J. Bot., 45:2225-2227.
- Rashid, A. (1971). Reproductive biology of the ferns: Lygodium and Ceratopteris. In International Symposium Morphogenesis in Plant Cell, tissue and organ cultures. U.G.C. Centre of Advanced Study in Botany, Dept. of Botany, University of Delhi, Delhi, India.
- * Schleiden, M.J. (1838). Beitrage zur Phytogenesis. Arch. Anat., Physiol. U. Wiss. Med. (J. Miller ed.).
- Schwabe, W.W. (1951). Physiological studies in plant nutrition. XVI. The mineral nutrition of bracken. Part I. Prothallial culture and the effects of phosphorus and potassium supply on leaf production in the sporophyte. Ann. Bot., 15:417-446.
- * Schwann, TH. (1839). Mikroskopische Untersuchungen uber die ubereinstimmung in der Struktur und dem Wachstume der Tiere und Pflanzen. Leipzig: W. Engelmann, Nr. 176, Ostwalds Klassiker der exakten Wissenschaften 1910.
- Sinnott, E.W. (1960). Plant morphogenesis. McGraw-Hill, New York.
- Stange, L. (1965). Plant cell differentiation. Ann. Rev. Pl. Physiol., 16:119-140.

- Steebin, G.L. (1950). Variation and evolution in plants. Columbia University Press, New York.
- Steeves, T.A., Sussex, I.M. and Partanen, C.R. (1955). In vitro studies on abnormal growth of prothalli of the bracken fern. Am. J. Bot., 42:232-245.
- Steeves, T.A. and Sussex, I.M. (1957). Studies on the development of excised leaves in sterile culture. Am. J. Bot., 44:665-674.
- Steil, W.N. (1939). Apogamy, apospory and Parthenogenesis in the Pteridophytes. Bot. Rev., 5:433-453.
- Steil, W.N. (1951). Apogamy, apospory and parthenogenesis in Pteridophytes II. Bot. Rev., 17:90-104.
- Steward, F.C. (1958). Growth and organised development of cultured cells. III. Interpretations of the growth from free cells to carrot plant. Am. J. Bot., 45:709-713.
- Steward, F.C. (1968). Growth and organisation in plants. Addison-Wesley Publishing Co., Reading, Mass.
- Steward, F.C., and Shantz, E.M. (1955). The chemical induction of growth in plant tissue cultures. In "The Chemistry and Mode of Action of Plant Growth Substances" (R.L. Wain and F. Wightman, eds.). Butterworth, London.
- Steward, F.C., Mapes, M.O. and Ammirato, P.V. (1969). Growth and Morphogenesis in Tissue and free cell culture. In "Analysis of Growth: The responses of cells and Tissues in culture" (F.C. Steward ed.), Academic Press, New York and London.

- Stokey, A.G. (1951). The contribution of the gametophyte to the classification of the homosporous ferns. *Phytomorphology*, 1:39-58.
- Street, H.E. (1966). The nutrition and metabolism of plant tissue and organ cultures. In "Cells and tissues in culture". (E.N. Willmer ed.), Academic Press, London.
- Street, H.E. (1969). Growth in organised and unorganised systems Knowledge gained by Culture of Organs and Tissue Explants. In "Analysis of Growth: The responses of cells and tissues in culture" (F.C. Steward ed.). Academic Press, New York and London.
- Street, H.E. and Henshaw, G.G. (1966). Introduction and Methods Employed in Plant Tissue Culture. In "Cells and tissues in culture" (E.N. Willmer ed.). Academic Press, London.
- Sussex, I.M. (1958). A morphological and experimental study of leaf development in Leptopteris hymenophylloides (A, Rich.) Presl. *Phytomorphology*, 8:96-107.
- Sussex, I.M. and Steeves, T.A. (1958). Experiments on the control of fertility of fern leaves in sterile culture. *Bot. Gaz.*, 119:203-208.
- Takahashi, C. (1962). Cytological studies on induced apospory in ferns. *Cytologia*, 27:79-96.
- Takebe, I., Labib, G. and Melchers, G. (1971). Regeneration of whole plants from isolated mesophyll protoplasts of tobacco. *Naturwissenschaften*, 58:318-320.

- Treanor, L.L. and Whittier, D.P. (1969). The effect of mineral nutrition on apogamy in Pteridium. Can. J. Bot., 47:733-777.
- Vajranabhaiah, S.N. (1969). The changes in nucleic acids in relation to auxins, kinetin and sugar during growth in cultures of Cucumis melo L. and Datura metel L. Ph.D. Thesis, M.S. University of Baroda.
- Ward, M. (1954a). Fertilization in Phlebodium aureum J. sm. Phytomorphology, 4:1-17.
- Ward, M. (1960). Callus tissue from the mosses Polytrichum and Atrichum. Science, 132:1401-1402.
- Ward, M. (1963). Developmental pattern of adventitious sporophytes in the fern Phlebodium aureum. J. Linn. Soc. (Bot.) Lond., 58:377-380.
- Wetmore, R.H. and Rier, J.P. (1963). Experimental induction of vascular tissues in callus of Angiosperms. Am. J. Bot., 50:418-430.
- White, P.R. (1937). Vitamin B₁ in the nutrition of excised tomato roots. Plant Physiol., 12:803-811.
- White, P.R. (1939). Potentially unlimited growth of excised plant callus in an artificial nutrient. Am. J. Bot., 26:59-64.
- White, P.R. (1954). "The cultivation of animal and plant cells". Thames and Hudson, London.
- White, P.R. (1963). The cultivation of Animal and Plant Cells. 2nd Ed., Ronald Press Co., New York.

- White, R.A. (1971). Experimental and developmental studies of fern sporophyte. *Bot. Rev.*, 37:509-540.
- Whittier, D.P. (1962). The origin and development of apogamous structures in the gametophyte of Pteridium in sterile culture. *Phytomorphology*, 12:10-20.
- Whittier, D.P. (1964a). The influence of cultural conditions on the induction of apogamy in Pteridium gametophytes. *Am. J. Bot.*, 51:703-736.
- Whittier, D.P. (1965). Obligate apogamy in Cheilanthes tomentosa and Cheilanthes alabamensis. *Bot. Gaz.*, 126:275-281.
- Whittier, D.P. (1966b). Induced apogamy in diploid gametophytes of Pteridium. *Can. J. Bot.*, 44:1717-1721.
- Whittier, D.P. (1971). The value of ferns in an understanding of the alternation of generations. *Bioscience*, 21:225-227.
- Whittier, D.P. and Steeves, T.A. (1960). The induction of apogamy in bracken fern. *Can. J. Bot.*, 38:925-930.
- Whittier, D.P. and Steeves, T.A. (1962). Further studies on induced apogamy in ferns. *Can. J. Bot.*, 40:1525-1531.