

BIBLIOGRAPHY

BIBLIOGRAPHY

- Alfermann, A.W. and Reinhard, E. (1978). Possibilities and problems in production of natural compounds by cell culture methods. In : Production of Natural Compounds by Cell Culture Methods, (eds. Alfermann, A.W. and Reinhard,E.) BPT. Report. Gesellschaft für strahlen und Umweltforschung, Munich,3-15
- Ammirato, P.V. (1987) organization events during somatic embryogenesis. In : Plant Tissue and Cell Culture (eds. Green, C.E, Somres, D.A.; Hackelt and Biesboer, D.D.) Alan R Liss, Inc, New York.
- Anderson, L.A.; Phillipson, J.D. and Roberts, M.F. (1986). Aspects of alkaloid production by plant cell cultures. In : Secondary Metabolism in Plant Cell Cultures (eds. Morris, P.; Scragg, A.H.; Stafford, A. and Fowler, M.W.). Cambridge University Press, Great Britain.
- Anderson, W.C. (1978). Rooting of tissue cultured *Rhododendrons*. Proc.Int.Plant Pro.Soc., 28 : 135-139.
- Attfield, E.M. and Evans, P.K. (1991). Developmental pattern of root and shoot organogenesis in cultured leaf explants of Nicotina tobacum cv. Xanthi nc. J.Experimental Botany, 42, 51-57.

Bahadur, B;Reddy, K.R.K. and Krishna Murthy, K.V. (1991). Experimental Xylogenesis in castor bean callus cultures. *Phytomorphology*, **41**, 299-303.

Bapat, V.A. and Rao, P.S. (1988) Somatic seeds of Sandalwood (*Santalum album* L.) and Mulberry (*Morus indica* L.) In : Proceedings of National Seminar on advances in seed science and technology (eds. Shokara Shetty, H. and Prakash, H.S.). 372

Basu, N.K. (1957). The chemical constitution of achyranthine. *J. and Proc. Inst. Chem.* **29**, 73-76.

Basu, N.K. and Sharma, S.N. (1947). Investigation on Indian medicinal plant. *Qr. J.Pharm and Pharma.* **20**, 38-42.

Benjamin, B.D.; Heble, M.R. and Chadha, M.S. (1979). Alkaloid synthesis in tissue culture and regenerated plants of *Tylophora indica* Merr. *Z. Pflanzenphysiol*, **92**, 74-84.

Bhalla, T.N.; Gupta, M.B. and Bhargava, K.P. (1971). Anti-inflammatory and biochemical study of *Boerhaavia diffusa* *J.Res.Ind.Med.* **6**, 11-15

Bhattacharya, S. and Gupta, S. (1987). Growth and regenerating potentiality of roots of *Nigella sativa* L. in long term culture. *Phytomorphology* , **37**, 303-306.

Bhojwani, S.S. and Johri, B.M. (1970). Cytokinin induced shoot bud

differentiation in mature endosperm of *Scurrula pulverulenta*.
Z .Pflanzenphysiol., 63,269-275.

Bhojwani, S.S. and Rajdan, M.K. (1983). Plant Tissue Culture : Theory and Practice. Elsevier, Amsterdam.

Blakely, L.M.; Roadway, S.J.; Hollen, L.B. and Croker, S.G.(1972). Control and Kinetics branch root formation in cultured root segments of *Haplopappus ravenii*. Plant Physiology, 50, 35-42.

Bohm, H. (1980). The formation of secondary metabolites in plant tissue and cell cultures. Int. Rev. Cytol. Supp. 11B, 183-208.

Boll, W.G. (1965). The growth curve and growth correlations of excised tomato roots in sterile culture. Can J. Bot., 43, 287-304.

Bonnett, H.T. and Torrey, J.G. (1965). Chemical control of organ formation in root segments of *Convolvulus* cultured in *in vitro*. Plant Physiology, 40, 1228-1236.

Borgman, C.A. and Mungae, K.W. (1986). Factors affecting the establishment and maintenance of 'Titan' red raspberry root organ cultures. Plant Cell Tissue and Organ Culture, 6, 127-137.

Butcher, D.N. (1977). Secondary products in tissue culture. In : Plant Cell Tissue and Organ Culture, (eds. Reinert,J and Bajaj, Y.P.S.). Berlin, Heidelberg and New York : Springer Verlag, 668-693.

Butcher, D.N. and Street, H.E. (1960). The effects of gibberellins on the growth of excised tomato roots. *J.Expt.Bot.*, 11, 206-216.

Butcher, D.N. and Street, H.E. (1964). Excised root cultures. *Bot.Rev.*, 30, 513-586.

Caplin, S.M. and Steward, F.C. (1948). Effect of coconut milk on the growth of explants from carrot root. *Science*, **108**, 655-657.

Carlson, P.S. (1973). The use of protoplast for genetic research. *Proc. Natl. Acad. Sci. U.S.A.*, **70**, 598-602.

Chakraborti, K.K. and Handa, S.S. (1989). Antidiheptotoxic investigations on *Boerhaavia repanda* Willd. *Indian Drugs*, **27**, 19-24.

Chopra, R.N., Chopra, I.C., Handa, K.L. and Kapur, L.D. (1982). Indigenous Drugs of India, Academic Publisher, New Delhi.

Chowdhury, A.R. and Chaturvedi, H.C. (1979). Cholesterol and biosynthesis of diosgenin by tuber callus of *Dioscorea deltoidea*. *Current Science*, **49**, 273-278.

Christianson, M.L. and Warnick, D.A., (1984). Phenocritical times in the process of *in vitro* shoot organogenesis. *Developmental Biology*, **101**, 382-390.

Coleman, W.K. and Greyson, R.T. (1976). Root regeneration from leaf

cuttings of *Lycopersicon esculentum* Mill. Applications on the leaf plastochnor index and responses to exogenous Gibberellic acid. J. Exp. Bot., 27, 1339-1351.

Collins, G.B. and Sunderland, N. (1974). Pollen derived haploids of *Nicotina knightiana*, *N.raimondii* and *N.alternuala*, J.Exp. Bot. 25, 1030-1039.

Dodds, J.H. and Roberts, L.W. (1985). Root cultures, In : Experiments In Plant Tissue Culture. Cambridge University Press, Cambridge, 94-103.

Dormer, K.J. and Street, H.E. (1949). The carbohydrate nutrition of tomato roots. Ann. Bot. 13, 199-217.

Dougall, D.K. (1979). Plant Cell and Tissue Culture - Principles and Applications. In : "Proceedings of the Fourth Annual Biological Sciences, Colloquium.". Ohio State University Press, 727-743.

Eapen, S.; Rangan, T.S.; Chadha, M.S. and Heble, M.R. (1978). Morphogenetic and biosynthetic studies in tissue cultures of *Atropa belladonna* L. Plant Science Letter, 13, 83-89.

Evans, D.A.; Sharp, W.R.; Ammirato, P.V. and Yamada, Y. (1983). In : Hand Book of Plant Culture, Techniques for Propagation and Breeding, Macmillan Publishing Co. New York.

Fadia, V.P. and Mehta, A.R. (1973). Tissue culture studies on cucurbits :
The effect of NAA, sucrose and Kinetin on tracheal differentiation in
Cucumis tisues and cultured *in vitro*. *Phytomorphology*, **23**, 212-215.

Fairbrain, J.W.; and Wassel, G. (1964). *Phytochemistry* **3**,253.

Flores, H.E.; Dai, Y.R.; Cuello, J.L.; Mendoza, I.E.M. and Vargas, V.
M.L. (1993). Green roots : Photosynthesis and Photoautotrophy in an
underground plant organ. *Plant Physiology*, **101**, 363-371.

Fosket, D.E. and Torrey, J.G. (1969). Hormonal control of cell
proliferation and xylem differentiation in cultured tissue of *Glycine
max* var. Biloxi. *Plant Physiology*, **44**, 871-880.

Fowler, M.W. (1983). Commercial applications and economic aspects of
mass plant cell culture. In: *Plant Biotechnology* (eds. Mantell, S.H.
and Smith, H.). Cambridge University Press, 3-37.

Fukuda, H. and Komamine, A. (1980). Establishment of an experimental
system for the study of tracheary element differentiation from single
cells isolated from the mesophyll of *Zinnia elegans*. *Plant Physiol*, **65**,
57-60.

Fulzele, D.P.; Sipahimalani, A.T. and Heble, M.R. (1991). Tissue cultures
of *Artemisia annua* : Organogenesis and artemisinin production.
Phytotherapy Research, **5**, 149-153.

Gamborg, O.L. ; Constable, F. ; Larau, T.A.G.; Miller,R.A. and Steck, W. (1971). The influence of hormones on secondary metabolite formation in plant cell cultured. In : Les cultures de Tissue de Plants, Centre National de la Recherche Scientifique Paris, 335-344.

Gautheret, R.J. (1939). Sur la possibilite de realiser la culture indefinie des tissue de tubercules de carotte C.R.Hebd. Seances Acad. Sc. 208, 118-120.

Gautheret, R.J. (1959). La culture des tissue vegetaux. Masson Publ., 483-490.

Gautheret, R.J. (1966). Factors affecting diffferentiation of plant tissue grown *in vitro*.In : Cell Differentiation and Morphogenesis ; Intern. Lectures, Wageningen. 55-95.

(1984).
George, E.F. and Sherrington, P.D. In : Plant Propagation by Tissue Culture, Eastern Press.

Ghoshal, S. and Bhattacharya, S.K.(1972). Desmodium Alkaloids Part II Chemical and Pharmacological evaluation of *D.gangeticum*. Planta Medica, 22, 434-440.

Goswami, P and Sharma, T.C. (1993). Effect of *Boerhaavia diffusa* Linn. Extract on the activities of enzyme system. J. Res. Aur. Siddha, 13, 48-55.

Gramberg, J.J. (1971). The first stage of the formation of adventitious roots in petioles of *Phaseolus vulgaris*. Proc of the Koninlijke Nederlandse Academice Van Wetenschappen, **74**, 42-45

Gresshoff, P.M. (1978) Phytohormones and thier role in differentiation and growth of plant cells *in vitro*. In : Phytohormones and Related Compounds. A Comprehensive Treatise (eds. Letham D,S.; Higgins.T,J and Goodwin. P,J). Elsevier/North Holland Press; 1-29.

Guha, S.and Maheshwari, S.C. (1964). *In vitro* production of embryos from anther of *Datura*. Nature, **204**,497.

Gupta, R.B.L.; Singh, S. and Dayal, Y. (1962). Effect of punarnava on the visual acuity and refractive errors. Ind. J. Med. Res., **50**, 428-434.

Gupta, S.S. ; Bhagwat, A.W. and Ram, A.K., (1972). Cardiac stimulant activity of the saponin of *Achyranthes aspera* L. Ind. J. Med. Res., **60**, 462-471.

Gurel, E. and Wren, M.J. (1995). *In Vitro* development from leaf explants of sugar beet (*Beta vulgaris* L.) rhizogenesis and the effect of sequential exposure to auxin and cytokinin. Ann. Bot.,**75**, 31-38.

Heble, M.R (1985). Multiple shoot cultures : A viable alternative *in vitro* system for the production of known and new biologically active plant

constituents. In : Primary and Secondary Metabolism of Plant Cell Cultures (eds. Neumann, K.H.; Barz, W. and Reinhard, E.) Springer-verlag, Berlin. 282-289.

Heble, M.R. (1993). High value chemicals by tissue culture. Chem. Indus. Digest 3, 113-118.

Heble , M.R; Narayanswami, S. and Chadha, M.S. (1971). Hormonal control of steroid synthesis in *Solanum xanthocarpum* in tissue culture. Phytochem.,**10**, 2393-2394.

Hiraoka, N. and Tabata, M. (1974). Alkaloid production by plants regenerated from cultured cells of *Datura innoxia*. Phytochemistry, **13**, 1671-1675.

Huber, W. (1967). In 'Titration in Non Aqueous Solvent'. Academic Press, New York, London.

Ikeda, T; Matsumoto, T and Noguchi, M. (1976) Effect of nutritional factors on the formation of ubiquinone by tobacco plant cells in suspension culture. Agri. Bio. Chem., **40**, 1765.

Jacobs, W.P., (1952). The role of auxin in differentiation of xylem around a wound. Am. J. Bot., **39**, 301-309.

Jain, S.C. and Khanna, P. (1974). Quercetin from *Crotalaria Juncea* L. tissue cultures. Indian J. Exp. Biol. **12**, 466.

Jensen, W.A. (1962) Botanical Histochemistry - Principles and Practice,
W.H. Freeman and Co. San Francisco.

Kapoor, V.K. and Singh, H. (1967) Investigation of *Achyranthes aspera* L.
Ind. J. Pharm., 29, 285-288.

Kartnig, T. (1977) Cardiac glycosides in cell cultures of Digitalis. In : Plant
Tissue Culture and its Biotechnological Applications (eds. W, Barz,
Reinhard and Zenk). Berlin, Heidelberg and New York; Springer -
Verlag, 44-51.

Kinnersley, A.M. and Dougall, D.K. (1981). Correlation between nicotine
content of tobacco plants and callus cultures. In Cell Science Centre
Annual report (W.Alton Jones) Lake Placid : W. Alton Jones Cell
Science Centre, 7-8.

Kirtikar, K.R. and Basu, B.D. (1975). Indian Medicinal Plants, 3, Vivek
Vihar, Delhi, 2066.

Kotte, W. (1922). Kalturoersuche mit isolierten warzel spitzen. Beitr. allg.
Bot. 2, 413-434.

Lindsey, K. and Yeoman, M.M. (1983). Relationship between growth rate,
differentiation and alkaloid accumulation in cell cultures. J.Expt. Bot.,
34, 1055-1065.

Mantell, S.H.; Mathews, J.A. and Mckae, R.A. (1985) Principles of Plant

Biotechnology, An Introduction to Genetic Engineering in Plants
Black well Scientific Publication, Oxford, 186-220.

Mantell, S.H.; Pearson, D.W.; Hazell, L.P. and Smit, H. (1983). The effect of initial phosphate and sucrose levels on nicotine accumulation in batch suspension cultures of *Nicotina tobacum* L. Plant Cell Reports, 2, 73-77.

Mantell, S.H. and Smith, M. (1983). Cultural factors that influence secondary metabolite accumulations in plant cell and tissue cultures. In 'Plant Biotechnology' (eds. Mantell, S.H. and Smith, H.). Cambridge University Press, Great Britain.

Miller, C.; Skoog, F; Okumura, F.S.; Von Saltza, M.H. and Strong, F.M. (1955). Structure and Synthesis of Kinetin J. Am. Chem. Soc. 77, 2662 - 2663.

Misawa, M. (1982). Industrial and government research. In 'Plant Tissue Culture as a source of Biochemicals' (ed. Staba. E.J.).CRC Press, Boca Raton, Florida, 167-190.

Misra, T.N; Singh, R.S.; Pandey, H.S. and Prasad, C. (1991). An aliphatic dihydroxyketone from *Achyranthes aspera* Phytochemistry, 30, 2076-2078.

Mitra, G.C. (1968). *In Vitro* growth of excised root of *Rauwolfia serpentina* in continuous culture. Ind. Jr. Exp. Bio., 6, 230-231.

Mitra, G.C. (1975). Studies on the growth of excised roots of *Rauwolfia serpentina* : Effect of certain auxins, "antiauxins", amino acid and Kinetin. In Form Structure and Functions in Plant. (eds. Mohanram, H.Y.; Shah, J.J. and Shah C.K.), Sarita Prakashan, Meerut, India.

Mitra, G.C. and Chaturvedi, H.C. (1972). Embryoids and complete plants from unpollinated ovaries and from ovules of *in vivo* grown emasculated flower buds of *Citrus*. Bull. Torrey Botan. Cl., **99**, 184-189.

Mitra, R. and Jain, S.K. (1994). Medicinal Plants Research in India A Review. In : Advances in Plant Sciences Research (ed. Sohni, K.C.) International book distributors, **1**, 47-73.

Morel, G. (1960). Producing virus free cymbidium. Am. Orchid Soc. Bull., **29**, 495-497.

Mudgal,V. (1975). Studies on medicinal properties of *Convolvulus pluricaulis* and *Boerhaavia diffusa*. Planta Medica, **28**, 62-68.

Murashige, T. (1974). Plant Propagation through tissue cultures. Annual Review of Plant Physiology. **25**, 135-166.

Murashige, T. and Skoog, F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiol. Plant. **15**, 473-497.

Narayanswamy, S. (1977). Regeneration of plants from tissue cultures. In :

'Applied and Fundamental Aspects of Plant Cell, Tissue, and Organ Culture (eds. Reinert, J. and Bajaj, Y.P.S.). Springer-Verlag, New York.

Neogi, N.C.; Garg, R.D. and Rathor, R.S. (1970). Preliminary pharmacological studies on Achyranthine. Ind. J. Pharmacy, 43-46.

Nitsch, J.P. and Nitsch, C. (1969). Haploid plants from pollen grains. Science, 163, 85-87.

Nitsch J,P; Nitsch, C.; Rossini, L.M.E. and Ha, B.D. (1967) The role of adenine in bud differentiation Phytomorphology, 17, 446-453.

Nobecourt, P. (1939). Sur la perennite et l' augmentation de volume des cultures de tissus vegetaux. C.R. Seances Soc. Biol. Ses. Fil. 130, 1270-1271.

Nogel, M. and Reinhard, E. (1975). Das Aetherische ole der Calluskulturen von *Ruta graveolus* L. Die zuzammensetzung des Oeles, Plant Med. 27, 151-158.

Noguchi, M.; Matsumoto, T. ; Hirata, Y. ; Yamamoto, K.; Katsuyama, A.; Kato, A. ; Azechi, S. and Kato, K. (1977). Improvement of growth rate of Plant cell cultures. In : Plant Tissue Culture and its Biotechnological Applications' (eds., Barz, W.; Reinhard, E. and Zenk, M.H.), Springer-Verlag, Berlin.

Padhya, M.A. (1993). *In Vitro* studies on *Boerhaavia diffusa* L. At XVI Annual Meeting of IAPTC, Jaipur.

Pakrashi, A. and Bhattacharya, N. (1977). Abortifacient principle of *Achyranthes aspera* L. Ind. J. Exp. Bio., 15, 856-858.

Parr, A.J. and Hamill, J.D. (1987). Relationships between *Agrobacterium rhizogenesis* transformed hairy roots and intact, uninfected *Nicotina* plants. Phytochemistry, 26, 3241-3245.

Ramawat. K, G (1991) Factors affecting production of alkaloids in plant cell cultures. In : Lectures, Short Term Training Course in Plant Tissue/Cell Cultures for the Production of Phytochemicals and Plant Improvement, DBT Sponsored at CFTRI, Mysore, 30-34.

Ramesh, K. and Padya, M.A. (1988). Isolation of nimbin from *Azadirachta indica* leaves and its callus cultures. Indian Drugs, 25, 526-527.

Ramesh, K. and Padhye. M,A. (1991). Propagation of neem-*Azadirachta indica* A. Juss through *in vitro* culture. In : Proc. National Seminar on Biotechnological Approaches for Upgrade of Agricultural and Horticultural crops, Coimbatore. Agri. Univ. 50-51.

Rao, P.S.; Handro, W. and Harada, H. (1973). Hormonal control of differentiation of shoots, roots and embryos in leaf and stem cultures of *Petunia inflata* and *Petunia hybrida*, Physiol. Plant. 28, 458-463.

Ratra, P.S. and Misra, K.C. (1970). Seasonal variation in chemical composition of *Achyranthes aspera* L. and *A. bidentata* Bl. Indian Forester, **96**, 372-375.

Reinhard, E. and Alfermann, A.W. (1980). Biotransformation by plant cell cultures, In "Advances in Biochemical Engineering" (ed. Fiechler), 16, Plant cell culture, I, 49-84. Springer-verlag, New York.

Rhodes, M.J.C.; Robins, R.J.; Hamill, J. and Parr, A.T. (1986). Potential for the production of biochemicals by plant cell culture, New Zealand, J. Technology, **2**, 59-70.

Robins, R.J.; Walton, N.J.; Hamill, J.D.; Parr, A.J. and Rhodes, M.J.C. (1991). Strategies for the genetic manipulation of alkaloid-producing pathways in plants. *Planta Medica*, **57**, 527-535.

Roddick, J.G. and Butcher, D.N. (1972). Isolation of tomatine from cultured excised roots and callus tissue of tomato. *Phytochemistry*, **11**, 2019-2024.

Roja, P.C.; Sipahimalani, A.T.; Heble, M.R. and Chadha, M.S (1987). Multiple shoot culture of *Rauwolfia serpentina* : growth and alkaloid production. *J. Nat. Prod.*, **50**, 872-875.

Ross, M.K and Thorpe, T.A (1973). Physiological gradients and shoot initiations in tobacco callus cultures. *Plant Cell Physiol.* **14**, 473-480.

Routier, J.B. and Nickell, L.G. (1956). Cultivation of plant tissue, U.S. Patent, 2, 747.

Said, A.G.E. and Murashige, T. (1979). Continuous cultures of tomato and Citron roots *in vitro*. In *vitro*, 15, 593-602.

Sander, H. (1956). *Planta* 47, 374

Schieder, O. (1980). Somatic hybrids between a herbaceous and two tree *Datura* Species. *Z. Pflanzenphysiol*, 98, 119-127.

Seabrook, J.E.A. (1980). Laboratory culture. In : Plant Tissue Culture As a Source of Biochemicals (ed. Staba, E.J.), CRC Press, Boca Ratan, Florida, 2-20.

Seth, R.K.; Khanna, M.; Chaudhary, M.; Singh, S. and Sarin, J.P.S. (1986). Estimations of Punarnavoside; a new antifibrinolytic compound from *Boerhaavia diffusa*, *Indian Drug*, 23, 583-584.

Sharma, K.K. and Bhojwani, S.S. (1990). Histological aspects of *in vitro* root and shoot differentiation from cotyledon explants of *Brassica juncea* (L.) Czern. *Plant Science*, 69, 207-214.

Sharma, V. (1991). Rapid Propagation of *Crataeva nurval* Through Tissue Culture. Ph.D. Thesis, M.S.University, Baroda.

Sharma, V. and Padhya, M.A. (1991). Detection of glucosinolate, glucocapparin in axenic root cultures of *Crataeva nurvala*. In : Proceedings of Sympsoium of Plant Tissue Culture, Shillong (ed. Tandon, T.).

Shio, I. and Ohta, S. (1973). Nicotine production by tobacco callus tissues and effect of plant growth regulators. Agricultural and Biological Chemistry, **37**, 1857-1864.

Shrivastava, N and Padhya, M.A. (1991). *In vitro* studies on excised root cultures of *Boerhaavia diffusa* L. In : Abstract, National Seminar on Recent Advances in Plant Sciences, University of Bombay.

Shrivastava, N. and Padhya, M.A. (1992). *In vitro* excised root cultures of *Boerhaavia diffusa* a source of alkaloid. In : Souvenir, National Seminar on Trends in Formulations, Standardisation and Commerce of Herbal Products, Baroda.

Shrivastava, N. and Padhya, M.A. (1995). Punarnavine profile in the regenerated roots of *Boerhaavia diffusa* L. from leaf segments Current Science, **68**, 653-656.

Shukla, N.K.; Narang, R. ; Nair, N.G.K.; Radhakrishnan, S. and Satyavati, G.V. (1991). Multicentric randomized controlled clinical trial of "Kshaarasootra" in the management of fistula-in-ano. Ind. J. Med. Res. **94**, 177-185.

Singh, A.; Singh, R.G; Singh, R.H.; Mishra, N. and Singh, N. (1991). An experimental evaluation of possible teratogenic potential in *Boerhaavia diffusa* in Albino roots. *Planta Medica*, **57**, 315-316

Singh, R.H. and Udupa, K.N. (1972). Studies on the Indian indigenous drug, Punarnava (*Boerhaavia diffusa*), *J.Res.Ind.Med.* **1**, 13-16.

Solt, M.L.; Dawson, R.F. and Christman, D.R. (1960). Biosynthesis of anabasine and of nicotine by excised root cultures of *Nicotina glauca*. *Plant Physiol*, **35**, 887-894.

Staba, E.J. (1980). Secondary metabolism and biotransformation. In : *Plant Tissue Cultures, As a source of Biochemicals* (ed. Staba, E.J.) CRS Press Inc, Bonca Raton, Florida.

Staba, E.J. (1982). Production of useful compounds from plant tissue cultures. In *Plant Tissue Culture*, (ed. A Fujiwara; Tokyo) Japanese Association for Plant Tissue Culture, 25-30.

Steward, F.C. (1958). Growth and development of cultivated cells. III Interpretations of the growth from free cell to carrot plant. *Am.J. Bot.*, **45** : 709-713.

Street, H.E. and Jones, O.P. (1963). Nutritional problems raised by work with root cultures. In *Plant Tissue and Organ Culture A Sympsonium* (eds : Maheswari, P. and Rangaswami N.S.) The International Society

of Plant Morphologist, Delhi.

Street, H.E. and Winter, A (1963). Studies on hormonal control of the growth of excised roots. In : Plant Tissue and Organ Culture A Symposium, (ed. Maheshwari, P. and Rangaswamy N.S.), . The International Society of Plant Morphologist, Delhi.

Sudarshana, M.S. and Shanthamma, C. (1988). Plant regeneration from inflorescence culture of *Boerhaavia diffusa* L. Current Science, 57, 268-270.

Szweykowska, A. (1974). The role of cytokinins in the control of cell Growth and differentiation in culture, In : Tissue Culture and Plant Science (ed. Street, H.E.). Academic Press, London.

Takebe, I.; Labib, C.; and Melchers, G. (1971). Regeneration of whole plants from isolated mesophyll protoplasts of tobacco. Naturwissenschaften, 58, 318-320.

The Wealth of India (1978), Raw Materials, 12, CSIR, New Delhi.

Thomas, D.R.; Carigie, J.S. and Street, H.E (1963). The carbohydrate nutrition of the excised tomato root. In Plant Tissue and Organ Culture A Sympsonium, (ed. Maheshwari P. and Rangaswamy N.S., The International Society of Plant Morphologist, Delhi.

Thomas, E. and Street, H.E. (1970). Organogenesis in cell suspension

cultures of *Atropa belladonna* L. and *Atropa belladonna* cultivar lutea
DOLL. Ann. Bot. 34, 657-669.

Torrey, J.G. (1954). The role of vitamins and micronutrient elements in the nutrition of the apical meristem of pea roots. Plant Physiology, 29, 279-287.

Verma, H.N. and Awasthi, L.P. (1979). Antiviral activity of *Boerhaavia diffusa* root extract and the physical properties of the virus inhibitors. Can. J. Bot. 57, 926-932.

Von Arnold, S. (1988). Tissue Culture methods for clonal propagation of forest trees. News Letter, I.A.P.T.C., 56, 2-13.

Wadhwa, V.; Singh, M.M.; Gupta, D.N.; Singh, C. and Kamboj, V.P. (1986). Contraceptive and hormonal properties of *Achyranthes aspera* in Rats and Hamsters. Planta Medica, 231-233.

Waller, G.R. and Nowacki, E.K. (1978). In : 'Alkaloid Biology and Metabolism in Plants'. Plenum Press, New York.

West, F.R. and Mika, E.S. (1957). Synthesis of atropine by isolated roots and root-callus cultures of belladonna, Bot. Gaz. 50, 119.

White, P.R. (1934). Potentially unlimited growth of excised tomato root tips in a liquid medium. Plant Physiol. 9, 585-600.

White, P.R. (1939). Potentially unlimited growth of excised plant callus in an artificial nutrient. *Am.J.Bot.* **26**, 59-64

White, P.R. (1943). "A Handbook of Plant Tissue Culture". Jacques Cattel Press.

White, P.R. (1954). The Cultivation of Animal and Plant Cells. The Ronald Press, New York, 239.

Widholm, J.M. (1972). Cultured Nicotiana tabacum cells with an altered anthranilate synthetase which is less sensitive to feedback inhibitor. *Biochem. Biophys. Acta*, **261** : 52-58.

Widholm, J.M. (1980) Selection of plant cell lines which accumulate compounds. In : 'Plant Tissue Culture As a Source of Biochemicals (ed. Staba, E.J.). CRC Press, Florida, 99-114.

Yeoman, M.M. (1987). In : Plant Cell Culture Technology (ed. Yeoman, M.M.). Narosa Publishing House, New Delhi.

Yeoman, M.M.; Miedzybrodzka, M.B.; Lindsey, K. and McLauchan, W.R. (1980). In Plant Cell Cultures : Results and Perspectives (ed. Sala, E.; Parisi, R.; Cella, R. and Cifervi). Elsevier, 327-343.

Zenk, M.H. (1978). The Impact of cell on industry. In : Frontiers of Plant Tissue Culture (ed. Thorpe, T.A.). University of Calgary, Calgary, 1-13.

Zenk, M.H; El-Shagi, H.; Arens, H; Stockigt, J; Weiler, E.W. and Dells, B. (1977). Formation of the indole alkaloids serpentine and ajmalicine in cell suspension cultures of *Catharanthus roseus*. In : "Plant Tissue Culture and Its Biotechnological Application" (eds. W.Barz, E.Reinhard, and M.H.Zenk), . Springer-Verlag, 27-43.

Zenk, M.H.; El-Shagi, H. and Schulte, U. (1975). Anthraquinone production by cell suspension culture of *Morinda citrifolia*. *Planta Med. Suppl.* **79**, 101.