

## Materials & Methods

## CHEMICALS

Benzamide, trypan blue, pepstatin A, ALLN, NAD<sup>+</sup>, agarose, Annexin V-FITC/PI dual staining kit, APO-BRDU kit, Caspase-3 assay kit, 1,6 diphenyl 1,3,5-hexatriene (DPH), 4',6-diamidino-2-phenylindole (DAPI), 3,3'-dihexyloxycarbocyanine iodide (DiOC<sub>6</sub>), 2',7' dichlorodihydrofluorescein diacetate (H2DCFDA-AM), Protease Inhibitor Cocktail, anti-mouse IgG (whole molecule) FITC conjugate, anti rabbit IgG (whole molecule) TRITC conjugate, sheep anti-cytochrome C polyclonal antibodies raised against rabbit, anti sheep IgG (whole molecule) TRITC conjugate (Sigma, USA), anti-PAR mouse mAb (10H) (Calbiochem, Germany), N-Succinyl-LLVY-AMC (Calbiochem, Germany), TRIZOL reagent (Invitrogen, USA), rabbit anti-AIF polyclonal antibodies raised against amino acids 151-180 of human AIF (Cayman Chemical, USA). Other reagents were of analytical grade from Merck and Himedia.

## METHODS

### *D. discoideum* culturing

*D. discoideum*, Ax-2 strain which is an axenic derivative of Raper's wild type NC-4 (a mutant in at least two genes i.e. *axe* A and *axe* B, Newell *et al.*, 1982 and Deering *et al.*, 1988) was used. *D. discoideum* was grown under different culture conditions. The growing cells (unicellular) were maintained in a liquid suspension (HL5 medium). *D. discoideum* cells were grown in modified HL-5 medium, pH 6.5 with 150 rpm shaking at 22°C (Watts and Ashworth, 1970). Log phase cells at a density of  $\sim 2.5 \times 10^6$  cells/ml were used for experiments.

It was also maintained on a solid substratum containing Phosphate Buffered Agar (PBA). *D. discoideum* was also cultured on bacterial lawn of *Klebsiella* which is its natural food. For this a loop full of overnight grown culture of *Klebsiella* was taken and *D. discoideum* spores (4-5) were mixed with it. This 'mixture' of two cell types is then pour plated on PBA plates (90mm). *D. discoideum* cells fed on *Klebsiella* and when no more *Klebsiella* was left the *D. discoideum* cells undergo developmental changes and form fruiting bodies.

**Preparation of Axenic Medium (Oxoid) – (modified HL5, Ashworth and Watts, 1970; Kosta *et al.*, 2001)**

Proteose peptone	-	14.3 gm
Yeast extract	-	7.5 gm

---

Maltose	-	18 gm
Na <sub>2</sub> HPO <sub>4</sub> 2H <sub>2</sub> O	-	0.616 gm
KH <sub>2</sub> PO <sub>4</sub> 2H <sub>2</sub> O	-	0.486 gm

The above constituents were added to 1000ml of distilled water and pH was adjusted to 6.7. The medium (10 ml) was then dispensed into conical flasks and were autoclaved at 15 psi for 15 minutes. Care was taken so that the medium would not be charred by over heating.

#### **Phosphate Buffer / Sorenson's buffer / KK<sub>2</sub> buffer (50X SB)**

2mM Na<sub>2</sub>HPO<sub>4</sub>

15mM KH<sub>2</sub>PO<sub>4</sub>

pH 6.4

It was sterilized by autoclaving at 15 psi for 15 minutes. 1X SB buffer was obtained by adding 1 ml of 50X SB to 49 ml of autoclaved distilled water.

#### **Phosphate Buffered Agar (PBA)**

2 gm agar was added to 98 ml distilled water and autoclaved at 15psi for 15 minutes. After autoclaving 2 ml 50X SB was added.

### **Culture Maintenance**

#### **Growth of Ax-2 cells in Axenic medium (Levraud *et al.*, 2001)**

- i). 10 ml of the autoclaved HL-5 medium was taken and approximately 100  $\mu$ l (inoculum size = 1% of medium volume, approx  $0.05 \times 10^5$  cells/ml) of log phase culture with a cell density of  $2 \times 10^6$  cells/ml were used for inoculation in the medium under sterile conditions.
- ii). The flasks were incubated at 22 °C under shaking conditions maintained at a speed of 150 rpm.
- iii). Subculturing can be done twice a week (till one month), after which the culture must be revived from the spores. The first or second subculture from this spore revival was used for experimental purposes.

#### **Differentiation of *D. discoideum* cells on non nutrient agar (Levraud *et al.*, 2001)**

- i). Exponentially growing cells from HL-5 medium were pelleted down at 300g/5min/4°C.
- ii). The pellet was washed with 1X SB (Sorenson's buffer) thrice at 300g/5min/4°C.

- iii). The final cell density was adjusted to  $2.5 \times 10^6$  cells / ml and 100 $\mu$ l of cell suspension was then spread on PBA (non-nutrient agar) containing 35 mm plate and allowed to differentiate at 22 °C to form fruiting bodies.

### Spore Revival

- 5 ml HL-5 medium was dispensed in 25 ml conical flasks and autoclaved.
- The fruiting bodies were picked up from the surface PBA plates with a sterile Nichrome loop. The loop must not be allowed to touch the agar surface at any point, and all other precautions must also be taken to avoid bacterial contamination.
- Few fruiting bodies were inoculated in each flask, and then incubated under shaking conditions at 22 °C /150 rpm. Spores germinate with overnight incubation but log phase culture that can be used for further sub-culturing, could be obtained after about a week.

### Cell viability test by trypan blue exclusion technique (Kosta *et al.*, 2001)

This is a vital staining procedure, where Trypan blue stains dead cells exclusively as live cells can pump out the dye. The stained dead cells then appear blue, while viable cells remain colourless. 20  $\mu$ l of the culture to be tested is taken and 10  $\mu$ l of 0.1% Trypan blue solution is added. A cell count is taken using the haemocytometer. The percentage viability is then calculated using the following formula

$$\frac{\text{Number of dead cells}}{\text{Total number of cells}} \times 100 = \% \text{ Dead cells}$$

Note: Trypan blue is commercially available as 0.4% stock solution. Diluted it 4 times with 1X SB to make 0.1% working solution.

### UV-C irradiation of *D. discoideum* cells

Cells were given different UV-C exposures in 1XSB / KK<sub>2</sub> buffer in a laminar as follows

Shaker	= 150 rpm
Distance from UV source	= 23 cm
Radius of petriplate	= 3.5 cm
Time of exposure	= 0.8, 1, 2, 3, 5 or 10 min
Transmission	= 72.3 %

$$\frac{\text{Time of exposure} \times \text{specimen's area in m}^2 \times \text{power flux}}{(\text{Distance})^2} = \text{Actual energy incident on sample in J/m}^2$$

2-3 X10<sup>6</sup> cells were harvested by centrifugation at 300g / 5 min / 4<sup>0</sup> C. Washed the cells 2-3 times with 1X SB and finally resuspended in 1.0 ml 1XSB and transferred in Tarson plastic petri plate (35 mm). Different doses (0, 1.0, 2.0, 3.0, 5.0, 10.0, min) of UV-C were given to different petriplates and all the plates were shifted immediately to BOD at 22°C for 30 min. under dark condition to avoid photo-reactivation recovery. After incubation, irradiated cells from each plate were transferred to different eppendorfs for centrifugation and pelleted down by 300g /5min /4<sup>0</sup> C. To the pellet HL<sub>5</sub> medium, pH 6.5 was added and kept on shaker at 120 rpm / 21<sup>0</sup> C for growth. Cell death was monitored by Trypan blue exclusion method till 22hour.

#### **Assessment of cell death by AnnexinV-FITC/PI dual staining (Miller, 2004)**

To differentiate between apoptotic and necrotic cell death, dual staining with Annexin V-FITC/PI was performed using Apoptosis detection kit (Molecular Probes, USA). Phosphatidyl serine (PS) exposure on the surface of the cells is one of the characteristic features of apoptotic cells. It is also a signal for macrophages to clear such cells by phagocytosis which is important to prevent inflammation and damage to the surrounding cells. PS has an affinity to bind to Annexin in a Ca<sup>+2</sup> dependent manner. Annexin can bind to PS only when it is exposed outside as it cannot cross the membrane. Propidium Iodide (PI) is a DNA binding dye which can enter the cell only when membrane integrity is completely lost. Thus early apoptotic cells will be Annexin+ and PI- while late apoptotic and necrotic cells will be Annexin+ and PI+.

2-3 X10<sup>6</sup> cells were pelleted and washed twice with 1X SB. *D. discoideum* cells were then suspended in binding buffer provided in the kit and incubated with Annexin V and PI for 10 minutes in dark at 22°C. Fluorescence was monitored under 60X magnification using fluorescent microscope (Nikon Eclipse TE2000-S, Japan). Dose and time dependent study was done to standardize the apoptotic and necrotic doses for further experiments.

**Measurement of PARP activity by indirect immunofluorescence (Cole and Perez-Polo, 2002)**

PARP activity was assayed indirectly by using antibodies against the product of PARP i.e. PAR. For assaying PARP, indirect immunofluorescence was done using anti-PAR mouse mAb (10H) (Calbiochem, Germany) at a concentration of 0.5ug/ml and anti-mouse IgG (whole molecule) FITC conjugate (Sigma) at a dilution of 1:200. *D. discoideum* cells were pelleted and washed once with phosphate buffered saline (PBS) pH 7.4, fixed in 70% chilled methanol for 10 minutes at -20 °C and then washed with blocking solution (1.5% BSA with 0.05% Tween 20 in PBS) followed by incubation for 1 hour in primary antibody. After incubation the cells were washed 2-3 times with blocking solution and further incubated for 1 hour with FITC labeled secondary antibody. Finally these cells were washed 2-3 times with PBS and the fluorescence was observed at 490 nm under 60X magnification. PARP assay was performed at different time intervals as described in the experiments later.

**ROS measurement by DCFDA dye (Esposti, 2002)**

In order to observe the formation of reactive oxygen species a fluorescent dye is employed. The acetate ester form of 2',7' dichlorodihydrofluorescein diacetate (H2DCFDA-AM) is a membrane permeable molecule that passes through the cell membrane. Once inside the cell, cellular esterases act on the molecule to form the non-fluorescent moiety H2DCFDA, which is ionic in nature and therefore trapped inside the cell. Oxidation of H2DCFDA by ROS converts the molecule to 2',7' dichlorodihydrofluorescein (DCF), which is highly fluorescent. Upon stimulation, the resultant production of ROS causes an increase in fluorescence signal over time.

$2 \times 10^6$  cells were harvested and washed with 1X SB twice. DCFDA (50 nM) was added to cells and was incubated for 15 min at 22 °C with shaking, followed by two washes with 1X SB. Fluorescence unit was measured by fluorimeter (F7000, Hitachi, Japan) using 200 µl sample diluted 5 times using KK2 buffer. Excitation ( $\lambda_{ex}$ ) and emission ( $\lambda_{em}$ ) wavelengths used for fluorimetric studies were 480 and 525 nm respectively.

**Estimation of NAD<sup>+</sup> and ATP levels (Bernofsky and Swan, 1973; Leoncini *et al.*, 1987)**

Intracellular levels of NAD<sup>+</sup> in *D. discoideum* were determined by enzymatic recycling method employing alcohol dehydrogenase to reduce NAD<sup>+</sup> to NADH. NAD<sup>+</sup> present in

the cell is used to convert ethanol into acetaldehyde, the reaction being catalyzed by alcohol dehydrogenase. NADH produced in the first reaction is used to reduce MTT in to Formazan and  $\text{NAD}^+$  is regenerated. In this way cellular  $\text{NAD}^+$  keeps on getting recycled and thereby enhances the yield of Formazan, a chromophore which gives absorbance at 570 nm. In brief, cells were exposed to 0.05 and 0.1 mM cumene  $\text{H}_2\text{O}_2$  for 1 hour and cultures were washed twice with ice cold PBS and  $\text{NAD}^+$  was extracted with 1 ml of 0.5 M perchloric acid and then neutralized with 1 N KOH.  $\text{NAD}^+$  levels were estimated by taking the absorbance at 570 nm following protein estimation with Lowry method (Lowry *et al.*, 1951).

ATP was determined by HPLC in aliquots extracted with alkali according to Leoncini *et al.*, (1987). Cells exposed to 0.05 and 0.1 mM cumene  $\text{H}_2\text{O}_2$  were washed with cold saline solution (containing 10mM glucose and 2mM EDTA) after 1 hour of stress induction. Nucleotides were extracted by treating the cell pellet with 0.5 M KOH which were kept on ice for 2 minutes and immediately sonicated (four 10 sec bursts) to ensure maximal extraction. The extract was neutralized with 1 M  $\text{KH}_2\text{PO}_4$ , centrifuged at 8000g for 10 minutes and the supernatant was used for HPLC analysis using rp-C18 column (5 $\mu\text{m}$ ) and UV detector. 1 M  $\text{KH}_2\text{PO}_4$  was used as mobile phase and flow rate was maintained at 1 ml/min.

#### **Evaluation of mitochondrial membrane potential (Koning *et al.*, 1993)**

Potential sensitive dye DiOC<sub>6</sub> (3,3'-dihexyloxycarbocyanine iodide) (Sigma) was used to evaluate changes in mitochondrial membrane potential (MMP). DiOC<sub>6</sub> is a membrane permeable lipophilic cationic dye which binds specifically to mitochondria proportionally to the difference between the negativity of the cytoplasmic compartment and the mitochondrial matrix. Dissipation of the mitochondrial membrane potential reduces the affinity of binding of the dye.

To observe the change in MMP, time dependent study was done using 0.05 mM cumene  $\text{H}_2\text{O}_2$  dose (non-apoptotic dose) and 0.1 mM cumene  $\text{H}_2\text{O}_2$  dose (necrotic dose) as standardized by PS-PI dual staining.  $\sim 2.0 \times 10^6$  cells were pelleted and washed twice with 1X SB. Cells were stained with DiOC<sub>6</sub> (400 nM) for 15 minutes in dark and then washed once with 1X SB and monitored for the fluorescence as above.

**Monitoring AIF release by indirect immunofluorescence (Bidere *et al.*, 2003)**

Release of AIF from mitochondria to cytosol and its translocation to nucleus was monitored by detecting immunofluorescence at different time intervals. Same protocol as for PAR assay was followed using rabbit anti-AIF polyclonal antibodies raised against amino acids 151-180 of human AIF (Cayman Chemical) at 1:1000 dilution and anti rabbit IgG (whole molecule) TRITC conjugate (Sigma) at 1:400 dilution. Nuclear counterstaining with DAPI (1 $\mu$ g/ml) for 5 minutes was performed after the removal of excess secondary antibody and observed for the fluorescence.

**Cytochrome c release (Granville *et al.*, 2001)**

Release of cytochrome c from mitochondria into cytosol was monitored by detecting immunofluorescence at different time intervals. Same protocol as for AIF was followed using sheep anti-cyt c polyclonal antibodies (Sigma) raised against rabbit at 1:200 dilution and anti sheep IgG (whole molecule) TRITC conjugate (Sigma) at 1:400 dilution. Nuclear counterstaining with DAPI (1 $\mu$ g/ml) for 5 minutes was performed after the removal of excess secondary antibody and fluorescence was monitored.

**Monitoring DNA fragmentation by agarose gel electrophoresis and TUNEL assay (Hershberger *et al.*, 1994; Noegel *et al.*, 1985; Gavrieli *et al.*, 1992)**

Genomic DNA and large scale fragments (>50kb) were run on 0.8% agarose gel. To monitor internucleosomal fragmentation cells were harvested by low-speed centrifugation. Apoptotic vesicles were pelleted from the growth medium supernatant by centrifugation (14,000 x g, 15 min) at 4°C. Cells and vesicles were pooled, suspended in lysis buffer (10 mM Tris [pH 7.5], 10 mM EDTA, 0.2% Triton X-100), and extracted for low-molecular-weight DNA as previously described (Ishida *et al.*, 1992). DNA samples were treated with 0.6 mg of RNase A per ml prior to electrophoresis, using 2% agarose-Tris-borate-EDTA gels and visualization with ethidium bromide.

The DNA fragmentation creates a large number of 3'-hydroxyl sites at the DNA breaks. This property is used in the APO-BRDU kit (Sigma, USA) to identify apoptotic cells by labeling the 3'-hydroxyl sites with bromodeoxyuridine triphosphate (Br-dUTP). Br-dUTP is enzymatically attached to the 3-hydroxyl sites of double- or single-stranded DNA by terminal transferase (TdT). Non-apoptotic cells do not incorporate Br-dUTP due to the lack of available 3-hydroxyl sites.

Initial standardization of DNA fragmentation analysis was done at different time intervals. After exposure of cells to UV-C paraptotic and necrotic dose respectively DNA isolation was carried out from the cells by the conventional method as described by Noegel *et al* (1985) and was subjected to electrophoresis using 0.8% agarose gel. TUNEL assay (Terminal deoxynucleotidyl transferase dUTP nick end labeling) was performed according to the procedure supplied by the manufacturer (ApoBrdU DNA fragmentation kit; Sigma). Fluorescence was monitored by using Anti Brdu-FITC antibody.

#### **Monitoring stress induced DNA damage by immunofluorescence (Minami *et al.*, 2005)**

Histones get phosphorylated during oxidative stress. Hence presence of phosphorylated histone indicates oxidative stress induced DNA damage. Anti-H2AX at a concentration of 0.5ug/ml and anti-mouse IgG (whole molecule) TRITC conjugate (Sigma) at a dilution of 1:200 were used to study the presence of oxidative stress. Cells were pelleted and washed once with phosphate buffered saline (PBS) pH 7.4, fixed in 70% chilled methanol for 10 minutes at -20°C and then washed with blocking solution (1.5% BSA with 0.05% Tween 20 in PBS), incubated for 1 hour in primary antibody. After incubation the cells were washed 2-3 times with blocking solution and further incubated for 1 hour with TRITC labeled secondary antibody. Washed 2-3 times with PBS and observed the fluorescence under 60X.

#### **PARP inhibition by benzamide (Szabo and Dawson, 1998)**

Log phase culture of *D. discoideum* with a cell count of  $2 \times 10^6$  cells was incubated with different doses (1.0, 2.0, 3.0, 4.0 mM) of benzamide (Sigma, USA) for 24 hours. For further experiments, 1 mM benzamide was used as it showed 2% cell death only. Cells were preincubated with 1 mM benzamide and then stressed with different agents and observed for different cell death parameters. To show the specificity of benzamide similar studies were also carried out with non inhibitory structural analog of benzamide, benzoic acid.

#### **Detection of caspase activity (Olie *et al.*, 1998)**

The substrate DEVD-AMC is an oligopeptide that is covalently linked with the fluorophore 7- amino 4- methyl coumarin (AMC). DEVD is the cleavage site for

Caspases (3 and 7). The cleavage takes place at the C terminal of the last aspartate residue, thus liberating the fluorophore AMC which can be estimated and/or visualized under the fluorescence microscope (AMC  $\lambda_{ex}$  = 380 nm).

$2-3 \times 10^6$  cells were harvested and washed with PBS. Cells were resuspended in 1ml PBS and an aliquot of 100  $\mu$ l was taken from it and 10 $\mu$ l of DEVD-AMC substrate (1mg/ml) was added. After an incubation of 1 hour cells were observed under the fluorescence microscope. AMC liberated from the fluorogenic substrate was measured at 380 nm using UV filter. Caspase activity as a function of AMC fluorescence was monitored at different time intervals. Effect of caspases on cell death was also studied by using caspase-3 specific inhibitor. The entire procedure was carried out using Caspase-3 assay kit (Sigma) as per manufacturer's instructions.

#### **Characterization of apoptotic vesicles (Gautam and Sharma, 2002)**

Isolation of apoptotic vesicles was performed from *D. discoideum* culture after 16 hours of UV-C stress as mentioned by Gautam and Sharma (2002). The culture was centrifuged at 1000g for 4minutes at 4°C; the supernatant was collected and centrifuged at 21,000g for 45minutes at 4°C. The obtained pellet was washed once with 1X SB and used for further analysis. Formation of apoptotic vesicles was monitored at different time intervals. Isolated vesicles were stained separately with a fluorescent membrane probe, 1,6 diphenyl 1,3,5-hexatriene (DPH) at a concentration of 1 $\mu$ M, 4',6-diamidino-2-phenylindole (DAPI), PS-PI and then observed under fluorescence microscope.

#### **Investigation of proteases involved in cell death (Moubarak *et al.*, 2007)**

Cells were pre-incubated with protease inhibitor cocktail (Sigma, USA) followed by stress and MMP changes and PS exposure were checked. Similarly MMP changes, PS exposure and vesicle formation were followed in stressed cells pre-incubated for 12 hours with 0.5  $\mu$ g/ml pepstatin A, a cathepsin D inhibitor and 10  $\mu$ M ALLN, a Calpain inhibitor.

#### **DNA Isolation from *D. discoideum* (Pilcher *et al.*, 2007)**

$10^7$  cells/ml were harvested and washed once with 1X SB, followed by one wash with ice cold 0.2 % NaCl (0.034M). One volume 2% SDS was added and incubated at 65 °C for 15'. One volume TE buffer (pH 9.5) was added and then extracted with one volume of

Phenol: Chloroform (1:1). The upper aqueous layer was collected after centrifugation at 9000g for 5' and extracted it with one volume of chloroform. Centrifuge at 9000g for 5' and collect the upper aqueous layer. 1/10<sup>th</sup> volume of sodium acetate and one volume of isopropanol was added. Complete precipitation was done by incubating at -20 °C for more than 30'. The pellet was obtained by centrifuging at 9000g for 5'. Pellet was washed once with 75% ethanol. Dried the pellet in oven and resuspended it in autoclaved DDW (~20 µl). The DNA preparation was stored at -20 °C till further use.

### PCR Standardization for Amplification of *aif* gene

PCR reaction was setup as follows:

DNA	3.0µl (100 ng)
Forward Primer	0.5µl (10 pmol)
Reverse Primer	0.5µl (10 pmol)
Master Mix (2X)	12.5µl
DDW	8.5µl
Total	25µl

Reaction parameters were as follows:

Reaction parameters	Temperature and Time
Initial Denaturation	94 °C for 6 min
Denaturation	94 °C for 45 sec
Annealing	63 °C for 30 sec
Polymerization	72 °C for 45 sec
Repeat cycle 29 times	
Final Extension	72 °C for 10 min
Hold	22 °C

### Plasmid isolation from *E. coli*

Alkaline lysis method for isolation of plasmid from *E. coli* has been followed (Sambrook and Russell, 2001):

2 ml over night grown culture was pelleted down. Washed once with 0.25 volume of STE and 100 µl of AL-I was added and vortex vigorously. 200 µl AL-II was added and mixed well. 150 µl AL-III was added and mixed well, stored on ice for 10 min. Centrifuged at

9K/5' and collected supernatant. It was extracted with one volume of Phenol: Chloroform (1:1). Centrifuged at 9K/5' and the upper aqueous layer was transferred in fresh eppendorf. It was then extracted with one volume of Chloroform. Centrifuge at 9K/5' and again collect the upper aqueous layer. 1/10<sup>th</sup> volume of Sodium acetate and one volume of Isopropanol was added to it and incubated at -20 °C for 30'. Centrifuge at 9K/5' and supernatant was discarded. Washed once with 75% ethanol and evaporated the ethanol completely. Finally pellet was resuspended in autoclaved DDW (~20 µl).

#### **Transformation of plasmid DNA in *E.coli* (Sambrook and Russell, 2001)**

Inoculated 3 ml LB with *E.coli DH5α* and incubated at 37°C till it reaches OD ~ 0.4 - 0.6 (log phase). 3 ml log phase culture of *E.coli* was pelleted down at 5000g/5min. Added 1.8 ml 80 mM MgCl<sub>2</sub> + 20 mM CaCl<sub>2</sub> and kept it on ice for 20'. Pelleted the cells and washed once with 1 ml 100 mM CaCl<sub>2</sub>. Added 100 µl 100 mM CaCl<sub>2</sub> and incubated on ice for ~ 1 hour to prepare competent cells. Added DNA and incubated on ice for 30 minutes. Heat shock was given at 42 °C for 90 seconds and immediately transferred on ice for 2-3 minutes and then added 800 µl of sterile luria broth (LB) and incubated at 37 °C for 45 minutes under shaking conditions. Pelleted down the cells and resuspended in 40 µl LB. Spread it on LA plates containing ampicillin.

#### **Transformation of plasmid DNA in *D. discoideum* (Gaudet *et al.*, 2007)**

Pelleted down 5x10<sup>6</sup> cells. Washed twice with ice cold H-50 buffer. Added 2 µl (1-10 µg DNA) of cloned plasmid DNA. Incubated on ice for 5 minutes. The cell suspension was transferred to a cold 0.2 cm electroporation cuvette. Electroporated in cold at 0.8kV/25 µF pulse, twice for about 5 second gap between 2 pulses. Incubated the cuvette on ice for 5 minutes. Set up 10cm Petri dish for selection with sterile 10ml HL5 and cells were transferred out of cuvette with a few ml of HL5 to the plate. Incubated in BOD at 22 °C for overnight.

Added antibiotics from the next day with medium change as follows:

	1 <sup>st</sup> day	3 <sup>rd</sup> day	5 <sup>th</sup> day	Alternate day
Ampicillin	1 µg/ml	1 µg/ml	1 µg/ml	1 µg/ml
G418	5 µg/ml	10 µg/ml	15 µg/ml	15 µg/ml

### RNA Isolation Protocol (Pilcher *et al.*, 2007)

Harvested  $2 \times 10^7$  cells and washed with 1 ml KK2. The pellet can be stored at  $-80^\circ\text{C}$ . Re-suspended the pellet in 500  $\mu\text{l}$  Trizol reagent (Invitrogen, USA) by pipetting up and down. Incubated at room temperature (RT) for 5 min. Added 200  $\mu\text{l}$  chloroform and stored at RT for 5 min. Centrifuged for 10 min at 10,000 rpm. The upper phase was transferred in a new tube and added 1 ml 100% ethanol. Incubated for 20 min at  $-20^\circ\text{C}$ . Centrifuged for 5 min at 13,000 rpm and then removed the supernatant completely. Air dried the pellet for 5 min without allowing the pellet to completely dry out. The pellet was dissolved in 10  $\mu\text{l}$  nuclease free water. Measured the RNA content in a spectrophotometer (used a 1:100 dilution) and also checked RNA quality with an agarose gel.

### Expression analysis of developmentally regulated genes by RT-PCR

*D. discoideum* cells were exposed to oxidative stress as mentioned earlier. After one hour pretreatment cells were pelleted and washed with 1X SB and finally resuspended in 1X SB. Total RNA was isolated from the cells at two time points (0 and 9 hour) using TRIZOL reagent (Invitrogen, USA) according to the manufacturer's instructions. The expression kinetics of selected genes was examined by RT-PCR using following primers

*acaA*,

*cAR1* (5'TGTGGACTTTATGTCTTGCAATTAG3' & 5'CCAATACTGCTGAAATTGCCC3'),

*yakA* (5'GCTTAGAGGACTTTCAACCAATTT3' &

5'GATTTTTTCATAACAAGCAGATCCA 3')

*gp80* (5'GCTGATTCAGTATCCCCAACAATTTATTTC3' &

5'CAAGGTTTGTGGGCTTGAGGTTCCCATG3')

*gp24* (5'CCAGGAGCTTTTCAATGGGCAGTTGATG3' &

5'GTGTAACAGTCATATTCTTTGGGAATGTCTC3'),

*dsn1* (5'CCACCCATTAACCTGGAATG3' & 5'TGGTGGCATCAGTACAATCG3')

*ctn* (5'ATGAATAAATTATTTTCATTAATTTTAGCTTTATTCCTTGCAACTCCGC3' &

5'TTAAAATAAAGCAAAACCTGAACCTGAACCAGAGGCGGCACC3')

*ctn2* (5'GTGGTGCCGTTTGTTCATTACTCCC3' &

5'CCAGTTGGGTCAGTTACCATAACAGCAAC3')

*rnl A* (5'TTACATTTATTAGACCCGAAACCAAGCG3' & 5'TTCCCTTTAGACCTATGGACCTTAGCG3') was used as an internal control.

The reactions were performed according to the manufacturer's instructions (Fermentas, Ontario, Canada). DNA fragments were amplified 24 cycles after reverse transcription and signal intensities were analyzed on 2% agarose gel.

#### **cDNA synthesis protocol:**

##### Mixed the following:

- 1-5 µg total RNA
- 1 µl oligo dT primer (supplied at 100uM)
- 1 µl 10 mM dNTPs
- Added water to 10 µl total volume
- Incubated 10 min at 65°

##### In the meantime prepared:

- 4 µl 5x Reaction Buffer
- 1 µl Ribonuclease Inhibitor
- 4 µl water
- Added 9 µl of the mixture to 10 µl samples and incubate 5 min at 37°C
- Added 1 µl M-MuLV Reverse Transcriptase.
- Incubated 60 min at 37°C. Stopped the reaction at 70°C for 10 min.
- 1 µl of the cDNA should be around 50-100 ng and can be used in a usual PCR with gene specific primers

#### **Developmental studies**

##### **Effect of GSH/ benzamide/PIC on the growth of stressed *D. discoideum* cells**

$0.5 \times 10^6$  cells were treated with the respective chemicals for 1 hour. The cells were then exposed to different UV-C doses (10.4, 13, 26, 65 and 130 J/m<sup>2</sup>) in SB in 35mm petri plate. Following this, the cells were harvested and finally resuspended in 5 ml of sterile HL5 containing flask and growth was monitored for 6 days at an interval of 12 hours under shaking conditions at 22°C.

##### **Addition of GSH/ benzamide**

*D. discoideum* cells were subjected to 1 mM GSH/1mM benzamide treatment for 1 hour, then subjected to UV-C treatment after washing with SB. Cells were then harvested and

pellet was resuspended in 100µl of Sorenson's buffer (1X SB) and spread on PBA plate (Sroka *et al.*, 2002).

#### **Measurement of cAMP levels and exogenous addition of cAMP (1 µM and 10 µM) in *D. discoideum* cells**

5 x10<sup>6</sup> cells were harvested by centrifugation at 700g/5 minutes/ 4<sup>0</sup>C and subjected to stress. Cells were resuspended in 1X SB for 6 hours. Cells were then collected and extracellular cAMP was estimated in the buffer by ELISA kit method as per the manufacturer's instructions (Calbiochem, USA).

2-3 X10<sup>6</sup> cells pretreated (150 rpm/2.5 hours /22<sup>0</sup>C) with cAMP were harvested, resuspended in 1X SB containing 1 µM/ 10 µM cAMP and subjected to development after UV-C treatment.

#### **Nitric oxide estimation in *D. discoideum* cells (Green *et al.*, 1982)**

NO generation was estimated according to Green et al (1982). 5 x 10<sup>6</sup> cells were suspended in 1 ml 1X SB and incubated at 21<sup>0</sup>C for 20 minutes to allow the accumulation of NO generation. 1 ml Griess reagent was added and mixed well. Again incubated at 21<sup>0</sup>C for 15-30 minutes and took absorbance at 546 nm.

#### **References**

- Bernofsky C. and Swan M. (1973) An improved cycling assay for nicotinamide adenine dinucleotide. *Anal. Biochem.* 53, 452-458.
- Bidere N., Lorenzo H.K., Carmona S., Laforge M., Harper F., Dumont C. and Senik A. (2003) Cathepsin D Triggers Bax Activation, Resulting in Selective Apoptosis-inducing Factor (AIF) Relocation in T Lymphocytes Entering the Early Commitment Phase to Apoptosis *J. Biol. Chem.* 278, 31401-31411.
- Cole K.K. and Perez-Polo J.R. (2002) Poly(ADP-ribose) polymerase inhibition prevents both apoptotic-like delayed neuronal death and necrosis after H<sub>2</sub>O<sub>2</sub> injury. *J. Neurochem.* 82, 19-29.
- Esposti M.D. (2002) Measuring mitochondrial reactive oxygen species. *Methods* 26, 335-340.

- Gaudet P., Pilcher K.E., Fey P. and Chisholm R.L. (2007) Transformation of *Dictyostelium discoideum* with plasmid DNA. *Nature Protocols* 2, 1317 – 1324.
- Gautam S. and Sharma A. (2002) Rapid cell death in *Xanthomonas campestris* pv. *glycines*. *J. Gen. Appl. Microbiol.* 48, 67-76.
- Gavrieli Y., Sherman Y. and Ben-Sasson S.A. (1992) Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation. *J. Cell Biol.* 119, 493-501.
- Granville D.J., Cassidy B.A., Ruehlmann D.O., Choy J.C., Brenner C., Kroemer G., van Breemen C., Margaron P., Hunt D.W. and McManus B.M. (2001) Mitochondrial Release of Apoptosis-Inducing Factor and Cytochrome c during Smooth Muscle Cell Apoptosis. *Amer. J Pathol.* 159, 305-311.
- Green L. C., Wagner D. A., Glogowski J., Skipper P.L., Wishnok J.S. and Tannenbaum, S.R. (1982) Analysis of nitrate, nitrite, and [15N]nitrate in biological samples. *Anal. Biochem.* 126, 131-138.
- Hershberger, P. A., LaCount, D. J. and Friesen, P.D. (1994) The apoptotic suppressor P35 is required early during baculovirus replication and is targeted to the cytosol of infected cells. *J Virol.* 68, 3467-3477.
- Ishida, Y., Agata, Y., Shibahara, K. and Honjo, T.(1992) Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. *EMBO J.* 11,3887-3895.
- Koning A.J., Lum P.Y., Williams J.M. and Wright R. (1993) DiOC6 staining reveals organelle structure and dynamics in living yeast cells. *Cell Motil Cytoskeleton.* 25, 111-128.
- Kono and Fridovich. (1983) Isolation and characterization of the pseudocatalase of *Lactobacillus plantarum*. *J. Biol. Chem.* 258, 6015-6019.
- Kosta A., Laporte C., Lam D., Tresse E., Luciani M.F. and Golstein P. (2001) How to Assess and Study Cell Death in *Dictyostelium discoideum*. *Methods in Cell Biology.* 346, 535-550.

- Leoncini G., Buzzi E. Maresca M., Mazzel M. and Balbi A. (1987) Alkaline extraction and reverse phase high-performance liquid chromatography of adenine and pyridine nucleotides in human platelets. *Anal. Biochem.* 165, 379-383.
- Levraud J.P., Adam M., Cornillon S. and Golstein P. (2001) *Methods in Cell Biology.* 66, 469-498.
- Lowry O. H., Rosebraugh N.J., Farr A.L. and Randall R.J. (1951) Protein measurement with the Folin Phenol reagent *Anal. Biochem.* 193, 265-275.
- Miller E. (2004) Apoptosis Measurement by Annexin V Staining. *Methods Mol. Med.* 9, 191-202.
- Minami,T., Adachi, M., Kawamura, R., Zhang, Y., Shinomura, Y. and Imai, K. (2005) Sulindac enhances the proteasome inhibitor bortezomib-mediated oxidative stress and anticancer activity. *Clin. Cancer Res.* 11, 5248-5256.
- Moubarak R.S., Yuste V.J., Artus C., Bouharrou A., Greer P.A., de Murcia J.M. and Susin S.A. (2007) Sequential activation of poly(ADP-ribose) polymerase 1, calpains, and Bax is essential in apoptosis-inducing factor-mediated programmed necrosis. *Mol. Cell. Biol.* 27, 4844-4862.
- Pilcher K.E., Fey P., Gaudet P., Kowal, A.S. and Chisholm R.L. (2007) A reliable general purpose method for extracting genomic DNA from *Dictyostelium* cells. *Nature Protocols* 2, 1325-1328.
- Olie R.A., Durrieu F., Cornillon S., Loughran G., Gross J., Earnshaw W.C. and Golstein P. (1998) Apparent caspase independence of programmed cell death in *Dictyostelium*. *Current Biol.* 8, 955-958.
- Sroka J., Madeja Z., Michalik M., Przystalski S. and Korohoda W. (2002) Folic acid, ascorbic acid and sodium selenite restore the motility of *Dictyostelium discoideum* inhibited by triethyllead. *Toxicol.* 180, 275-292.
- Szabo C. and Dawson V.L. (1998) Role of Poly(ADP-ribose) synthetase in inflammation and ischaemia- reperfusion. *Trends Pharmacol. Sci.* 19, 287-298.
- Watts D.J. and Ashworth J.M. (1970) Growth of myxamoebae of the cellular slime mould *Dictyostelium discoideum* in axenic culture. *Biochem J.* 119, 171-174.