

REFERENCES

References

- Anderson, T.F., 1969. Self-diffusion of carbon and oxygen in calcite by isotope exchange with carbon dioxide. *J. Geophysical Research*, 74: 3918-3932.
- Andrews, J.N., Balderer, W., Bath, A.H., Calusen, H.B., Evans, G.V., Florkowski, T., Goldbrunner, J.E., Ivanovich, M., Lossli, H. and Zojer, H., 1984. Environmental isotope studies in two aquifer systems, in *Isotope Hydrology 1983 (Proceedings of the International Symposium)*: Vienna, International Atomic Energy Agency, pp. 535-576.
- Amundson, R.G., Chadwick, O.A., Sowers, J.M. and Doner, H.E., 1989. The stable isotope chemistry of pedogenic carbonates at Kyle Canyon, Nevada. *Soil Science*, 53:201-210.
- Apple, E., Rosler, W. and Corvinus, G., 1991. Magnetostratigraphy of the Miocene-Pleistocene Surai Khola, Siwaliks in West Nepal. *Journal of Geophysical International*, 105: 191-198.
- Bath, A.H., Edmunds, W.M. and Andrews, J.N., 1979. Paleoclimatic trends deduced from the hydrochemistry of a Triassic sandstone aquifer, United Kingdom, in *Isotope Hydrology 1978 (Proceedings of the Symposium, Neuherberg, 1978)*, 1: Vienna, International Atomic Energy Agency, pp. 545-568.
- Berner, R. A., Lasaga, A. C. and Garrels, R. M., 1983. The carbonate-silicate geochemical cycle and its effect on atmospheric carbon dioxide over the past 100 million years. *American Journal of Science*, 283: 641-683.
- Berner, R.A., 1991. A model for atmospheric CO₂ over Phanerozoic time. *American Journal of Science*, 291:339-376.
- Berner, R.A. and Kothavala, Z., 2001. GEOCARB III: A revised model of atmospheric CO₂ over Phanerozoic time. *American Journal of Science*, 301: 182-204.
- Biscaye, P.E., 1965. Mineralogy and sedimentation of recent deep sea clay in the Atlantic Ocean and adjacent seas and oceans. *Geological Society of American Bulletin*, 76: 803-832.
- Bird, M.I. and Chivas, A.R., 1988. Stable isotope evidence for low temperature kaolinitic weathering and post-formational hydrogen-isotope exchange in Permian kaolinites. *Chemical Geology*, 72:249-265.
- Burbank, D.W., Derry, L.A. and Lanord, C.F., 1993. Reduced Himalayan sediment production 8 Myr ago despite an intensified monsoon. *Nature*, 364:48-50.

- Burbank, D.W., Beck, A.R. and Mulder, T., 1996. The Himalayan foreland basin. In: An Yin and Mark Harrison (Eds.), *The tectonic evolution of Asia*. Cambridge University Press, USA. pp. 149-188.
- Cande, S.C. and Kent, D.V., 1995. Revised calibration of the geomagnetic polarity timescale for the late Cretaceous and Cenozoic. *Journal of Geophysical Research*, 100: 6093-6095.
- Capuano, R.M., 1992. The temperature dependence of hydrogen isotope fractionation between clay minerals and water: evidence from a geopressured system. *Geochimica et Cosmochimica Acta*, 56:2547-2554.
- Casshyap, S.M., Tewari, R.C. and Khan, A., 1993. Alluvial fan origin of the Bagra Formation (Mesozoic Gondwana) and tectono-stratigraphic implications. *Journal Geological Society of India*, 42:269-279.
- Cerling, T.E., 1984. The stable isotopic composition of modern soil carbonate and its relationship to climate. *Earth and Planetary Science Letters*, 71: 229-240.
- Cerling, T.E., 1991. Carbon dioxide in the atmosphere: Evidence from Cenozoic and Mesozoic paleosols. *American Journal of Science*, 291:377-400.
- Cerling, T.E., Quade, J., Wang, Y. and Bowman, J.R., 1989. Carbon isotopes in soils and paleosols as ecology and paleoecology indicators. *Nature*, 341: 138-139.
- Cerling, T.E., Harris, M. J., MacFadden J.B., Leakey, G. M., Quade, J., Eisenmann, V. and Ehleringer, R. J., 1997, Global vegetational change through the Miocene/Pliocene boundary. *Nature*, 389: 153-158.
- Chatterjee, S. and Roychowdhury, T., 1974. Triassic Gondwana vertebrates from India. *Indian Jour. Earth Sciences*, 1: 96-112.
- Clark, J.F., Stute, M., Schlosser, P., Drenkard, S. and Bonani, G., 1997. A tracer study of the Floridian aquifer in southeastern Georgia: implications for groundwater flow and paleoclimate. *Water Resources Research*, 33:281-289.
- Coleman, M. and Hodges, K., 1995. Evidence for the Tibetan plateau uplift before 14 Myr ago from a new minimum age for east-west extension. *Nature*, 374:49-52.
- Coplen, T.B., Kendall, C. and Hopple, J., 1983. Comparison of stable isotope reference samples. *Nature*, 302: 236-238.
- Corvinus, G., 1990. Litho and biostratigraphy of the Siwalik succession in Surai Khola area, Nepal. In Jain, K.P. and Tiwari, R.S. (eds.), *Proc. Symp. 'Vistas in Indian Palaeobotany'*, *Palaeobotanist*, 38: 293-297.

- Criss, R.E., 1999. Principles of stable isotopic distribution. Oxford University Press, New York.
- Dalai, T.K., Bhattacharya, S.K. and Krishnaswami, S., 2002. Stable isotopes in the source waters of the Yamuna and its tributaries: Seasonal and altitudinal variations and relation to major cations. *Hydrological Processes*, 16: 3345-3364.
- Davidson, G.R., 1995. The stable isotopic composition and measurement of carbon in soil CO₂. *Geochimica et Cosmochimica Acta*, 59:2485-2489.
- Deak, J., 1979. Environmental isotopes and water chemical studies for groundwater research in Hungary. In *Isotope Hydrology 1978 (Proceedings of the Symposium, Neuherberg, 1978)*, 1: Vienna, International Atomic Energy Agency, pp. 221-249.
- Decelles, P.G., Gehrels, G.E., Quade, J., Ojha, T.P., Kapp, P.A. and Upreti, B.N., 1998. Neogene foreland basin deposits, erosional unroofing and kinematic history of the Himalayan fold-thrust belt, western Nepal. *Geological Society of American Bulletin*, 110: 2-21.
- Dörr, H. and Munnich, K.O., 1987. Annual variation in soil respiration in selected areas of the temperate zone. *Tellus*, 39B: 114-121.
- Dubey, A.K., 1997. Simultaneous development of noncylindrical folds, frontal ramps and transfer faults in a compressional regime-experimental investigations of Himalayan examples. *Tectonics*, 16: 336-346.
- Edwards, M. A., Kidd, W.S.F., Li, J., Yue, Y. and Clark, M., 1996. Multi stage development of the southern Tibet detachment system near Khula Kangri. New data from Gonto La. *Tectonophysics*, 260: 1-20.
- Ekart, D.D., Cerling, T.E., Montanez, I.P. and Tabor, N.J., 1999. A 400 million year carbon isotope record of pedogenic carbonate: implications for paleoatmospheric carbon dioxide. *American Journal of Science*, 299: 805-827.
- Freeman, K.H. and Colarusso, L.A., 2001. Molecular and isotopic records of C₄ grassland expansion in the late Miocene. *Geochimica et Cosmochimica Acta*, 65:1439-1454.
- Friedman, I. and O'Neil, J.R., 1977. Compilation of stable isotopic fractionation factors of geochemical interest. In: M. Fleischer (Editor), *Data of Geochemistry*. U.S. Geol. Surv. Prof. Pap., 440-K, pp-12.

- Fritz, P., Reardon, E.J., Barker, J., Brown, R.M., Cherry, J. A., Killey, R. W. D. and Mc-Naughton, D., 1978. The carbon isotope geochemistry of a small groundwater system in northern Ontario. *Water Resources Research*, 14: 1059-1067.
- Gautam, P., and Rösler, W., 1999. Depositional chronology and fabric of Siwalik group sediments in Central Nepal from magnetostratigraphy and magnetic anisotropy. *Journal of Asian Earth Science*, 17: 659-682.
- Gaudry, A., Polian, G., Ardouin, B. and Lambert, G., 1990. Radon-calibrated emission of CO₂ from South Africa. *Tellus*, 42B: 9-19.
- Gerhard, E., 2000. *Sedimentary Basins- Evolution, Facies and Sediment budget*. Springer-Verlag, New York.
- Ghosh, S.K. and Kumar, R., 2000. Petrography of Neogene Siwalik sandstone of the Himalayan foreland basin, Garhwal Himalaya: Implications for source area tectonics and climate. *Journal Geological Society of India*, 55: 1-15.
- Ghosh, S.K., Kumar, R. and Suresh, N., 2003. Influence of Mio-Pliocene drainage re-organization in the detrital modes of sandstone, Subathu sub-basin, Himalayan foreland basin. *Journal of Himalayan Geology*, 24: 35-46.
- Ghosh, P., Bhattacharya, S.K. and Jani, R.A., 1995. Palaeoclimate and palaeovegetation in central India during the Upper Cretaceous based on stable isotope composition of the palaeosol carbonates. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 114: 285-296.
- Ghosh, P., Ghosh, P. and Bhattacharya, S.K., 2001. CO₂ levels in the Late Palaeozoic and Mesozoic atmosphere from soil carbonate and organic matter, Satpura basin, Central India. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 170:219-236.
- Ghosh, P., Padia, J.T. and Mohindra, R., 2004. Stable isotopic studies of palaeosol sediment from Upper Siwalik of Himachal Himalaya: evidence for high monsoonal intensity during late Miocene? *Palaeogeography, Palaeoclimatology, Palaeoecology*, 206:103-114.
- Gilg, H.A. and Sheppard, S.M.F., 1996. Hydrogen isotope fractionation between kaolinite and water revisited. *Geochimica et Cosmochimica Acta*, 60:529-533.

- Graham, C.M., Atkinson, J. and Harmon, R.S., 1984. Hydrogen isotope fractionation in the system chlorite-water. NERC 6th Progress Report of Research 1981-1984, NERC Publication Series D, No. 25: p139.
- Grocke, D.R., Hesselbo, S.P. and Jenkyns, H.C., 1999. Carbon isotope composition of lower Cretaceous fossil wood: Ocean atmosphere chemistry and relation to sea-level changes. *Geology*, 27: 155-158.
- Harland, W.B., Cox, A.V., Llewellyn, P.G., Pickton, C.A.G. and Smith, A.G., 1982. A geological Time Scale, Cambridge University Press, Cambridge, UK.
- Harris, N.B., 1992. Burial diagenesis of Brent sandstone: a study of statford, hutton and Lyell fields. In: Morton, A.C., Haszeldine, R.S., Giles, M.R. and Brown, S. (eds) *Geology of the Brent Group*. Geological Society, London. Special Publications, 61: 351-376.
- Harrison, T.M., Copeland, P., Kidd, W.S.F. and Yin, A., 1992. Raising Tibet. *Science*, 255:1663-1670.
- Harrison, T.M., Yin, A. and Ryerson, J.F., 1998. Orographic evolution of the Himalaya and Tibetan Plateau. In: Crowley J.T., Burke C.K., *Tectonic boundary condition for climate resolution*. Oxford monograph on Geology and Geophysics. No.39: 39-72.
- Hisamoti, K., 1990. The sandstone petrography of the Churia (Siwalik) Group in the Arung Khola-Binai Khola Area, West Central Nepal. *Bull. Fac. Edu. Wakayama Univ., Nat. Sci.*, 39: 5-29.
- Hodell, D.A., Mueller, P.A., McKenzie, J.A and Mead, G.A., 1989. Strontium isotope stratigraphy and geochemistry of the late Neogene ocean. *Earth and Planetary Science Letters*, 92:165-178.
- Humphrey, J.D. and Ferring, C.R., 1994. Stable isotopic evidence for latest Pleistocene and Holocene climatic change in north-central Texas. *Quaternary Research*, 41: 200-213.
- Hyeong, K. and Capuano, R.M., 2000. The effect of organic matter and the H₂O₂ organic matter removal method on the δD of smectite-rich samples. *Geochimica et Cosmochimica Acta*, 64, no 22: 3829-3837.
- IAEA, 2003. <http://isohis.iaea.org>
- Indian Meteorological Department, 1970. Printed by Manager, Govt. of India Press (P.L. Wing), Nasik. pp.195-196.



- Irwin, H., Curtis, C. and Coleman, M., 1977. Isotopic evidence for source of diagenetic carbonates formed during burial of organic-rich sediments. *Nature*, 269: 209-213.
- Jackson, M.L., 1969. Soil chemical analysis-advance course: 2nd edition, published by the author, Madison, Wis., pp 895.
- Jaeger, J.J., Courtillot, V. and Tapponier, P., 1989. Paleontological view of the ages of Deccan Traps, the Cretaceous/Tertiary boundary, and the India-Asia collision. *Geology*, 17:463-469.
- Johnson, M.N., Stix, J., Tauxe, L., Cervený, P.F. and Tahirkheli, R. A. K., 1985. Paleomagnetic chronology, fluvial process and tectonic implications of the Siwalik deposits near Chinji Village, Pakistan. *J. Geol.*, 93:27-40.
- Karunakaran, C. and Rao, R.A., 1979. Status of exploration for hydrocarbons in the Himalayan region - contributions to stratigraphy and structure. In *Himalayan Geology Seminar-1976*, Geological Survey of India, Miscellaneous publications 41:1-66.
- Kelly, E.F., Amundson, R.G., Marino, B.D. and DeNiro, M.J., 1991a. Stable carbon isotopic composition of carbonate in Holocene grassland soils. *Soil Science Society of American Journal*. 55: 1651-1658.
- Kelly, E.F., Amundson, R.G., Marino, B.D. and Deniro, M.J., 1991b. Stable isotope ratios of carbon in Phytoliths as a quantitative method of monitoring vegetation and climate change. *Quaternary Research*, 35:222-233.
- Keller, W.D., 1970. Environmental aspects of clay minerals. *Journal Sedimentary Petrology*, 40: 785-813.
- Knoepp, J.D., Tieszen, L.L. and Fredlund, G.G., 1998. Assessing the vegetation history of three southern Appalachian Balds through soil organic matter analysis. Research Paper, SRS-13. U.S department of Agriculture, Forest service, Southern Research Station. 1-12.
- Kotzer, T.G. and Kyser, T.K., 1991. Retrograde alteration of clay minerals in uranium deposits: radiation catalyzed or simply low-temperature exchange? *Chemical Geology*, 86: 307-321.
- Krishnamuthy, S.V. and Bhattacharya, S.K., 1991. Stable oxygen and hydrogen isotope ratio in shallow ground waters from India and a study of the role of evapotranspiration in the Indian Monsoon. *Isotope Geochemistry: A tribute to*

- Samuel Epstein; Special publication No.3 of The Geochemical Society (ed.H.P.Taylor, J.R.O'Neil and I.R. Kaplan) pp.1-7.
- Kumar, R.,1993. Coalescence megafan: multistorey sandstone complex of the late-orogenic (Mio-Pliocene) sub-Himalayan belt, Dehra Dun, India. *Sedimentary Geology*, 85: 327-337.
- Kumar, R. and Nanda, A.C., 1989. Sedimentology of the Middle Siwalik Subgroup of Mohand Area, Dehra Dun Valley, India. *Journal Geological Society of India*, 34: 597-616.
- Kumar, R., Ghosh, S.K. and Sangode, S.J., 1999. Evolution of a Neogene fluvial system in a Himalayan foreland basin, India. *Himalaya and Tibet: Mountain Roots to Mountain Tops: Boulder, Colorado, Geological Society of America Special Paper*. Edited by Macfarlane A., Sorkhabi, R. B., Quade J. pp 239-256.
- Kumar, R., Ghosh, S.K. and Sangode, S. J., 2003a. Mio-Pliocene sedimentation history in the northwestern part of the Himalayan foreland basin, India. *Current Science*, 84, no 8: 1006-1013.
- Kumar, R., Ghosh, S. K., Mazari, R. K. and Sangode, S. J., 2003b. Tectonic impact on fluvial deposits of Plio-Pleistocene Himalayan foreland basin, India. *Sedimentary Geology*, 158: 209-234.
- Kumar, R., Sangode, S.J. and Ghosh, S.K., 2004. A Multistory sandstone complex of the Himalayan foreland basin, NW Himalaya, India. *Jour. Asian Earth Sci.*, In Press.
- Kroon, D., Steens, T. and Troelstra, S.R., 1991. Onset of monsoonal related upwelling in the western Arabian Sea as revealed by planktonic foraminifers. In: *Prell, W.L., Niitsuma, N et al., (Eds.) Proceedings of the Ocean Drilling Program, Scientific Results*, 117: 257-263.
- Land, L.S., 1980. The isotopic and trace element geochemistry of dolomite: the state of art, in Zenger, D.H., Dunham, J.B. and Ethington, R.A., eds., *Concepts and Models of dolomitization: Soc. Econ. Paleontologists, Special Paper 28*: 87-110.
- Lanord, C.F., Derry, L. and Michard, A., 1994. Evolution of the Himalaya since Miocene time: isotopic and sedimentological evidence from Bengal Fan.

- From Treloar, P.J and Searle, M.P. (eds.), Himalayan Tectonics. Geological Society Special Publication, London. No.74: 603-621.
- Leaney, F.W. and Herczeg, A.L., 1995. Regional recharge of a karst aquifer estimated from chemical and isotopic composition of diffuse and localized recharge, South Australia. *Journal of Hydrology*, 164: 363-387.
- Longstaffe, F.J. and Ayalon, A., 1990. Hydrogen-isotope geochemistry of diagenetic clay minerals from Cretaceous sandstones, Alberta, Canada: evidence of exchange. *Applied Geochemistry*, 5: 657-668.
- Lydekker, R., 1877. Vertebrata from Indian Tertiary and Secondary rocks. *Rec. Geol. Surv. India.*, 10, no.1: 34.
- Maulik, P.K., Chakraborty, C., Ghosh, P. and Rudra, D., 2000. Meso and Macro-Scale Architecture of a Triassic Fluvial Succession: Denwa Formation, Satpura Gondwana Basin, Madhya Pradesh. *Journal Geological Society of India*, 56: 489-504.
- Marumo, K., Nagasawa, K. and Kuroda, Y., 1980. Mineralogy and hydrogen isotope of clay minerals in the Ohnuma geothermal area, Northeastern Japan. *Earth and Planetary Science Letters*, 47:255-262.
- Mehra, O.P. and Jackson, M.L., 1960. Fe oxide removal from soils and clays by dithionite-citrate system buffered with sodium bicarbonate. In *Proceeding of the 7th National conference on Clays and Clay minerals* (ed. A.Swineford. Pergamon Press) pp. 317-327.
- Mehta, Y.P., Thakur, A.K., Lal, N., Shukla, B. and Tandon S.K., 1993. Fission track age of zircon separates of tuffaceous mudstones of the Upper Siwalik subgroup of Jammu-Chandigarh sector of the Panjab Sub-Himalaya. *Current Science*, 64, no.7: 519-521.
- Metivier, F., Gaudemer, Y., Tapponnier, P. and Klein, M., 1999. Mass accumulation rates in Asia during the Cenozoic. *Geophysical Journal International*, 137: 280-318.
- Milliken, K.L., 1989. Petrography and composition of authigenic feldspars, Oligocene Frio formation, South Texas. *Journal of Sedimentary Petrology*, 59: 361-374.
- Mora, C.I., Driese, S.G. and Fastovsky, D.E., 1993. Geochemistry and stable isotopes of paleosols. A geological society of America continuing education short course. Knoxville Publication, University of Tennessee.

- Mora, C.I., Driese, S.G. and Colarusso, L.A., 1996. Middle to Late Paleozoic atmospheric CO₂ levels from soil carbonate and organic matter. *Science*, 271:1105-1107.
- Morgan, M.E., Kingston, J.D. and Marino, B.D., 1994. Carbon isotopic evidence for the emergence of C₄ plants in the Neogene from Pakistan and Kenya. *Nature*, 367: 162-165.
- Mugnier, J.L., Charlaron, E., Mascle, G., Pradier, B. and Herail, G., 1995. Structural and thermal evolution of the Siwaliks of western Nepal. *Bulletin of the Geological Society, Nepal*, 11: 171-179.
- Nakayama, K. and Ulak, P.D., 1999. Evolution of fluvial style in the Siwalik Group in the foothills of the Nepal Himalaya. *Sedimentary Geology*, 125: 205-224.
- Nigrini C., 1991. Composition and biostratigraphy of radiolarian assemblages from an area of upwelling (Northwestern Arabian Sea, Leg 117). *Proceeding of the Ocean Drilling Program, Scientific Results*, 117: 89-101.
- Pagani, M., Freeman, K.H. and Arthur, M.A., 1999. Late Miocene atmospheric CO₂ concentrations and the expansion of C₄ grasses. *Science*, 285: 876-879.
- Pearson, F.J., Jr. and Hanshaw, B.B., 1970. Source of dissolve carbonate species in groundwater and their effects on ¹⁴C dating, in *Isotope Hydrology 1970: Vienna, International Atomic Energy Agency*, pp. 271-286.
- Pendall, E. and Amundson, R., 1990. The stable isotope chemistry of pedogenic carbonate in an alluvial soil from the Punjab. *Soil Science*, 149: 199-211.
- Phadtare, N.R., Kumar, R. and Ghosh, S.K., 1994. Stratigraphic palynology, floristic succession and the Tatrot/Pinjur boundary in upper Siwalik sediments of Haripur Khol area, district Sirmaur(H.P), India. *Himalayan Geology*, 15:69-82.
- Pilgrim, G.E., 1913. The correlation of the Siwaliks with mammal horizons of Europe. *Rec.Geo.surv.India*, 43:264-236.
- Pollastro, R.M., 1989. Clay minerals as geothermometers and indicators of thermal maturity-application to basin history and hydrocarbon generation. *Am Assoc Petrol Geol Bull.*, 73: 1171.
- Powers, P.M., Lillie, R.J. and Yeats, R.S., 1998. Structure and shortening of the Kangra and Dehradun re-entrants, sub-Himalaya, India. *Geological Society of America Bulletin*, 110, no.8: 1010-1027.

- Prell, L.W., Murray, W.D., Clemens, C.S. and Anderson, D.M., 1992. Evolution and variability of the Indian Ocean Summer Monsoon: Evidence from the Western Arabian Sea drilling program. Synthesis of results from scientific drilling in Indian Ocean. *Geophysical Monograph*, 70:447-469.
- Prell, L.W. and Kutzbach, J.E., 1992. Sensitivity of the Indian Monsoon to forcing parameters and implications for its evolution. *Nature*, 360: 647-652.
- Quade, J., Cerling, T.E. and Bowman, J.R., 1989. Development of Asian monsoon revealed by marked ecological shift during the latest Miocene in the northern Pakistan. *Nature*, 342:163-166.
- Quade, J., Cater, M.L.J., Ojha, P.T., Adam, J. and Harrison, M.T., 1995. Late Miocene environmental change in Nepal and the northern Indian subcontinent: Stable isotopic evidence from paleosols. *Geological Society of America Bulletin*, 107, no. 12:1381-1397.
- Quade, J. and Cerling, T. E., 1995. Expansion of C₄ grasses in the Late Miocene of Northern Pakistan: evidence from stable isotopes in paleosols. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 115: 91-116.
- Quade, J. and Roe, J.L., 1999. The stable isotope composition of early ground water cements from sandstone in paleoecological reconstruction. *Journal of Sedimentary Research*, 69, no.3: 667-674.
- Rabenhorst, M.C., Wilding, L.P. and West, L.T., 1984, Identification of pedogenic carbonates using stable carbon isotope and microfacies analyses. *Soil Science*, 48:125-132.
- Raiverman, V., Kunte, S.V. and Mukherjee, A., 1983. Basin geometry, Cenozoic sedimentation and hydrocarbon prospects in north-western Himalaya and Indog-Gangetic plains. *Pet. Asia J*, 6 (IV): 67-92.
- Raiswell, R. and Fisher, Q.J., 2000. Mudrock-hosted carbonate concretions: a review of growth mechanisms and their influence on chemical and isotopic composition. *Journal of the Geological Society, London*, 157:239-251.
- Rao, Y.P., 1976. South West monsoon. *Meteorological Monograph Synoptic*. Meteorology No 1/1976. Published by Indian Meteorological Department. Pp.8.
- Rao, Y.P., 1981. The climate of the Indian subcontinent. In *climates of southern and western Asia* (ed. K.Takahshi and H. Aarakawa): *World survey of climatology* 9: 67-182.

- Rao, R.A., Agarwal, R.P., Sharma, U.N., Bhalla, M.S. and Nanda, A.C., 1988. Magnetic polarity stratigraphy and vertebrate palaeontology of the Upper Siwalik Subgroup of Jammu Hills, India. *Journal of Geological Society of India*, 31: 361-385.
- Raymo, M.E. and Ruddiman, W.F., 1992. Tectonic forcing of late Cenozoic climate. *Nature*, 359: 117-122.
- Retallack, G.J., 2001a. *Soils of the Past*. Blackwell Science. USA. pp-185.
- Retallack, G.J., 2001b. Cenozoic expansion of grasslands and climate cooling. *The Journal of Geology*, 109:407-426.
- Retallack, G.J., 2001c. A 300-million-year record of atmospheric carbon dioxide from fossil plant cuticles. *Nature*, 411:287-290.
- Retallack, G.J., 2002. Carbon dioxide and climate over 300 Myr. *Philosophical Transactions: Mathematical, Physical and Engineering Sciences*, 360:659-673.
- Richter, F.M., Rowley, D.B. and DePaolo, D.J., 1992. Sr isotope evolution of seawater: the role of tectonics. *Earth and Planetary Science Letters*, 109: 11-23.
- Romanek, C.S., Grossman, E.L. and Morse, J.W., 1992. Carbon isotopic fractionation in synthetic aragonite and calcite: Effects of temperature and precipitation rate. *Geochimica et Cosmochimica Acta*, 56: 419-430.
- Ruddiman, W.F. and Prell, W.L., 1997. Introduction to the uplift-climate connection. In: W.F. Ruddiman (Editor), *Tectonic Uplift and Climate Change*. Plenum, New York, pp. 3-19.
- Ruddiman, W.F. and Kutzbach, J.E. 1989. Forcing of late Cenozoic northern hemisphere climate by plateau uplift in Southern Asia and the American West. *Journal of Geophysical Research* 94(D15): 18, 409-18,427.
- Salomons, W., Goudie, A. and Mook, W.G., 1978. Isotopic composition of calcrete deposit from Europe, Africa and India. *Earth Surface Process*, 3: 43-57.
- Sangode, S.J., Kumar, R. and Ghosh, S.K., 1996. Magnetic polarity stratigraphy of the Siwalik sequence Haripur area (H.P.), NW Himalaya. *Journal of Geological Society of India*, 47: 683-704.
- Sangode, S.J., Kumar, R. and Ghosh, S.K., 1999. Palaeomagnetic and rock magnetic perspectives on the post-collisional continental sediments of the Himalaya,

- India. In: Radhakrishna, T., Piper J.D.A. (Eds.), *The Indian Subcontinent and Gondwana: A Palaeomagnetic and Rock Magnetic Perspective*. Geol. Soc. India Mem., 44:221-248.
- Sangode, S.J., Bloemendal, J., Kumar, R. and Ghosh, S.K., 2001. Plio-Pleistocene pedogenic change in the Siwalik Palaeosols: A rock magnetic approach. *Current Science*, 81,no.4: 387-392.
- Sangode, S.J., Kumar, R. and Ghosh, S.K., 2003. Magnetic Polarity Stratigraphy of the Late Miocene Siwalik Group sediments from Kangra Re-entrant, H.P., India. *Himalayan Geology*, 24, no 1: 47-61.
- Sanyal, P., Bhattacharya, S.K., Kumar, R., Ghosh, S.K. and Sangode, S.J., 2004a. Mio -Pliocene monsoonal record from Himalayan Foreland basin (Indian Siwalik) and its relation to the vegetational change. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 205:23-41.
- Sanyal, P., Bhattacharya, S.K., Kumar, R., Ghosh, S.K. and Sangode, S.J., 2004b. Palaeovegetational reconstruction in Late Miocene: a case study based on early diagenetic carbonate cement from Indian Siwalik. Under revision in *Palaeogeography, Palaeoclimatology, Palaeoecology*.
- Savin, S.M. and Epstein, S., 1970. The oxygen and hydrogen isotope geochemistry of clay minerals. *Geochimica et Cosmochimica Acta*, 34:25-42.
- Savin, S.M. and Hsieh, J.C.C., 1998. The hydrogen and oxygen isotope geochemistry of pedogenic clay minerals: principles and theoretical background. *Geoderma*, 82: 227-253.
- Scotese, C.R., 1998. Gondwana's climate changes. *J. Afr. Earth Science*, 27:172-173.
- Sheppard, S.M.F. and Gilg, H.A., 1996. Stable isotope geochemistry of clay minerals. *Clay Minerals*, 31:1-24.
- Spicer, A.R., Harris, B.W.N., Widdowson, M., Herman, B.A., Guo, S., Valdes, J.P., Wolfe, A.J. and Kelley, P.S., 2003. Constant elevation of Southern Tibet over the past 15 million years. *Nature*, 421:622-624.
- Spivac, A. J., You, C.F., Smith, J.H., 1993. Foraminiferal boron isotope ratios as a proxy for surface ocean pH over the past 21 Ma. *Nature*, 363: 149-151.
- Stern, A.L., Chamberlain, R.C., Reyhols, C.R. and Johnson, D.J., 1997. Oxygen isotope evidence of climate change from pedogenic clay minerals in the Himalayan molasse. *Geochimica et Cosmochimica Acta*, 61 no.4: 731-744.

- Suresh, N., Ghosh, S.K., Kumar, R. and Sangode, S.J., 2004. Clay-mineral distribution patterns in late Neogene fluvial sediments of the Subathu sub-basin, central sector of Himalayan foreland basin: implications for provenance and climate. *Sedimentary Geology*, 163: 265-278.
- Tandon, S.K. and Kumar, R., 1984. Discovery of tuffaceous mudstones in the Pinjor Formation of Panjab sub-Himalaya, India. *Current Science*, 53:982-984.
- Tandon, S.K. and Varshney, S.K., 1991. Origin of selective carbonate cemented (concretionary) layers within multistoried sandstone bodies of the Neogene Middle Siwalik Subgroup, NW Himalaya, India. Abstract, Birbal Sahni Birth Centenary Symposium on the Siwalik Basin, WIHG, Dehra Dun, India, p.45.
- Tandon, S.K., Sood, A., Andrews, J.E. and Dennis, P.F., 1995. Palaeoenvironments of the dinosaur-bearing Lameta Beds (Maastrichtian), Narmada Valley, Central India. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 117:123-154.
- Thomas, J.V., Prakash, B. and Mahindra, R., 2002. Lithofacies and paleosol analysis of the Middle and Upper Siwalik Groups (Plio-Pleistocene), Haripur-Kolar section, Himachal Pradesh, India. *Sedimentary Geology*, 150:343-366.
- Van Der Burgh, J., Visscher, H., Dilcher, L.D. and Kurschner, M. W., 1993. Paleoatmospheric signatures in Neogene fossil leaves. *Science*, 260:1788-1790.
- Virdi, N.S., 1979. On the geodynamic significance of mega-lineaments in the Outer and Lesser regions of western Himalaya. *Himalayan Geology*, 9, no 1:79-99.
- Wang, H. and Follmer, R. L., 1998. Proxy of monsoon seasonality in carbon isotopes from paleosols of the southern Chinese Loess Plateau. *Geology*, 26, no. 11: 987-990.
- Wassenaar, L.I., Aravena, R., and Fritz, P., 1992. Radiocarbon contents of dissolved organic carbon in shallow groundwater systems. In *Isotope Techniques in Water Resources Development 1991 (Proceedings of the international Symposium)*: Vienna, International Atomic Energy Agency, pp. 143-151.
- White, N.M., Parrish, R.R., Bickle, M.J., Najman, M.R., Burbank, D. and Maithani, A., 2001. Metamorphism and exhumation of the NW Himalaya constrained by U-Th-Pb analyses of detrital monazite grains from early foreland basin sediments. *Journal Geological Society London*, 158: 625-635.

- Wilkinson, M. and Haszeldine, R.S., 1996. Aluminum loss from arkoses produces diagenetic quartzites and secondary porosity: Fulmar Formation, North Sea. *Journal of the Geological Society, London*, 154: 747-751.
- Wilson, M.J., 1987. *A Handbook of Determinative Methods in Clay Mineralogy*. Blackie, Glasgow. pp 308.
- Yeh, H., 1980. D/H ratios and late-stage dehydration of shales during burial. *Geochimica et Cosmochimica Acta*, 44: 341-352.
- Yeh, H. and Epstein, S., 1978. Hydrogen isotope exchange between clay minerals and sea water. *Geochimica et Cosmochimica Acta*, 42:140-143.
- Yin, A., Harrison, T.M., Ryerson, F.J., Chen, W., Kidd, W.S.F. and Copeland, P., 1994. Tertiary structural evolution of Gangdese thrust system, South eastern Tibet. *J. Geophys. Res.*, 99: 18,175-18,201.
- Yurtsever, Y. and Gat, J.R., 1981. Atmospheric waters. In stable isotope hydrology: Deuterium and Oxygen-18 in the Water cycle. In: J.R. Gat and R. Gonfiantini (Eds.). *Technical Report Series 210*, IAEA, Vienna. pp. 103-142.
- Zaleha, M.J., 1997. Intra- and extra-basinal controls on fluvial deposition in the Miocene Indo-Gangetic foreland basin, northern Pakistan. *Sedimentology*, 44: 369-390.
- Ziegler, K. and Longstaffe, F.J., 2000. Clay mineral authigenesis along a mid-continental scale fluid conduit in Palaeozoic sedimentary rocks from southern Ontario, Canada. *Clay Minerals*, 35: 239-260.