

LIST OF FIGURES

Figure No.	Title of the figure	Page No.
1.1	Study area map	10
1.2	Diagram of the rDNA region of fungi and ITS primers	16
1.3	Pipeline of DNA barcoding	19
2.1	Schematic illustration of therapeutic applications of macro-fungi	36
3.1	Schematic representation of PCR reaction conditions	72
3.2	Schematic representation of PCR cycle	73
3.3	Geographical map showing the collection sites of fungi	77
3.4	Gel image showing the isolated bands of DNA from different fungi	79
3.5	Gel image showing the amplified bands of ITS rDNA from different fungi after PCR	79
3.6	Gel image showing the purified amplified bands of ITS rDNA from different fungi after PCR purification step	80
3.7	Maximum likelihood cladogram generated from ITS sequence dataset for <i>Clathrus delicatus</i> and related taxa. As an outgroup sequences of <i>Trappea darkeri</i> were utilised. Bootstrap values are indicated on the generated phylogenetic tree branches	82
3.8	(1) Various stages of fruiting body development of <i>Clathrus delicatus</i> . A-C, mature fruiting bodies (receptacles); D, developing fruiting bodies ('myco-eggs' stage) arising from the thick whitish mycelial mat; E, fully grown cluster of myco-egg ready to develop into the mature fruiting body; F, cluster of myco-eggs and volva (arrowhead); G, volva and H, I, expanded receptacle. (2) Nucleotide sequence submitted to NCBI. (3) Illustrative barcode generated after submission to the BOLD database	84
3.9	Maximum likelihood cladogram generated from ITS sequence dataset for <i>Trichaleurina javanica</i> and related taxa. <i>Pseudosarcosoma latahense</i> and <i>Neournula pouchetii</i> as outgroup and the bootstrap values are indicated on the tree branches	86
3.10	(1) A-J, Morphology of <i>Trichaleurina javanica</i> . (2) Nucleotide sequence submitted to NCBI. (3) Illustrative barcode generated after	88

	submission to the BOLD database	
3.11	Maximum likelihood cladogram created from concatenated LSU and ATP6 dataset for <i>Itajahya galericulata</i> and related taxa. <i>Phallobata alba</i> and <i>Trappea darkeri</i> were taken as outgroup. Bootstrap values are shown on the tree branches	90
3.12	(1) Morphology of <i>Itajahya galericulata</i> : A, greyish white immature eggs ready for opening; B, developing fruiting body emerging from the egg; C, mature fruiting body; D, habit and morphological features in natural habitat; E, mature fruiting body; F, calyptra and gleba; G, SEM of basidiospores; H, developing immature egg and I, egg dissected to show internal features. (2) Nucleotide sequence submitted to NCBI. (3) Illustrative barcode generated after submission to the BOLD database	92
3.13	Maximum likelihood cladogram generated from ITS sequence dataset for <i>Cyathus stercoreus</i> and related taxa. <i>Cystoderma amianthinum</i> was taken as an outgroup. Bootstrap values are shown on the tree branches	94
3.14	(1) Morphology of <i>Cyathus stercoreus</i> . A, B, C, the morphology of fruiting bodies growing on dung; D, E, mycelia cultures on PDA plate; F, G, section through the longitudinal line of cup showing attached peridioles and H, I peridioles with funicular cords. (2) Nucleotide sequence submitted to NCBI. (3) Illustrative barcode generated after submission to the BOLD database	96
3.15	Maximum likelihood cladogram generated from ITS sequence dataset for <i>Geastrum sp.</i> and related taxa. <i>Myriostoma coliforme</i> was taken as an outgroup. Bootstrap values are shown on the tree branches	98
3.16	RFLP analysis of three different species of <i>Geastrum</i> using three different restriction enzymes: (A) <i>EcoRI</i> digestion, (B) <i>HinfI</i> digestion and (C) <i>TaqI</i> digestion	99
3.17	(1) Morphology of <i>Geastrum saccatum</i> . A, immature brown bulb and growing fruiting bodies; B-F, mature fruiting bodies with 5-8-pointed non-hygroscopic saccate rays; G, mature fruit body	102

	measuring 0.6-2.5 cm in diameter and H, exposed endoperidium. (2) Nucleotide sequence submitted to NCBI. (3) Illustrative barcode generated after submission to the BOLD database	
3.18	(1) Morphology of <i>Geastrum rufescens</i> . A-C, mature fruiting bodies with 5-8-pointed starfish-like rays; D-E, endoperidium with Peristome surrounding the slight elevated apical pore and F, mature fruit body measuring 3-4 cm in diameter. (2) Nucleotide sequence submitted to NCBI. (3) Illustrative barcode generated after submission to the BOLD database	104
3.19	(1) Morphology of <i>Geastrum triplex</i> . A, immature basidiomata with coarsely fibrous to the squamulose surface; B, growing basidiomata; E-F, mature fruiting bodies with 5-6 nonhygroscopic starfish-like rays forming a prominent collar around the endoperidium; C, D, G and H, conically elevated peristome surrounding the ostiole. (3) Nucleotide sequence submitted to NCBI. (4) Illustrative barcode generated after submission to the BOLD database	107
3.20	Maximum likelihood cladogram generated from ITS sequence dataset for <i>Disciseda candida</i> and related taxa. <i>Mycenastrum corium</i> and <i>Lepiota cristata</i> were taken as out group. Bootstrap values are shown on the tree branches	109
3.21	(1) Morphology of <i>Disciseda candida</i> . A-E, basidiomata; F, basidiospores observed under SEM and G, basidiospores observed under a microscope. (2) Nucleotide sequence submitted to the NCBI. (3) Illustrative barcode generated after submission to the BOLD database	111
3.22	Maximum likelihood cladogram generated from ITS sequence dataset for <i>Scleroderma bovista</i> and related taxa. <i>Pisolithus albus</i> and <i>Pisolithus tinctorius</i> were taken as outgroup. Bootstrap values are indicated on the tree branches	113
3.23	(1) A-G, Morphology of <i>Scleroderma bovista</i> . (2) Nucleotide sequence submitted to NCBI. (3) Illustrative barcode generated after submission to the BOLD database	115
3.24	Maximum likelihood cladogram generated from ITS sequence	117

	dataset for <i>Pisolithus sp.</i> and related taxa. <i>Suillus luteus</i> and <i>Paxillus involutus</i> were taken as outgroup. Bootstrap values are indicated on the tree branches	
3.25	(1) Morphology of <i>Pisolithus albus</i> . A-D, fruit body and E-F, basidiospores under microscope (2) Nucleotide sequence submitted to the NCBI. (3) Illustrative barcode generated after submission to the BOLD database	119
3.26	(1) Morphology of <i>Pisolithus tinctorius</i> . A-B, fruit body and C-D, basidiospores (2) Nucleotide sequence submitted to NCBI. (3) Illustrative barcode generated after submission to the BOLD database	121
3.27	Maximum likelihood cladogram generated from ITS sequence dataset for <i>Dictyophora sp.</i> and related taxa. <i>Gelopellis sp.</i> , <i>Phallobata alba</i> and <i>Trappea darkeri</i> were taken as out group. Bootstrap values are shown on the tree branches	123
3.28	(1) A-H, Morphology of <i>Dictyophora indusiata</i> . (2) Nucleotide sequence submitted to the NCBI. (3) Illustrative barcode generated after submission to the BOLD database	126
3.29	(1) A-E, Morphology of <i>Dictyophora multicolor</i> . (2) Nucleotide sequence submitted to NCBI. (3) Illustrative barcode generated after submission to the BOLD database	128
4.1	Antioxidant potentials of different fungal crude extracts at different concentration against DPPH free radicals. (a) <i>P. albus</i> (b) <i>P. tinctorius</i> (c) <i>G. triplex</i> (d) <i>G. saccatum</i> (e) <i>D. indusiata</i> (f) <i>I. galariculata</i> (g) <i>C. stercoreus</i>	148
4.2	Antioxidant potentials of different fungal crude extracts at different concentration against H ₂ O ₂ . (a) <i>P. albus</i> (b) <i>P. tinctorius</i> (c) <i>G. triplex</i> (d) <i>G. saccatum</i> (e) <i>D. indusiata</i> (f) <i>I. galariculata</i> and (g) <i>C. stercoreus</i>	152
4.3	Anticancer potentials of different fungal crude extracts at different concentration on various cancerous cell lines. (a) <i>P. albus</i> (b) <i>P. tinctorius</i> (c) <i>G. triplex</i> (d) <i>G. saccatum</i> (e) <i>D. indusiata</i> (f) <i>I. galariculata</i> and (g) <i>C. stercoreus</i>	156

4.4	HR-LC/MS analysis of crude extract of <i>P. albus</i> in +ve and -ve mode	157
4.5	HR-LC/MS analysis of crude extract of <i>G. saccatum</i> in +ve and -ve mode	158
4.6	HR-LC/MS analysis of crude extract of <i>D. indusiata</i> in +ve and -ve mode	159
4.7	HR-LC/MS analysis of crude extract of <i>I. galericulata</i> in +ve and -ve mode	160
4.8	HR-LC/MS analysis of crude extract of <i>C. stercoreus</i> in +ve and -ve mode	161
4.9	Interaction of N3 in the binding cleft of SARS-CoV-2 M ^{pro} (A) hydrophobicity surface 3D representation (B) Interaction of N3 in a pocket site of SARS-CoV-2 M ^{pro} (C) the crystal structure of SARS-CoV-2 M ^{pro} in complex with an inhibitor N3 (D) 2D representation describing all possible bindings in active site of SARS-CoV-2 M ^{pro}	176
4.10	Docking analysis visualization of SARS-CoV-2 M ^{pro} binding with Bergenin (A) hydrophobicity surface 3D representation (B) Interaction of Bergenin in a pocket site of SARS-CoV-2 M ^{pro} (C) 3D representation of Bergenin-SARS-CoV-2 M ^{pro} interaction (D) 2D representation describing bindings of Bergenin in active site of SARS-CoV-2 M ^{pro}	181
4.11	Docking analysis visualization of SARS-CoV-2 M ^{pro} binding with Quercitrin (A) hydrophobicity surface 3D representation (B) Interaction of Quercitrin in a pocket site of SARS-CoV-2 M ^{pro} (C) 3D representation of Quercitrin-SARS-CoV-2 M ^{pro} interaction (D) 2D representation describing bindings of Quercitrin in active site of SARS-CoV-2 M ^{pro}	182
4.12	Docking analysis visualization of SARS-CoV-2 M ^{pro} binding with Dihydroartemisinin (A) hydrophobicity surface 3D representation (B) Interaction of Dihydroartemisinin in a pocket site of SARS-CoV-2 M ^{pro} (C) 3D representation of Dihydroartemisinin-SARS-CoV-2 M ^{pro} interaction (D) 2D representation describing bindings of Dihydroartemisinin in active site of SARS-CoV-2 M ^{pro}	183

4.13	Interactions of compounds with H-bond and hydrophobic in a pocket site of SARS-CoV-2 M ^{pro} (A), (C) and (E) show H-bond interaction of Bergenin, Quercitrin and Dihydroartemisinin; (B), (D) and (F) show hydrophobic interaction of Bergenin, Quercitrin and Dihydroartemisinin	184
------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----
