

List of Figures

Fig 2.01	Relationship between the six principles and adolescents nutritional status	31
Fig 3.01	Study Design	48
Fig 4.01	Mean Height as Percent Standards of the Adolescents (Boys + Girls) at the Beginning and After 12 months of the Study (N = 692)	69
Fig 4.02	Comparison of Mean Height for Age of 10 - 18 Years Old Study boys with the NCHS Standards and Other Indian Studies	72
Fig 4.03	Mean Increment in Height of Adolescent Boys (10 - 18 Years) During Study Period	72
Fig 4.04	Comparison of Mean Height for Age of 10 - 18 Years Old Study Girls with the NCHS Standards and Other Indian Studies	73
Fig 4.05	Mean Increment in Height of Adolescent Girls (10 - 18 Years) During Study Period	73
Fig 4.06	Comparison of Mean Height Velocity of the Study Boys Over a 12 Month Period at Each Age Compared with the NCHS Standards (N = 330)	76
Fig 4.07	Comparison of Mean Height Velocity of the Study Girls Over a 12 Month Period at Each Age Compared with the NCHS Standards (N = 362)	76
Fig 4.08	Mean Weight as Percent Standards of the Adolescents (Boys + Girls) at the Beginning and After 12 months of the Study (N = 692)	80
Fig 4.09	Mean Increment in Weight of Adolescent Boys (10 - 18 Years) During Study Period	83
Fig 4.10	Mean Increment in Weight of Adolescent Girls (10 - 18 years) During Study Period	83

		Pg No
Fig 4.11	Comparison of Mean Weight Velocity of the Study Boys Over a 12 Month Period at Each Age with the NCHS Standards (N = 330)	85
Fig 4.12	Comparison of Mean Weight Velocity of the Study Girls Over a 12 Month Period at Each Age with the NCHS Standards (N = 362)	85
Fig 4.13	Percent Prevalence of Under Nutrition in Boys (10 – 18 years) by Waterlow's Qualitative Classification at the Beginning and After 12 months of the Study	90
Fig 4 14:	Percent Prevalence of Under Nutrition in Girls (10 – 18 years) by Waterlow's Qualitative Classification at the Beginning and After 12 months of the Study	91
Fig 4.15	Percent Prevalence of Anaemia among Adolescent Boys (10 - 18 Years) at the Beginning of the Study	105
Fig 4.16	Percent Prevalence of Anaemia among Adolescent Boys (10 - 18 Years) after 12 months of the Study	105
Fig 4 17	Percent Prevalence of Anaemia among Adolescent girls (10 - 18 Years) at the Beginning of the Study	106
Fig 4.18	Percent Prevalence of Anaemia among Adolescent girls (10 - 18 Years) after 12 months of the Study	106
Fig 4 19	Nutrient Intake (as % RDA) of Adolescent Boy in An Average Household in Three Different Seasons During Study Period	131
Fig 4.20	Nutrient Intake (as % RDA) of Adolescent Girl in An Average Household in Three Different Seasons During Study Period	131
Fig 4.21	Frequency Distribution of Calorie Intake (as % RDA) of Adolescent Boys (10 - 18 Years) by BMI Status	149
Fig 4.22	Frequency Distribution of Calorie Intake (as % RDA) of Adolescent girls (10 - 18 Years) by BMI Status	149