LIST OF TABLES

Sr. No.	Title of tables			
1.	Introduction			
1	Effect of iodine deficiency by life stage			
2	Countries (number) by iodine status over the period 2003-2011	10		
3	Comparison of country data on iodine nutrition	11		
2.	Review of literature			
1	Etiology of hypothyroidism in pregnancy	32		
2	Etiology of hyperthyroidism in pregnancy	32		
3	Etiology of postpartum thyroid dysfunction	. 32		
4	Thyroid function during pregnancy	36		
5	lodothyronine concentrations in maternal and fetal serum and amniotic fluid	52		
6	Worldwide trimester specific reference intervals [percentiles- 5^{th} & 95^{th} , 2.5^{th} & 97.5^{th}]	66		
7	Worldwide trimester specific reference interval [mean/median]	69		
8	Pregnancy outcome associated with hypothyroidism: maternal aspects	71		
9	Pregnancy outcome associated with hypothyroidism: fetal & neonatal aspects	72		
10	lodine requirements during pregnancy, lactation and neonatal period	87		
11	Epidemiological criteria from the WHO for assessment of iodine nutrition in a population based on median or range of UI	89		
12	Neuropsychiatric and intellectual deficits in infants and schoolchildren born to mothers residing in conditions with mild to moderate iodine deficiency	93		
13	Screening test items (BDSTI)	96		
14	Iron requirements in pregnancy	100		
15	Comparative Physico-Chemical Features of different DFS Formulations	106		
16	Comparative bio-impact features of different DFS formula	107		
17	Responses from various Public health experts/ scientists towards Double fortification of salt	110		
3.	Materials and methods			
1.	BMI reference for adults	124		
2	Cut-offs used for classifying nutritional status of infants based on WHO reference standards (2006)	125		
3	Epidemiological criteria from the WHO for assessment of iodine nutrition in a population based on median or range of UI	130		
4	Hemoglobin cut-offs to define iron deficiency anemia during pregnancy	151		

Sr. No.	Title of tables	Page No.
5	Phatak's Screening Test items and age-grouping	152
6	Screening efficiency of BDSTI	153
	Results and discussion	
4.1.	Phase I	
1	General characteristics of pregnant women	165
2	Time trends in dietary intake (g/day) in pregnant and lactating women	168
3	Time trends in nutrient intake in pregnant and lactating women	169
4	Change in weight during pregnancy	170
5	Birth weight and socio economic status	170
6	Prevalence of anemia among pregnant women in India and other Asian countries	174
7	Hemoglobin level of pregnant women	175
8	Summary of studies on lodine nutrition status of pregnant women in India	177
9	Urinary iodine level of pregnant women	180
10	Thyroid Hormone level of pregnant women	182
4.2.	Phase II	
1	Anthropometric measurements of pregnant women	189
2 ·	Mean difference in BMI (kg/m²) from 3 rd to 9 th month	190
3	Nutritional Status during initial and final stages of pregnancy	190
4	Mean and 95 % CI for hemoglobin (g/dl) during each trimester	193
5	Repeated measures of ANOVA, trend analysis and post hoc test for hemoglobin	194
6	Prevalence (%) of IDA during each trimester	195
7	Median Urinary lodine (µg/L) during pregnancy	197
8	Non parametric test (Friedman) for urinary iodine during all 3 trimesters	199
9	lodine intake of pregnant women	200
10	Mean and median thyroid hormone of pregnant women	203
11	Thyroid dysfunction among pregnant women	207
12	Repeated measures of ANOVA for logTSH	212
13	Repeated measures of ANOVA for FT4	213
14	Repeated measures of ANOVA for TT4	215
15	Repeated measures of ANOVA for TG	216
16	Trimester specific reference intervals for TSH (µIU/mI) and FT4 (ng/dI)	217
17	Comparison of trimester specific reference intervals	218
18	Distribution of subjects according to 3 different reference intervals	221

Sr. No.	Title of tables	Page No.
19	Knowledge of pregnant women regarding iodine and iron	229
20	Maternal health indicators	238
21	Performance indicators for maternal health services	239
4.3.	Phase III	
1	Characteristics of neonates	242
2	Gestational age at birth and Anthropometric measurements of neonates	242
3	Bio-chemical profile of mothers during pregnancy	243
4	Thyroid hormone level of neonates	244
4.4.	Phase IV	
1	Characteristic of women during postpartum (6 months)	251
2	Comparison of maternal dietary intake	252
3	Prevalence of IDA and ID	254
4	Comparison of thyroid dysfunction	256
5	Comparison of 2 diagnostic methods for hypothyroidism	257
6	Comparison of biochemical parameters of subjects during pregnancy (I, II and III trimester) and postpartum	258
7	Characteristic of infants	263
8	Nutritional status of infants (at six and twelve months)	264
9	Comparison of maternal and infant characteristics in 2 groups	266
10	Prevalence of IDA and ID before and after 6 months supplementation of DFS	267
11	Mean BDSTI scores, anthropometric measurements and thyroid hormones in both groups	272

,