

# LIST OF TABLES

<u>TABLE NO.</u>	<u>TITLE</u>	<u>PAGE NO.</u>
1.	Reactions with further attenuated live measles vaccine	46
2.	Seroconversion rates (%) by age at vaccination (other countries)	54
3.	Seroconversion rates (%) by age at vaccination (Indian studies)	58
4.	Percent distribution of children (0-6 years) according to age and sex at baseline	101
5.	Percent distribution of children (0-6 years) according to age and vaccination coverage for measles and BCG	102
6.	Association between sex and nutritional status of children (0-6 years) at baseline	103
7.	Percent distribution of children (0-6 years) with measles in relation to their age and sex at baseline	104
8.	Percent distribution of children (0-6 years) with measles in relation to their age and nutritional status at baseline	107
9.	Mean age of onset of measles in children (0-6 years) stratified for age and sex and its comparison in various age groups at baseline	108
10.	Beliefs and perceptions of respondents regarding causation of measles at baseline	110
11.	Rituals observed by families during measles at follow up	111
12.	Complications of measles observed in children (0-6 years) at follow up	113
13.	Distribution of complications of measles as per nutritional status at baseline observed in children (0-6 years) recorded at follow up	115
14.	Percent distribution of children (0-6 years) suffering from measles stratified for nutritional status at baseline and at follow up	116

<b><u>TABLE NO.</u></b>	<b><u>TITLE</u></b>	<b><u>PAGE NO.</u></b>
15.	Mean change in weight of children (0-6 years) from baseline to follow up after an episode of measles	118
16.	Percent distribution of and time interval taken for medical treatment by children (0-6 years) suffering from measles recorded at follow up	119
17.	Percent distribution of newborns in relation to sex as per their birth weight and maturity	122
18.	Percent distribution of and correlation between maternal and cord blood measles HAI antibody titres	124
19.	Comparison between maternal and cord blood measles HAI antibody titres (GMT) in relation to the maturity and birth weight of newborns	126
20.	Comparison of cord blood passive measles HAI antibody titres (GMT) in relation to sex, maturity and birth weight and maturity of newborns	127
21.	Percent distribution of children (0-72 months) in relation to their age and sex	130
22.	Percent distribution of passive measles HAI antibody titres in relation to age in children (0-72 months)	132
23.	Percent distribution of and association between the presence of passive measles HAI antibody and age in children (0-72 months)	133
24.	Association between passive measles HAI antibody titres (GMT) and age in children (0-72 months)	134
25.	Percent distribution of and association between the presence of passive measles HAI antibody and nutritional status in children (6-72 months)	138
26.	Association between passive measles HAI antibody titres (GMT) and nutritional status in children (6-72 months)	139
27.	Percent distribution and association between the presence of passive measles HAI antibody and sex in children (0-72 months)	140

<u>TABLE NO.</u>	<u>TITLE</u>	<u>PAGE NO.</u>
28.	Association between passive measles HAI antibody titres (GMT) and sex in children (0-72 months)	142
29.	Percent distribution of children (6-72 months) in relation to their age and sex	145
30.	Percent distribution of prevaccination measles HAI antibody titres according to age at vaccination in children (6-72 months)	146
31.	Percent distribution of postvaccination measles HAI antibody titres according to age at vaccination in children (6-72 months)	147
32.	Percent distribution according to prevaccination measles HAI antibody titres and age at vaccination in children (6-72 months) who seroconverted (four-fold rise)	149
33.	Percent distribution according to age at vaccination and postvaccination measles HAI antibody titres in children (6-72 months) who seroconverted (four-fold rise).	150
34.	Percent distribution according to pre- and post-vaccination measles HAI antibody titres in children (6-72 months) who seroconverted (four-fold rise)	151
35.	Association between seroconversion (four-fold rise) and age at vaccination in children (6-72 months)	152
36.	Association between seroconversion (four-fold rise) and prevaccination titre in children (6-72 months)	155
37.	Association between seroconversion (four-fold rise) and nutritional status in children (6-72 months)	157
38.	Association between seroconversion (four-fold rise) and sex in children (6-72 months)	159
39.	Percent distribution according to postvaccination rise in measles HAI antibody titres (GMT) and age at vaccination in children (6-72 months) who seroconverted (four-fold rise)	160
40.	Distribution of and association between mean post-vaccination rise in measles HAI antibody titres (GMT) and age at vaccination in children (6-72 months) who seroconverted (four-fold rise)	162

<b><u>TABLE NO.</u></b>	<b><u>TITLE</u></b>	<b><u>PAGE NO.</u></b>
41.	Association between postvaccination rise in measles HAI antibody titres (GMT) and prevaccination titres in children (6-72 months) who seroconverted (four-fold rise)	163
42.	Association between postvaccination rise in measles HAI antibody titres (GMT) and nutritional status in children (6-72 months) who seroconverted (four-fold rise)	165
43.	Association between postvaccination rise in measles HAI antibody titres (GMT) and sex in children (6-72 months) who seroconverted (four-fold rise)	166