

REFERENCES

REFERENCES

- Ahokas, J.T., Saarni H., Nehert D.W. and Pelkonen O. (1979) The in vitro metabolism and covalent binding of benzo(a)pyrene to DNA catalysed by trout liver microsomes. *Chem.-Biol. Int.*, 25, 103-111.
- Alexandrov K., Vendrely C. and Vendrely R. (1970) A comparative study of the action of carcinogenic substances on the RNA synthesis in mouse skin. *Canc. Res.*, 30, 1192-1196.
- Alexandrov K. and Vendrely C. (1972) Action de condensats de fumee du tabac sur la synthese de RNA dans la peau des souris. *Chem. -Biol. Int.*, 4, 155-161.
- Alexandrov K., Brookes P., King H.W.S., Osborne M.R. and Thompson M.H. (1976) Comparison of metabolism of benzo(a)pyrene and binding to DNA caused by rat liver nuclei and microsomes. *ibid*, 12, 269-277.
- Ames B.N. (1983) Dietary carcinogens and anticarcinogens, *Science*, 221, 1256-1264.
- Anderson L.M., Priest L.J., Deschner E.E. and Bundin J.M. (1983) Carcinogenic effects of intracolonic benzo(a)pyrene in β -naphthoflavone induced mice. *Canc. Lett.*, 20, 117-124.
- Anderson M.W., Boroujerdi M. and Wilson A.G.E. (1981) Inhibition in vivo of formation of adducts between metabolites of benzo(a)pyrene and DNA by butylated hydroxyanisole. *Canc. Res.*, 41, 4309-4315.
- Anderson T.W. (1975) Large scale trials of vitamin C. *Ann. N.Y. Acad. Sci.*, 258, 498-503.
- Arrand J.E. and Murray A.M. (1982) Benzpyrene groups bind preferentially to the DNA of active chromatin in human lung cells. *Nucl. Acid Res.*, 10, 1547-1555.
- Aszalos A. and Berdy J. (1981) List of known natural anti-tumor compounds, according to chemical types. In: *Antitumor Compounds of Natural Origin: Chemistry and Biochemistry* (Ed. Aszalos A.), vol. I, pp. 1-78, CRC Press, Boca Raton, Florida.
- Austin G.E. and Moyer G.H. (1979) Inhibition of RNA synthesis by derivatives of carcinogen 2-acetylaminofluorene. *Proc. Soc. Exptl. Biol. Med.*, 161, 220-224.

Autrup H., Harris C.C., Trump B.F. and Jeffrey A.M. (1978) Metabolism of benzo(a)pyrene and identification of the major benzo(a)pyrene-DNA adducts in cultured human colon. *Canc. Res.*, 38, 3689-3696,

Aviv H. and Leder P. (1972) Purification of biologically active globin messenger RNA by chromatography on oligothymidylic acid cellulose. *Proc. Natl. Acad. Sci.*, 69, 1408-1412.

Baer-Dubowska W. and Alexandrov K. (1981) The binding of benzo(a)pyrene to mouse and rat skin DNA. *Canc. Lett.*, 13, 47-52.

Baer-Dubowska W. and Alexandrov K. (1985) The binding of benzo(a)pyrene metabolites to the DNA of isolated rat liver nuclei and nuclear matrix. *Biochem. Pharmacol.*, 34, 385-386.

Baird W.M., Harvey R.G. and Brookes P. (1975) Comparison of the cellular DNA-bound products of benzo(a)pyrene with benzo(a)pyrene-4,5-oxide. *Canc. Res.*, 35, 54-57.

Baird W.M. (1979) The use of radioactive carcinogen to detect DNA modifications. In: *Chemical Carcinogens and DNA* (Ed. Grover P.L.), vol. I, pp. 59-83, CRC Press Inc., Florida.

Basu T.K., Chan U. and Fields A. (1984) Vitamin A (retinol) and epithelial cancer in man. In: *Vitamins, Nutrition and Cancer*. (Ed. Prasad K.N.) pp 33-45, S Karger A G. Basel.

Benedict W.F., Wheatly W.L. and Jones P.A. (1980) Inhibition of chemically induced morphological transformation and reversion of the transformed phenotype by ascorbic acid in C3H/10T1/2 cells. *Canc. Res.*, 40, 2796-2801.

Ben-Itzhak J., Levi B.-Z., Shor R., Lanir A., Bassan H.M. and Ulitzur S. (1985) The formation of genotoxic metabolites of benzo(a)pyrene by the isolated perfused rat liver, as detected by the bioluminescence test. *Mut. Res.*, 147, 107-112.

Bernardi G. (1971) Chromatography of nucleic acids on hydroxyapatite columns. *Methods in Enzymology*, 21, 95-139.

Berry D.L., Zachariah P.K., Slaga T.J. and Jachau M.R. (1977) Analysis of biotransformation of benzo(a)pyrene in human foetal and placental tissues with HPLC. *Eur. J. Canc.*, 13, 667-676.

- Bevan D.R. and Manger W.E. (1985) Effect of particulates on metabolism and mutagenicity of benzo(a)pyrene. *Chem.-Biol. Inter.*, 56, 13-28.
- Bhattacharya R.K. (1980) Factors controlling carcinogenic responses to environmental chemicals. In: *Management of Environment* (Ed. Patel B.) pp. 193-202. Wiley Eastern, New Delhi.
- Bhattacharya R.K., Firozi P.F. and Aboobaker V.S. (1984) Factors modulating the formation of DNA adduct by aflatoxin B₁ in vitro. *Carcinogenesis*, 5, 1359-1362.
- Bhattacharya R.K. (1985) Interaction of certain dietary factors with chemical carcinogens. In: *Proc. Nutr. Soc. (India) XVII, Annual Meeting*, in press.
- Birt D.F., Walker B., Tibbels M.G. and Bresnick E. (1986) Antimutagenesis and anti-promotion by apigenin, robinetin and indole-3-carbinol. *Carcinogenesis*, 7, 959-963.
- Bishop J.M. (1985) Trends in oncogenes. *Trends in genetics*, 1, 245-249.
- Biswas B.B., Ganguly A. and Das A. (1975) Eukaryotic RNA polymerases and the factors that control them. *Prog. N.A. Res. Molec. Biol.*, 15, 145-184.
- Bjeldanes L.F. and Chang G.W. (1977) Mutagenic activity of quercetin and related compounds. *Science*, 197, 577-578.
- Blanton R.H., Lyte M., Myres M.J. and Bick P.H. (1986) Immunomodulation by polycyclic aromatic hydrocarbons in mice and murine cells. *Canc. Res.*, 46, 2735-2739.
- Blatti S.P., Ingles C.J., Lindell T.J., Morris P.W., Weaver R.F., Weinberg J. and Rutter W.J. (1970) Structure and regulatory properties of eucaryotic RNA polymerase. *Cold Spr. Harb. Symp. Quant. Biol.*, 35, 649-657.
- Blobel G. and Potter V.R. (1966) Nuclei from rat liver: Isolation method that combines purity with high yield. *Science*, 154, 1662-1665.
- Bollum F.J. (1968) Filter paper disk techniques for assaying radioactive macromolecules. *Methods in Enzymology*, 12(B), 169-173.
- Borgen A., Darvey H., Castagnoli N., Crocker T.T., Rasmussen R.E. and Wang T.Y. (1974) Metabolic conversion of benzo(a)pyrene by Syrian hamster liver microsomes and binding of metabolites to deoxyribonucleic acid. *J. Med. Chem.*, 16, 502-506.

- Boroujerdi M., Kung H.-C., Wilson A.G.E. and Anderson M.W. (1981) Metabolism and DNA binding of benzo(a)pyrene in vivo in the rat. *Canc. Res.*, 41, 951-957.
- Bowden G.T., McGovern V., Ossanna N. and Rosenthal H. (1982) Concentration dependent alterations of DNA replication initiation and elongation by benzo(a)pyrene diol epoxide. *Carcinogenesis*, 3, 473-480.
- Boyd J.N., Sell S. and Stoesswand G.S. (1979) Inhibition of aflatoxin-induced serum α -fetoprotein in rats fed cauliflower. *Proc. Soc. Exptl. Biol. Med.*, 161, 473-475.
- Boyland E., Busby E.R., Dukes C.E., Grover P.L. and Manson D. (1964) Further experiments on implantation of materials into urinary bladder of mice. *Brit. J. Canc.*, 18, 575-581.
- Bradford M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. *Anal. Biochem.*, 72, 248-254.
- Bresnick E. (1966) Ribonucleic acid polymerase activity in liver nuclei from rats pretreated with 3-methylcholanthrene. *Mol. Pharmacol.*, 2, 406-410.
- Brookes P. and Lawley P.D. (1964) Evidence for binding of polynuclear aromatic hydrocarbons to the nucleic acids of mouse skin: Relation between carcinogenic power of hydrocarbons and their binding to deoxyribonucleic acid. *Nature*, 202, 781-784.
- Brookes P., Jones P. and Amos J. (1975) The nature of the deoxyribonucleosides involved in the binding of carcinogenic hydrocarbons to the DNA of mouse embryo cells. *Int. J. Canc.*, 15, 912-917.
- Brown J.P. (1980) A review of the genetic effects of naturally occurring flavonoids, anthroquinones and related compounds. *Mutat. Res.*, 75, 243-277.
- Brunette D.M. and Katz M. (1975) The interaction of benzo(a)pyrene with cell membranes: Uptake into Chinese hamster ovary cells and fluorescence studies with isolated membranes. *Chem.-Biol. Int.*, 11, 1-14.
- Bryan G.T. and Pamukcu A.M. (1982) Sources of carcinogens and mutagens in edible plants: Production of urinary bladder and intestinal tumors by bracken fern (*Pteridium aquilium*). In: *Carcinogens and Mutagens in the Environment*, vol. 1, Food Products (Ed. Stich H.F.), pp. 75-82, CRC Press Inc., Boca Raton, Florida.

Buening M.K., Wislocki P.G., Levin W., Yagi H., Thakker D.R., Akagi H., Koreeda M., Jerina D.M. and Conney A.H. (1978) Tumorigenicity of optical enantiomers of diastereomeric benzo(a)pyrene-7,8-diol-9,10-epoxide in newborn mice: Exceptional activity of (+)-7 β -8 α -dihydroxy-9 α ,10 α -epoxy-7,8,9,10-tetrahydrobenzo(a)pyrene. Proc. Natl. Acad. Sci., 75, 5358-5361.

Burch H.B. (1961) Methods for detecting and evaluating ascorbic acid deficiency in man and animals. Ann. N.Y. Acad. Sci., 92, 268-276.

Burke M.D., Vadi H., Jernstrom B. and Orrenius S. (1977) Metabolism of benzo(a)pyrene with isolated hepatocytes and the formation and degradation of DNA-binding derivatives. J. Biol. Chem., 252, 6424-6431.

Burton K. (1956) A study of the conditions and mechanism of diphenylamine reaction for the colorimetric estimation of deoxyribonucleic acid. Biochem. J., 62, 315-323.

Buty S.G., Thompson S. and Slaga T.J. (1976) The role of epidermal aryl hydrocarbon hydroxylase in the covalent binding of PAH to DNA and its relationship to tumor initiation. Biochem. Biophys. Res. Comm., 70, 1102-1108.

Cardesa A., Mirvish S.S., Haven G.T. and Shubik P. (1974) Inhibitory effect of ascorbic acid on acute toxicity of dimethylamine plus nitrite in the rat. Proc. Soc. Exptl. Biol. Med., 145, 124-128.

Carver J.H., Carrano A.V. and MacGregor J.T. (1983) Genetic effects of the flavonols quercetin, kaempferol and galangin on Chinese hamster ovary (CHO) cells in vitro. Mutat. Res., 113, 45-60.

Cavalieri E. and Rogan E. (1985) Role of radical cations in aromatic hydrocarbon carcinogenesis. Environ. Hlth. Perspect., 64, 69-84.

Cedar H. and Felsenfeld G. (1973) Transcription of chromatin in vitro. J. Molec. Biol., 77, 237-254.

Cerioti G. (1952) A microchemical determination of deoxyribonucleic acid. J. Biol. Chem., 198, 297-303.

Cerioti G. (1955) Determination of nucleic acids in animal tissues. J. Biol. Chem., 214, 59-70.

Cerutti P.A., Sessions F., Hariharan P.V. and Lusby A. (1978) Repair of DNA damage induced by benzo(a)pyrene diol-epoxides I and II in human alveolar tumor cells. Canc. Res., 38, 2118-2124.

- Chambon P. (1975) Eukaryotic nuclear RNA polymerases. *Ann. Rev. Biochem.*, 44, 613-638.
- Chan W.C. and Fong Y.Y. (1977) Ascorbic acid prevents liver tumor production by aminopyrine and nitrite in the rat. *Int. J. Canc.*, 20, 268-270.
- Chang G.T., Harvey R.G., Hsu W.-T. and Weiss S.B. (1979) Inactivation of SV40 replication by derivatives of benzo(a)pyrene. *Biochem. Biophys. Res. Comm.*, 88, 688-695.
- Chen F.-M. (1985) Covalent binding of benzo(a)pyrene diol epoxides to polynucleotides. *Biochemistry*, 24, 5045-5052.
- Chiha P.A., Clarke R.H. and Stratton R.D. (1975) Optically detected zero field magnetic resonance studies of proflavine-DNA and 3,4-benzpyrene-DNA complexes. *Biochem. Biophys. Res. Comm.*, 67, 403-406.
- Chipman J.K., Frost G.S., Hiron P.C. and Millburn P. (1981) Biliary excretion, systemic availability and reactivity of metabolites following intraportal infusion of (³H)benzo(a)pyrene in the rat. *Carcinogenesis*, 2, 741-745.
- Church R.B., Luther S.W. and McCarthy B.J. (1969) RNA synthesis in Taper hepatoma and mouse-liver cells. *Biochem. Biophys. Acta*, 190, 30-37.
- Clawson G.A. and Smuckler E.A. (1982) Increased amounts of double-stranded RNA in the cytoplasm of rat liver following treatment with carcinogens. *Canc. Res.*, 42, 3228-3231.
- Clifford J.I. and Rees K.R. (1966) Aflatoxin: A site of action in the rat liver cell. *Nature*, 209, 312-313.
- Cohen G.M., Haws S.M., Moore B.P. and Bridges J.W. (1976) Benzo(a)pyrene-3-yl hydrogen sulfate, a major ethylacetate-extractable metabolite of benzo(a)pyrene in human, hamster and rat lung cultures. *Biochem. Pharmacol.*, 25, 2561-2570.
- Cohen G.M., Uotila P., Hartiala J., Suolinna E.M., Simberg N. and Pelkonen O. (1977) Metabolism and covalent binding of (³H)benzo(a)pyrene by isolated perfused lungs and short term tracheal organ culture of cigarette smoke-exposed rats. *Canc. Res.*, 37, 2147-2155.
- Conney A.H., Miller E.C. and Miller J.A. (1957) Substrate-induced synthesis and other properties of benzo(a)pyrene hydroxylase in rat liver. *J. Biol. Chem.*, 228, 753-766.

- Cook J.W., Hewett C.L. and Hieger I. (1932) Coal tar constituents and cancer. *Nature*, 130, 926.
- Idem (1933) The isolation of a cancer-producing hydrocarbon from coal tar, Part II: Isolation of 1:2- and 4:5-benzopyrenes, perylene and 1:2-benzanthracene. *J. Chem. Soc.*, 396-398.
- Das M., Srivastava S.P., Seth P.K. and Mukhtar H. (1985) Brain microsomal enzyme mediated covalent binding of benzo(a)pyrene to DNA. *Canc. Lett.*, 25, 343-350.
- deAngelo A.B., Hudgins W.R. and Kerby S.B. (1978) 7,12-Dimethylbenzanthracene-induced alteration of mammary epithelial cell. RNA-synthetic patterns of rat strains of high and low tumor incidence. *Canc. Res.*, 38, 384-389.
- Delclos K.B. and Blumberg P.M. (1982) Identification of ascorbic acid as the heat stable factor for brain which inactivates the phorbol ester receptors. *ibid*, 42, 1227-1232.
- DelTito (Jr) B.J., Mukhtar H. and Bickers D.R. (1983) Inhibition of epidermal metabolism and DNA binding of benzo(a)pyrene by ellagic acid. *Biochem. Biophys. Res. Comm.*, 114, 388-394.
- Diamond L., McFall R., Miller J. and Gelboin H.V. (1972) The effects of two isomeric benzoflavones on aryl hydrocarbon hydroxylase and the toxicity and carcinogenicity of polycyclic hydrocarbons. *Canc. Res.*, 32, 731-736.
- DiGiovanni J., Decina P.C., Prichette W.P., Fisher E.P. and Aalfs K.K. (1985) Formation and disappearance of benzo(a)pyrene DNA-adducts in mouse epidermis. *Carcinogenesis*, 6, 741-747.
- Dixit R., Teel, R.W. and Stoner G.D. (1984) Inhibition of DNA binding and metabolism of benzo(a)pyrene and benzo(a)pyrene-7,8-diol in cultured strain A mouse lung tissues by ellagic acid. *Proc. Amer. Assoc. Canc. Res.*, 25, 93.
- Dock, L., Waern F., Martinez M., Grover P.L. and Jernstrom B. (1986) Studies on the further activation of benzo(a)pyrene diol epoxide by rat liver microsomes and nucleic acids. *Chem.-Biol. Int.*, 58, 301-318.
- Doll R. (1971) Introduction. In: *Origin of Human Cancer* (Ed. Hiatt H.H., Watson J.D. and Winsten J.A.), Book A, pp. 1-12.
- Dunn B.P. (1982) Polycyclic aromatic hydrocarbons (PAH) In: *Carcinogens and Mutagens in the Environment. Vol. I: Food Products* (Ed. Stich H.F.), pp. 175-183, CRC Press Inc., Florida.

Eastman A., Sweetenham J. and Bresnick E. (1978) Comparison of in vivo and in vitro binding of polycyclic hydrocarbons to DNA. *Chem.-Biol. Int.*, 23, 345-353.

Ebright R.H., Wong J.R. and Chen L.B. (1986) Binding of 2-hydroxybenzo(a)pyrene to estrogen receptors in rat cytosol. *Canc. Res.*, 46, 2349-2351.

Edgar J.A. (1974) Ascorbic acid and biological alkylating agents. *Nature (New Biol.)*, 248, 136-137.

Emeh J.K., Niranjana B.G., Bhat N.K. and Avadhani N.G. (1981). Modulation of hepatic transcription and translation during early stages of aflatoxin B₁ carcinogenesis. *Carcinogenesis*, 2, 373-378.

Farber J.L., Shinozuka H., Serrani A. and Farmer R. (1974) Reversal of ethionine-induced inhibition of rat liver RNA polymerase in vivo by adenine. *Lab. Invest.*, 31, 462-472.

Feldman G., Remson J., Shinohara K. and Cerutti P. (1978) Excisability and persistence of benzo(a)pyrene DNA adducts in epithelial human lung cells. *Nature*, 274, 796-798.

Firozi P.F., Aboobaker V.S. and Bhattacharya R.K. (1986) Action of vitamin A on DNA adduct formation by aflatoxin B₁ in a amicrosome catalyzed reaction. *Canc. Lett.*, in press.

Flamm W.G., Counts W.B. and Banerjee M.R. (1966) Inhibition of ribonucleic acid synthesis in mouse skin by actinomycin D and 7,12-dimethylbenzanthracene. *Nature*, 210, 541-543.

Frolik C.A. and Roller P.P. (1981) The role of retinoids (vitamin A and its analogs) in the prevention of epithelial cancer. In: *Antitumor Compounds of Natural Origin: Chemistry and Biochemistry*, Vol II (Ed. Aszalos A.), pp. 87-128. CRC Press, Boca Raton, Florida.

Fukushima S., Imaida K., Sakata T., Okamura T., Shibata M.-A. and Nobuyuki I. (1983) Promoting effects of sodium L-ascorbate on the two-stage urinary bladder carcinogenesis in rats. *Canc. Res.*, 43, 4454-4457.

Fukushima S., Kurata Y., Shibata M.-A., Ikawa E. and Ito N. (1984) Promotion by ascorbic acid, sodium erythorbate and ethoxyquin of neoplastic lesions in rats initiated with N-butyl-N-(4-hydroxybutyl)nitrosamine. *Canc. Lett.*, 23, 29-37.

Gamper H.B., Tung A.S.C., Straub K., Bartholomew J.C. and Calvin M. (1977) DNA strand scission by benzo(a)pyrene diol epoxides. *Science*, 197, 671-674.

Gamper H.B., Straub K., Calvin M. and Bartholomew J.C. (1980) DNA alkylation and unwinding induced by benzo(a)pyrene diol epoxide: Modulation by ionic strength and superhelicity. *Proc. Natl. Acad. Sci.*, 77, 2000-2004.

Garyfallides S., Kyrtopoulos S.A. and Sekeris C.E. (1984) Effects of dimethylnitrosamine on RNA synthesis and metabolism in mouse liver, *Canc. Res.*, 44, 5110-5117.

Geacintov N.E., Yoshida H., Ibanez V. and Harvey R.G. (1982) Noncovalent binding of 7 β -, 8 α -dihydroxy-9 α , 10 α -epoxytetrahydrobenzo(a)pyrene to deoxyribonucleic acids and its catalytic effect on the hydrolysis of the diol epoxide to tetrol. *Biochemistry*, 21, 1864-1896.

Geacintov N.E. (1985) Mechanisms of interaction of polycyclic aromatic diol epoxides with DNA and structures of the adducts. In: ACS Symp. Series No. 283, Polycyclic Hydrocarbons and Carcinogenesis (Ed. Harvey R.G.), pp. 107-124, Amer. Chem. Soc.

Gelboin H.V., Worthan J.S., Wilson R.G., Freidman M. and Wogan G.N. (1966) Rapid and marked inhibition of rat liver RNA polymerase by aflatoxin B₁. *Science*, 154, 1205-1206.

Gelboin H.V. (1967) Carcinogens, enzyme induction and gene action. *Adv. Canc. Res.* 10, 1-81.

Gelboin H.V. (1969) A microsome-dependent binding of benzo(a)pyrene to DNA. In: The Jerusalem Symposium on Quantum Chemistry and Biochemistry, Vol. I: Physico-Chemical Mechanisms of Chemical Carcinogenesis (Ed. Bergmann E.D. and Pullman B.), pp. 175-182, The Israel Acad. Sciences and Humanities.

Gelboin H.V., Okuda T., Selkirk J.K., Nemoto N., Yang S.K., Rapp H.J. and Bast (Jr) R.C. (1976) Benzo(a)pyrene metabolism: Enzymatic and liquid chromatographic analysis and application to human tissues. In: Fundamentals in Cancer Prevention (Ed. Magee P.N. et al.) pp. 167-190, University of Tokyo Press, Tokyo / Univ. Park Press, Baltimore.

Gibson D.T., Mahadevan V., Jerina D.M., Yagi H. and Yeh H.J.C. (1975) Oxidation of the carcinogens benzo(a)pyrene and benzantracene to dihydrodiols by a bacterium. *Science*, 189, 295-297.

- Greenwald P. and Schreiber G. (1983) Perspectives on chemoprevention in man. In: Modulation and Mediation of Cancer by Vitamins (Ed. Meyskens F.L. and Prasad K.N.) pp. 40-46, S. Karger AG. Basel.
- Griffin G.D., Calle E.E., Morris M.D., Long K.Y., Schuffman S.S. and Mitchell W.M. (1986) Inhibition of murine interferon production following in vivo administration of benzo(a)pyrene. *J. Interferon Res.*, 6, 115-121.
- Grilli S., Rocchi P. and Prodi G. (1975) Non-enzymatic and microsome-dependent binding of polycyclic hydrocarbons to DNA or polynucleotides. *Chem.-Biol. Int.*, 11, 351-363.
- Grover P.L. and Sims P. (1968) Enzyme-catalysed reactions of polycyclic aromatic hydrocarbons with deoxyribonucleic acid and protein in vitro. *Biochem. J.*, 110, 159-160.
- Grover P.L., Hewer A. and Sims P. (1974) Metabolism of polycyclic hydrocarbons by rat lung preparations. *Biochem. Pharmacol.*, 23, 323-332.
- Grover P.L. (1979) Chemical Carcinogens and DNA. Vols I and II. CRC Press, Boca Raton, Florida.
- Grunberger G., Yu, F.-L., Grunberger D. and Feigelson P. (1973) Mechanism of N-hydroxy-2-acetylaminofluorene inhibition of rat hepatic ribonucleic acid synthesis. *J. Biol. Chem.*, 248, 6278-6281.
- Grunberger D. and Weinstein I.B. (1979) Biochemical effects of the modification of the nucleic acids by certain polycyclic aromatic carcinogens. *Progr. Nucl. Acid Res. Molec. Biol.*, 23, 106-149.
- Grunberger D. and Weinstein I.B. (1979b) Conformational changes in nucleic acids modified by chemical carcinogens. In: Chemical Carcinogens and DNA, vol. II (Ed. Grover P.L.), pp. 59-93, CRC Press, Boca Raton, Florida.
- Guengerich F.P. and Liebler D.C. (1985) Enzymatic activation of chemicals to toxic metabolites. *CRC Crit. Rev. Toxicol.*, 14, 259-307.
- Guttenplan J.B. (1977) Inhibition by L-ascorbate of bacterial mutagenesis induced by two N-nitroso compounds. *Nature*, 268, 368-370.
- Hahon N. and Booth J.A. (1986) Coinhibition of viral interferon induction by benzo(a)pyrene and chrysotile asbestos. *Environl. Res.*, 40, 103-109.

- Hanson-Painton O., Griffin M.J. and Tang J. (1981) Evidence for cytosolic benzo(a)pyrene carrier proteins which function in cytochrome P₄₅₀ oxidation in rat liver. *Biochem. Biophys. Res. Commun.*, 101, 1364-1371.
- Harris C.C., Autrup H., Connor R., Barrett L.A., McDowell E.M. and Trump B.F. (1976) Interindividual variation in binding of benzo(a)pyrene to DNA in cultured human bronchi. *Science*, 194, 1067-1069.
- Harris C.C., Autrup H., Stoner G.D., Trump B.F., Hillman E., Schafer P.W. and Jeffrey A.M. (1979) Metabolism of benzo(a)pyrene, N-nitrosodimethylamine and N-nitrosopyrrolidine and identification of the major carcinogen-DNA adducts formed in cultured human esophagus. *Canc. Res.*, 39, 4401-4406.
- Harvey R.G., Goh S.H. and Cortez C. (1975) "K-region" oxides and related oxidised metabolites of carcinogenic aromatic hydrocarbons. *J. Amer. Chem. Soc.*, 97, 3468-3479.
- Harvey R.G. and Dunne F.B. (1978) Multiple regions of metabolic activation of carcinogenic hydrocarbons. *Nature*, 273, 566-568.
- Harvey R.G. (1981) Activated metabolites of carcinogenic hydrocarbons. *Acc. Chem. Res.*, 14, 218-226.
- Hatch F.T., MacGregor J.T. and Zeiger E. (1986) Review: Putative mutagens and carcinogens in foods; VII. Genetic toxicology of the diet. *Environl. Mutagenesis*, 8, 467-484.
- Havsteen B. (1983) Flavonoids - a class of natural products of high pharmacological potency. *Biochem. Pharmacol.*, 32, 1141-1148.
- Heidelberger C. (1975) Chemical carcinogenesis. *Ann. Rev. Biochem.*, 44, 79-121.
- Herzog J., Serroni A., Briesmeister B.A. and Farber J.L. (1975) N-hydroxy-2-acetylaminofluorene inhibition of rat liver RNA polymerases. *Canc. Res.*, 35, 2138-2144.
- Herzog J. and Farber J.L. (1976) Inhibition of rat liver RNA polymerases by action of the methylating agents dimethylnitrosamine in vivo and methyl methanesulfonate in vitro. *ibid.*, 36, 1761-1770.
- Hill D.L. and Shih T. (1974) Vitamin A compounds and analogues as inhibitors of mixed function oxidases that metabolise carcinogenic polycyclic hydrocarbons and other compounds. *ibid.*, 34, 564-570.

- Hirose M., Fukushima S., Sakata T., Inui H. and Ito N. (1983) Effect of quercetin on two stage carcinogenesis of the rat urinary bladder. *Cancer Lett.*, 21, 23-27.
- Hishizawa T., Otsuka H. and Terayama H. (1964) Metabolic behaviour of nucleic acids in subcellular fractions from liver of rats treated with 3-methylcholanthrene. *J. Biochem.*, 56, 97-100.
- Hoffman H.D., Lesko (Jr) S.A. and T'so P.O.P. (1979) Chemical linkage of polycyclic aromatic hydrocarbons to DNAs and polynucleotides in aqueous solution and in a buffer-ethanol solvent system. *Biochemistry*, 9, 2594-2604.
- Holder G.M., Tierney B. and Bresnick E. (1981) Nuclear uptake and subsequent nuclear metabolism of benzo(a)pyrene complexed to cytosolic proteins. *Canc. Res.*, 41, 4408-4414.
- Holliday R. and Jeggo P.A. (1985) Mechanisms for changing gene expression and their possible relationship to carcinogenesis. *Canc. Surveys*, 4, 557-581.
- Hose J.E., Hannak J.B., Landolt M.L., Miller B.S., Felton S.P. and Iwaoka W.T. (1981) Uptake of benzo(a)pyrene by gonadal tissue of flatfish (family - pleuronectidae) and its effects on subsequent egg development. *J. Toxicol Environ. Hlth.* 7, 991-1000.
- Hradec J. and Kolar G.F. (1984) Microsome-mediated stimulation of the aminoacylation of rat liver initiator transfer ribonucleic acid by benzo(a)pyrene and 3,3-dimethyl-1-phenyltriazenes. *Canc. Lett.*, 23, 115-120.
- Huang M.-T., Wood A.W., Newmark H.L., Sayer J.M., Yagi H., Jerina D.M. and Conney A.H. (1983) Inhibition of the mutagenicity of bay region diol epoxides of polycyclic aromatic hydrocarbons by phenolic plant flavonoids. *Carcinogenesis*, 4, 1631-1637.
- Huang M.-T., Wood A.W., Chang R.L., Yagi H., Sayer J.M., Jerina D.M. and Conney A.H. (1984) Inhibition of the mutagenicity of bay-region diol epoxides of PAHs by the isomeric phenols of B(a)P. *Proc. Amer. Assoc. Canc. Res.*, 25, 97.
- Huberman E. and Sachs L. (1977) DNA binding and its relationship to carcinogenesis by different polycyclic hydrocarbons. *Int. J. Canc.* 19, 122-127.

Huckle K.R., Smith R.J., Watson W.P. and Wright A.S. (1986) Comparison of hydrocarbon-DNA adducts formed in mouse skin in vivo and in organ culture in vitro following treatment with benzo(a)pyrene. *Carcinogenesis*, 7, 965-970.

Hurlbert R.B. and Potter V.R. (1954) Nucleotide metabolism: I: The conversion of orotic acid-6-¹⁴C to uridine nucleotides. *J. Biol. Chem.* 209, 1-21.

IARC (1973) Monographs on the evaluation of carcinogenic risk of the chemical to man. Vol. 3: Certain Polycyclic Aromatic Hydrocarbons and Heterocyclic Compounds. B(a)P: pp. 91-136. B(e)P: 137-158, I.A.R.C., Lyon, France.

IARC (1983) Monographs on the evaluation of carcinogenic risk of chemical to humans. vol. 32: Polynuclear Aromatic Compounds part 1: Chemical, Environmental and Experimental Data. B(a)P: pp. 211-224, B(e)P: pp. 225-237, I.A.R.C., Lyon, France.

Imamura T., Busby (Jr) W.F., Goldman M.E., Wogan G.N. and Brunengraber H. (1982) Effects of hydroxyurea and benzo(a)pyrene on DNA synthesis in the isolated perfused rat lung. *Bull. Environ. Cont. Toxicol.*, 28, 135-140.

Irving C.C. (1973) Interaction of chemical carcinogen with DNA. *Methods Canc. Res.*, VII, 189-244.

Ivanovic V., Geacintov N.E. and Weinstein I.B. (1976) Cellular binding of benzo(a)pyrene to DNA characterized by low temperature fluorescence. *Biochem. Biophys. Res. Comm.*, 70, 1172-1179.

Ivanovic V., Geacintov N.E., Yamasaki H. and Weinstein I.B. (1978) DNA and RNA adducts formed in hamster embryo cell cultures exposed to benzo(a)pyrene. *Biochemistry*, 17, 1597-1603.

Jackson K.G. and Jones J.K.N. (1965) The C- and O-benzylation of L-ascorbic acid. *Can. J. Chem.*, 43, 450-457.

Jacob S.T., Sajdel E.M., Muecke W. and Munro H.N. (1970) Soluble RNA polymerases of rat liver nuclei: Properties, template specificity and amanitin responses in vitro and in vivo. *Cold Spr. Harb. Symp. Quant. Biol.*, 35, 681-691.

Jacob S.T. (1973) Mammalian RNA polymerases. *Prog.Nucl. Acid Res. Molec. Biol.*, 13, 93-126.

Jacobs M.M. and Griffin A.C. (1981) Trace elements and metals as anticarcinogens In: Inhibition of Tumor Induction and Development (Ed. Zadek M.S. and Lipkin M.), pp. 169-188, Plenum. New York.

Jahn C.L. and Litman G.W. (1979) Accessibility of deoxyribonucleic acid in chromatin to the covalent binding of the chemical carcinogen benzo(a)pyrene. *Biochemistry*, 18, 1442-1449.

Jamaluddin M., Rajalaxmi S. and Sarma D.S.R. (1981) Evidence for a direct interaction of L-ascorbate with rat liver DNA. *Biochem. Biophys. Acta*, 653, 423-427.

Jeffrey A.M., Weinstein I.B., Jennette K.W., Grzeskowiak K., Nakanishi K., Harvey R.G., Autrup H. and Harris C. (1977) Structures of benzo(a)pyrene-nucleic acid adducts formed in human and bovine bronchial explants. *Nature*, 269, 348-350.

Jeffrey A.M., Grzeskowiak K., Weinstein I.B., Nakanishi K., Roller P.P. and Harvey R.G. (1979) Benzo(a)pyrene-7,8-dihydrodiol 9,10-oxide adenosine and deoxyadenosine adducts: Structure and stereochemistry. *Science*, 206, 1309-1311.

Jennette K.W., Jeffrey A.M., Blobstein S.H., Beland F.A., Harvey R.G. and Weinstein I.B. (1977) Nucleoside adducts from the in vitro reaction of BP-7,8-dihydrodiol 9,10-oxide or BP 4,5-oxide with nucleic acids. *Biochemistry*, 16, 932-938.

Jenson J.C., FitzGibbon J., Brennan L.A. and Litman G.W. (1982) Binding of benzo(a)pyrene to histones and altered affinity of modified histone 1 for deoxyribonucleic acid. *ibid*, 21, 601-607.

Jerina D.M. and Daly J.W. (1974) Arene oxide: New aspect of drug metabolism. *Science*, 185, 573-582.

Jernstrom B., Vadi H. and Orrenius S. (1976) Formation in isolated rat liver microsomes and nuclei of benzo(a)pyrene metabolites that bind to DNA. *Canc. Res.*, 36, 4107-4113.

Jernstrom B., Orrenius S., Undeman, O., Graslund A. and Ehrenberg A. (1978) Fluorescence study of DNA-binding metabolites of benzo(a)pyrene formed in hepatocytes isolated from 3-methylcholanthrene treated rats. *ibid*, 38, 2600-2607.

Jones C.A., Moore B.P., Cohen G.M., Fry J.R. and Bridges J.W. (1978) Studies on metabolism and excretion of benzo(a)pyrene in isolated adult rat hepatocytes. *Biochem. Pharmacol.*, 27, 693-702.

Kahl R., Deckers-Schmelzlc B. and Klaus E. (1978) Ethoxyquin feeding to rats increases liver microsome catalysed formation of benzo(a)pyrene diol epoxide-DNA adduct. *Biochem. Biophys. Res. Comm.*, 85, 938-945.

Kahl R. (1982) Antioxidants and carcinogen metabolism. *Trends Pharmacol. Sci.*, 3, 72-74.

Takefuda T. and Yamamoto H.-A. (1978) Modification of DNA by the benzo(a)pyrene metabolite, diol-epoxide r-7,t-8-dihydroxy-t-9,10-oxy-7,8,9,10-tetrahydrobenzo(a)pyrene. *Proc. Natl. Acad. Sci.*, 75, 415-419.

Kallistratos G., Fasske E., Donos A. and Vadalouka-Kalfakatou V. (1983) The prophylactic and therapeutic effect of vitamin C on experimental malignant tumors. In: *Protective Agents in Cancer*. pp. 221-236, Acad. Press, London.

Kamm J.J., Dashman T., Conney A.H. and Burns J.J. (1975) Effect of ascorbic acid on amine-nitrite toxicity. *Ann. N.Y. Acad. Sci.* 258, 169-174.

Kaneko M. and Cerutti P.A. (1982) Excision of benzo(a)pyrene diol epoxide I adducts from nucleosomal DNA of confluent normal human fibroblasts. *Chem.-Biol. Int.*, 38, 261-274.

Kapitulnik J., Levin W., Yagi H., Jerina D.M. and Conney A.H. (1976) Lack of carcinogenicity of 4-,5-,6-,7-,8-,9- and 10-hydroxybenzo(a)pyrene on mouse skin. *Canc. Res.*, 36, 3625-3628.

Kapitulnik J., Levin W., Conney A.H., Yagi H. and Jerina D.M. (1977) Benzo(a)pyrene 7,8-dihydrodiol is more carcinogenic than benzo(a)pyrene in newborn mice. *Nature*, 266, 378-380.

Kapitulnik J., Wislocki P.G., Levin W., Yagi H., Jerina D.M. and Conney A.H. (1978a) Tumorigenicity studies with diol-epoxides of benzo(a)pyrene which indicate that (+)-trans-7 β ,8 α -dihydroxy-9 α ,10 α -epoxy-7,8,9,10-tetrahydrobenzo(a)pyrene is an ultimate carcinogen in newborn mice. *Canc. Res.*, 38, 354-358.

Kapitulnik J., Wislocki P.G., Levin W., Yagi H., Thakker D.R., Akagi H., Koreeda M., Jerina D.M. and Conney A.H. (1978b) Marked differences in the carcinogenic activity of optically pure (+)- and (-)-trans-7,8-dihydroxy-7,8-dihydrobenzo(a)pyrene in newborn mice. *ibid.*, 38, 2661-2665.

Kasai H. and Nishimura S. (1984) Hydroxylation of deoxyguanosine at the C-8 position by ascorbic acid and other reducing agents. *Nucl. Acid Res.*, 12, 2137-2145.

- Kidson C. and Kirby K.S. (1965) Selective alteration of rapidly-labeled ribonucleic acid synthesis in rat liver during azo-dye carcinogenesis. *Canc.Res.*, 25, 472-476.
- Kihlstrom I. (1986) Placental transfer of benzo(a)pyrene and its hydrophilic metabolites in the guinea pig. *Acta Pharmacol. Toxicol.*, 58, 272-276.
- Kimura T., Kodama M. and Nagata C. (1977) Differences in benzo(a)pyrene metabolism between lung and liver homogenates. *Biochem. Pharmacol.*, 26, 671-674.
- King H.W.S., Thompson M.H. and Brookes P. (1975) The benzo(a)pyrene-deoxyribonucleoside products isolated from DNA after metabolism of benzo(a)pyrene by rat liver microsomes in the presence of DNA. *Cance. Res.*, 35, 1263-1269.
- King H.W.S., Thompson M.H. and Brookes P. (1976a) The role of 9-hydroxybenzo(a)pyrene in microsome mediated binding of benzo(a)pyrene to DNA. *Int. J. Canc.*, 18, 339-344.
- King H.W.S., Thompson M.H., Tarmy E.M., Brookes P. and Harvey R.G. (1976b) On the nature of the product which results when 7,8-dihydrodiolbenzo(a)pyrene is metabolized in vitro and bound to DNA. *Chem.-Biol. Int.*, 13, 349-352.
- King H.W.S., Osborne M.R. and Brookes P. (1979) The in vitro and in vivo reactions at the N-position of guanine of the ultimate carcinogen derived from benzo(a)pyrene. *ibid*, 24, 345-353.
- Kinoshita N., Shears B. and Gelboin H.V. (1973) K and non-K metabolism of benzo(a)pyrene by rat liver microsomes. *Canc. Res.*, 33, 1937-1944.
- Kitagawa, T., Hirakawa, T., Ishikawa T., Nemoto N. and Takayama S. (1980) Induction of hepatocellular carcinoma in rat liver by initial treatment with benzo(a)pyrene after partial hepatectomy and promotion by phenobarbital. *Toxicol. Lett.*, 6, 167-171.
- Kleeberg U., Szeberenyi J., Juhasz P., Tigyi A. and Klinger W. (1982) Influence of 3-methylcholanthrene on liver nucleolar and nucleoplasmic activities of protein kinases and RNA polymerases. *Biochem. Pharmacol.*, 31, 1063-1067.
- Kodama M., Ioki Y. and Nagata C. (1977) Binding of benzo(a)pyrene-semiquinone radicals with DNA and polynucleotides. *GANN*, 68, 253-254.

- Kodama M. and Nagata C. (1977) Binding of 6-oxybenzo(a)pyrene radical with DNA and polynucleotides. *ibid*, 68, 125-126.
- Konieczny M. and Harvey R.G. (1982) Oxidized metabolites of 7-methylbenzo(a)pyrene. *Carcinogenesis*, 3, 573-575.
- Kootstra A. and Slaga T.J. (1980) Binding of isomers of benzo(a)pyrene diol-epoxide to chromatin. *Biochem. Biophys. Res. Comm.*, 93, 954-959.
- Kootstra A. (1982a) Formation and removal of B(a)P diol epoxide-DNA adducts in human fibroblasts. *Carcinogenesis*, 3, 953-955.
- Kootstra A. (1982b) Effect of histone acetylation on the formation and removal of B(a)P chromatin adducts. *Nucl. Acid Res.*, 10, 2775-2789.
- Koreeda M., Moore P.D., Yagi H., Yeh H.J.C. and Jerina D.M. (1976) Alkylation of polyguanylic acid at the 2-amino group and phosphate by the potent mutagen (+)7 β ,8 α -dihydroxy-9 β ,10 β -epoxy-7,8,9,10-tetrahydrobenzo(a)pyrene. *J. Amer. Chem. Soc.*, 98, 6720-6722.
- Kubinski H., Gutzke G.E. and Kubinski Z.O. (1981) DNA-cell binding (DCB) assay for suspected carcinogens and mutagens. *Mutat. Res.*, 89, 95-136.
- Kulkarni M.S., Calloway K., Irigaray M.F. and Kaufman D.G. (1986) Species differences in the formation of benzo(a)pyrene-DNA adducts in rodent and human endometrium. *Canc. Res.*, 46, 2888-2891.
- Kummet T. and Mayskens F.L. (1983) Vitamin A: A potential inhibitor of human cancer. *Seminars in Oncology*, X, 281-289.
- Kurian P., Jeffrey A.M. and Milo G.E. (1985) Preferential binding of benzo(a)pyrene diol epoxide to the linker DNA of human foreskin fibroblasts in S phase in the presence of benzamide. *Proc. Natl. Acad. Sci.*, 82, 2769-2773.
- Kurl R.N., Lund J., Poellinger L. and Gustafsson J.-A. (1982) Differential effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on nuclear RNA polymerase activity in the rat liver and thymus. *Biochem. Pharmacol.*, 31, 2459-2462.
- Kurokawa M., Hibino Y. and Sugano N. (1982) Associations of benzo(a)pyrene with rat-liver chromatin and the chromatin protein. *Chem.-Biol. Int.*, 39, 17-30.

Kuroki T. and Heidelberger C. (1971) The binding of polycyclic aromatic hydrocarbons to DNA, RNA and proteins of transformable cells in culture. *Canc. Res.*, 31, 2168-2176.

Lafarge-Frayssinet C., Alexandrov K., Rimbaut C., Dansette P.M., Mousset S. and Frayssinet C. (1981) Presence of benzo(a)pyrene metabolizing activities in isolated rat liver nuclei. *Carcinogenesis*, 2, 1189-1193.

Lakowicz J.R. and Hylden J.L. (1978) Asbestos-mediated membrane uptake of benzo(a)pyrene observed by fluorescence spectroscopy. *Nature*, 275, 446-448.

Leffler S., Pulkrabek P., Grunberger D. and Weinstein I.B. (1977) Template activity of calf thymus DNA modified by a dihydrodiol epoxide derivative of benzo(a)pyrene. *Biochemistry*, 16, 3133-3136.

Lefkowitz S.M., Brenner H.C., Astorian D.G. and Clarke R.H. (1979) Optically detected magnetic resonance study of benzo(a)pyrene-7,8-dihydrodiol 9,10-epoxide covalently bound to DNA. *FEBS Lett.*, 105, 77-80.

Lehr R.E., Yagi H., Thakker D.R., Levin W., Wood A.W., Conney A.H. and Jerina D.M. (1978) The bay region theory of polycyclic aromatic hydrocarbon-induced carcinogenicity. In: *Carcinogenesis: vol 3, Polynuclear Aromatic Hydrocarbon* (Ed. Jones P.W. and Freudenthal R.I.) pp 231-241, Raven Press, New York.

Lesko S.A. (Jr), Ts'o P.O.P. and Umans R.S. (1969) Interaction of nucleic acids V. Chemical linkage of 3,4-benzpyrene to deoxyribonucleic acid in aqueous solution. *Biochemistry*, 8, 2291-2298.

Levin W., Wood A.W., Yagi H., Dansette P.M., Jerina D.M. and Conney A.H. (1976) Carcinogenicity of benzo(a)pyrene 4,5-, 7,8-, and 9,10-oxides on mouse skin. *Proc. Natl. Acad. Sci.*, 73, 243-247.

Levin W., Wood A.W., Chang R.L., Slaga T.J., Yagi H., Jerina D.M. and Conney A.H. (1977) Marked differences in the tumor-initiating activity of optically pure (+)- and (-)-trans-7,8-dihydroxy-7,8-dihydrobenzo(a)pyrene on mouse skin. *Canc. Res.*, 37, 2721-2725.

Liberator P.A. and Bresnick E. (1981a) Effect of 3-methylcholanthrene administration on hepatic RNA polymerase activities. *Chem.-Biol. Int.*, 35, 93-110.

Idem (1981b) Further characterization of the effects of 3-methylcholanthrene administration upon hepatic ribonucleic acid polymerase activities. *ibid*, 38, 53-73.

Little J.B., McGandy R.B. and Kennedy A.R. (1978) Interactions between Polonium-210 α -radiation, benzo(a)pyrene, and 0.9% NaCl solution instillations in induction of experimental lung cancer. *Canc. Res.*, 38, 1929-1935.

Loeb L.A. and Gelboin H.V. (1964) Methylcholanthrene-induced changes in rat liver nuclear RNA. *Proc. Natl. Acad. Sci.*, 52, 1219-1226.

Loew G.H., Wong J., Phillips J., Hjelmeland L. and Pack G. (1978) Quantum chemical studies of the metabolism of benzo(a)pyrene. *Canc. Biochem. Biophys.*, 2, 123-130.

Lombardi P.E., Mayhew J.W., Goldin B.R., Gregory M.E., Lynch M.A., Sullivan C.E. and Gorbach L.S. (1981) Enzymatic methylation of microsomal metabolites of benzo(a)pyrene. *Canc. Res.*, 41, 4415-4419.

Lowry O.H., Rosebrough N.J., Farr A.L. and Randall R.L. (1951) Protein measurement with Folin phenol reagent. *J. Biol. Chem.*, 193, 265-275.

Lutz W.K., Viviani A. and Schlatter C. (1978) Nonlinear dose-response relationship for the binding of the carcinogen benzo(a)pyrene to rat liver DNA in vivo. *Cancer Res.*, 38, 575-578.

Lyte M. and Bick P.H. (1986) Modulation of interleukin-1 production by macrophages following benzo(a)pyrene exposure. *Int. J. Immunopharmacol.*, 8, 377-381.

Mabry T.J. and Ulubelen A. (1980) Chemistry and utilization of phenylpropanoids including flavonoids, coumarins and lignans. *Agricultural and Food Chem.*, 28, 188-196.

MacGregor J.T. and Jurd L. (1978) Mutagenicity of plant flavonoids: Structural requirements for mutagenic activity in *S.typhimurium*. *Mut. Res.*, 54, 297-309.

MacGregor J.T., Wehr C.H., Manners G.D., Jurd L., Minkler J.L. and Carrano A.V. (1983) In vivo exposure to plant flavanols: Influence of frequencies of micronuclei in mouse erythrocytes and sister-chromatid exchange in rabbit lymphocytes. *ibid*, 124, 255-270.

Mackenzie K.M. and Angevine D.M. (1981) Infertility in mice exposed in utero to benzo(a)pyrene. *Biol. Reprod.*, 24, 183-191.

- MacLeod M.C., Cohen G.M. and Selkirk J.K. (1979) Metabolism and macromolecular binding of the carcinogen benzo(a)pyrene and its relatively inert isomer benzo(e)pyrene by hamster embryo cells. *Canc. Res.*, 39, 3463-3470.
- MacLeod M.C., Kootstra A., Mansfield B.K., Slaga T.J. and Selkirk J.K. (1981) Binding of benzo(a)pyrene derivatives to specific proteins in nuclei of intact hamster embryo cells. *ibid*, 41, 4080-4086.
- MacLeod M.C., Mansfield B.K. and Selkirk J.K. (1982) Benzo(a)pyrene-binding proteins of hamster embryo cell nuclei: Comparison of nuclear isolation procedures. *J. Cellular Biochem.*, 18, 507-513./*Mech. of Chem. Carcinogenesis*, 299-305.
- MacLeod M.C., Pelling J.C., Slaga T.J., Noghrei-Nikbakht P.A., Mansfield B.K. and Selkirk J.K. (1983) Specificity of interaction between carcinogenic polynuclear aromatic hydrocarbons and nuclear proteins: Widespread occurrence of a restricted pattern of histone binding of intact cells. *Prog. Nucl. Acid Res. Molec. Biol.*, 29, 111-115.
- MacLeod M.C. and Zachary K. (1985) Catalysis of carcinogen-detoxification by DNA: Comparison of enantiomeric diol epoxides. *Chem.-Biol. Int.*, 54, 45-55.
- Madix J.C. and Bresnick E. (1967) Increased efficacy of liver chromatin as a template for RNA synthesis after administration of 3-methylcholanthrene. *Biochem. Biophys. Res. Comm.*, 28, 445-452.
- Marcus C.B., Turner C.R. and Jefcoate C.R. (1985) Binding of benzo(a)pyrene by purified cytochrome P-450. *Biochemistry*, 24, 5115-5123.
- Markov G.G. and Arion V.J. (1973) Characteristics of nuclear-ribosomal and DNA-like ribonucleic acids differentially extracted by hot-phenol fractionation. *Eur. J. Biochem.*, 35, 186-200.
- Marmur J. (1961) A procedure for the isolation of deoxyribonucleic acid from micro-organisms. *J. Molec. Biol.*, 3, 208-218.
- Marquardt H., Grover P.L. and Sims P. (1976) In vitro malignant transformation of mouse fibroblasts by non-K-region dihydrodiols derived from 7-methylbenz(a)anthracene, 7,12-dimethylbenz(a)anthracene and benzo(a)pyrene. *Canc. Res.*, 36, 2059-2064.

- Marshall C.J., Vousden K.H. and Phillips D.H. (1984) Activation of c-Ha-ras-1 proto-oncogene by in vitro modification with a chemical carcinogen benzo(a)pyrene diol-epoxide. *Nature*, 310, 586-589.
- Mattei E., Delpino A., Sacchi A. and Ferrini U. (1982) Characterization of the steady-state population of the poly (adenylic acid) sequences in a rat hepatoma and in normal liver. *Canc. Biochem. Biophys.*, 6, 13-18.
- McCann J., Choi E., Yamasaki E. and Ames B.N. (1975) Detection of carcinogens as mutagens in the Salmonella/microsome test: Assay of 300 chemicals. *Proc. Natl. Acad. Sci.*, 72, 5135-5139.
- McCormick J.J. and Maher V.M. (1985) Cytotoxic and mutagenic effects of specific carcinogen-DNA adducts in diploid human fibroblasts. *Environ. Hlth. Persp.*, 62, 145-155.
- McManus M.E., Minchin R.F., Schwartz D.M., Wirth P.J. and Huber B.E. (1986) Induction by phenobarbital in McA-RH 7777 rat hepatoma cells of a polycyclic hydrocarbon inducible cytochrome P-450. *Biochem. Biophys. Res. Comm.*, 137, 120-127.
- Meehan T., Straub K. and Calvin M. (1977) Benzo(a)pyrene diol epoxide covalently binds to deoxyguanosine and deoxyadenosine in DNA. *Nature*, 269, 725-727.
- Meehan T. and Straub K. (1979) Double-stranded DNA stereoselectively binds to benzo(a)pyrene diol epoxides. *ibid*, 277, 410-412.
- Meehan T., Gamper H. and Becker J.F. (1982) Characterization of reversible, physical binding of benzo(a)pyrene derivatives to DNA. *J. Biol. Chem.*, 257, 10479-10485.
- Mengle L., Gamper H. and Bartholomew J. (1978) Base specificity in the binding of benzo(a)pyrene diol epoxide to Simian virus 40 DNA. *Canc. Lett.*, 5, 131-137.
- Merrick B.A., Mansfield B.K., Nikbakht P.A. and Selkirk J.K. (1985) Benzo(a)pyrene metabolism in human T/47D mammary tumor cells: Evidence for sulphate conjugation and translocation of reactive metabolites across cell membrane. *ibid*, 29, 139-150.
- Mettlin G. (1983) The development of research on the epidemiology of vitamin A and cancer. In: *Diet, Nutrition and Cancer: From Basic Research to Policy Implications* (Ed. Roe D.A.) pp 125-134, Alan R. Liss Inc., New York.

- Miertus S. and Majek P. (1981) Calculation of reactivity indices for benz(a)anthracene and benzo(a)pyrene and their approximate models of complex with enzyme epoxidase. *Neoplasma*, 28, 441-446.
- Miller E.C. (1951) Studies on the formation of protein-bound derivatives of 3,4-benzpyrene in epidermal reaction of mouse skin. *Canc. Res.*, 11, 100-108.
- Miller J.A. (1970) Carcinogenesis by chemicals: An overview. G.H.A. Clowes memorial lecture. *ibid*, 30, 559-576.
- Miller J.A. and Miller E.C. (1977) Ultimate chemical carcinogens as reactive mutagenic electrophiles. In: *Origins of Human Cancer* (Eds. Hiatt H.H., Watson J.D. and Winsten J.H.) Book B, pp 605-627, Cold Spr. Harb. Lab.
- Mirvish S.S. (1975) Blocking the formation of N-nitroso compounds with ascorbic acid in vitro and in vivo. *Ann. N.Y. Acad. Sci.*, 258, 175-180.
- Idem (1981) Ascorbic acid inhibition of N-nitroso compound formation in chemical, food and biological systems. In: *Inhibition of Tumor Induction and Development* (Ed. Zadek M.S. and Lipkin M.), pp. 101-126, Plenum, New York.
- Misumi H., Hidaka H., Shohmori T. and Oda T. (1982) RNA synthesis in the presence of transfer RNA by DNA-dependent RNA polymerase II from mouse ascites sarcoma cells. *Arch. Biochem. Biophys.*, 215, 179-182.
- Mizusawa H. and Kakefuda T. (1979) Inhibition of DNA synthesis in vitro by binding of benzo(a)pyrene-metabolite diol-epoxide I to DNA. *Nature*, 279, 75-78.
- Mizusawa H., Lee C.-H., Kakefuda T., McKenney K., Shimatake H. and Rosenberg H. (1981) Base insertion and deletion mutations induced in an *Escherichia coli* plasmid by benzo(a)pyrene-7,8-dihydrodiol-9,10-epoxide. *Proc. Natl. Acad. Sci.*, 78, 6817-6820.
- Moon R.C., McCormick D.L. and Mehta R.G. (1983) Chemoprevention of animal tumors by retinoids. In: *Modulation and Mediation of Cancer by Vitamins. A : Vitamin A* (Ed. Meyskens F.L. and Prasad K.N.), pp 47-57, S. Karger, AG, Basel.
- Morse M.A. and Carlson G.P. (1985) Distribution and macromolecular binding of benzo(a)pyrene in Sencar and BALB/c mice following topical and oral administration. *J. Toxicol. Environ. Hlth.*, 16, 263-276.

- Muller R. and Rajewski M.F. (1981) Antibodies specific for DNA components structurally modified by chemical carcinogens. *J. Canc. Res. Clin. Oncol.*, 102, 99-113.
- Munro H.N. and Fleck A. (1966) The determination of nucleic acids. *Methods Biochem. Anal.*, 14, pp 113-176.
- Muramatsu M., Smetana K. and Busch H. (1963) Quantitative aspects of isolation of nucleoli of the Walker carcinosarcoma and liver of the rat. *Canc. Res.*, 23, 510-518.
- Nagao M., Morita N., Yahagi T., Shimizu M., Kuroyanagi M., Fukuyoka M., Yoshihira K., Natori S., Fujino T. and Sugimura T. (1981) Mutagenicity of 61 flavonoids and 11 related compounds. *Environ. Mutagenesis*, 3, 401-419.
- Nakanishi Y., Sekimizu K., Tamura H. and Natori S. (1981) Purification of a new protein stimulating RNA polymerase II from Ehrlich ascites tumor cells: Comparison with proteins purified before. *J. Biochem. (Tokyo)*, 90, 805-814.
- Naslund B. and von der Decken A. (1981) Effects of dietary protein conditions on RNA synthesis in liver nuclei from dimethylnitrosamine treated mice. *Nutr. Rep. Int.*, 23, 565-575.
- Natori S. (1982) Stimulatory proteins of RNA polymerase II from Ehrlich ascites tumor cells. *Molec. Cellular Biochem.*, 46, 173-187.
- Neal G.E. (1972) The effect of aflatoxin B₁ on normal and cortisol-stimulated ~~rat~~ liver ribonucleic acid synthesis. *Biochem. J.*, 130, 619-629.
- Idem (1973) Inhibition of rat liver RNA synthesis by aflatoxin B₁. *Nature*, 244, 432-435.
- Nemoto N. (1986) Marked activation of benzo(a)pyrene to protein-binding forms in the presence of unsaturated fatty acids and heme-compounds. *Carcinogenesis*, 7, 267-271.
- Newberne P.M. and Rogers A.E. (1981) Vitamin A, retinoids and cancer. In: *Nutrition and Cancer: Etiology and Treatment* (Ed. Newell G.R. and Ellison N.M.) pp. 217-232, Raven Press, New York.
- Newbold R.F. and Brookes P. (1976) Exceptional mutagenicity of a benzo(a)pyrene diol epoxide in cultured mammalian cells. *Nature*, 261, 52-54.

- Newmark H.L. and Mergens W.J. (1981) α -Tocopherol (vitamin E) and its relationship to tumor induction and development. In: Inhibition of Tumor Induction and Development (Ed. Zadek M.S. and Lipkin M.), pp 127-168, Plenum Press, New York.
- Niranjan B.G., Avadhani N.G. and DiGiovanni J. (1985) Formation of benzo(a)pyrene metabolites and DNA adducts catalyzed by a rat liver mitochondrial monooxygenase system. Biochem. Biophys. Res. Comm., 131, 935-942.
- Nishimoto M. and Varanasi U. (1985) Benzo(a)pyrene metabolism and DNA adduct formation mediated by English sole liver enzymes. Biochem. Pharmacol., 34, 263-268.
- Nixon J.E., Hendricks J.D., Pawlowski N.E., Pereira C.B., Simhuber R.O. and Bailey G.S. (1984) Inhibition of aflatoxin B₁ carcinogenesis in rainbow trout by flavone and indole compounds. Carcinogenesis, 5, 615-619.
- Nordenskjold M., Svenson S.-A., Jernstrom B., Moldeus P., Dock L. and Soderhall S. (1981) Studies on the in vitro transfer of DNA binding benzo(a)pyrene metabolites from rat hepatocytes to human fibroblasts. *ibid*, 2, 1151-1160.
- Novicki D.L., Rosenberg M.R. and Michalopoulos G. (1985) Inhibition of DNA synthesis by chemical carcinogens in cultures of initiated and normal proliferating rat hepatocytes. Canc. Res., 45, 337-344.
- Osborne M.R., Beland F.A., Harvey R.G. and Brookes P. (1976a) The reaction of (+)-7 α ,8 β -dihydroxy-9 β ,10 β -epoxy-7,8,9,10-tetrahydrobenzo(a)pyrene with DNA. Int. J. Canc., 18, 362-368.
- Osborne M.R., Thompson M.H., Tarmy E.M., Beland F.A., Harvey R.G. and Brookes P. (1976b) The reaction of 7,8-dihydro, 7,8-dihydroxybenzo(a)pyrene 9,10-oxide with DNA in relation to benzo(a)pyrene-DNA products isolated from cells. Chem.-Biol. Int., 13, 343-348.
- Osborne M.R., Harvey R.G. and Brookes P. (1978) The reaction of trans 7,8-dihydroxy-anti-9,10-epoxy-7,8,9,10-tetrahydrobenzo(a)pyrene with DNA involves attack at the N position of guanine moieties. *ibid*, 20, 123-130.
- Osborne M. and Merrifield K. (1985) Depurination of benzo(a)pyrene-diolepoxide treated DNA. *ibid*, 53, 183-195.
- Pamukcu A.M., Yalciner S., Hatcher J.F. and Bryan G.T. (1980) Quercetin, a rat intestinal and bladder carcinogen present in bracken fern (Pteridium aquilinum). Cancer Res., 40, 3468-3472.

- Parsons J.T. and McCarty K.S. (1968) Rapidly labeled messenger ribonucleic acid-protein complex of rat liver nuclei. *J. Biol. Chem.*, 243, 5377-5384.
- Paterour M.J., Bignon J. and Jaurand M.C. (1985) In vitro transformation of rat pleural mesothelial cells by chrysotile fibres and/or benzo(a)pyrene. *Carcinogenesis*, 6, 523-529.
- Pauling L. (1974) Are recommended daily allowances of vitamin C adequate ? *Proc. Natl. Acad. Sci.*, 71, 4442-4446.
- Peacock A.C. and Dingman C.W. (1967) Resolution of multiple ribonucleic acid, species by polyacrylamide gel electrophoresis. *Biochemistry*, 6, 1818-1827.
- Pelling J.C. and Slaga T.J. (1982) Comparison of levels of benzo(a)pyrene diol epoxide diastereomers covalently bound in vivo to macromolecular components of the whole epidermis versus basal cell layer. *Carcinogenesis*, 3, 1135-1141.
- Peterson F.J., Holloway D.-E., Duquette P.H. and Rivers J.M. (1983) Dietary ascorbic acid and hepatic mixed function oxidase activity in the guinea pig. *Biochem. Pharmacol.*, 32, 91-96.
- Petropoulos C.J., Lemire J.M., Goldman D. and Fausto N. (1985) Homology between rat liver RNA populations during development, regeneration and neoplasia. *Canc. Res.*, 45, 5114-5121.
- Pezzuto J.M., Lea M.A. and Yang C.S. (1976) Binding of metabolically activated benzo(a)pyrene to nuclear macromolecules. *ibid*, 36, 3647-3653.
- Phillips D.H. and Sims P. (1979) Polycyclic aromatic hydrocarbon metabolites: their reactions with nucleic acids. In: *Chemical carcinogens and DNA* (Ed. Grover P.L.), vol II, pp. 29-57, CRC Press, Florida.
- Phillips D.H., Hewer A. and Grover P.L. (1985) Aberrant activation of benzo(a)pyrene in cultured rat mammary cells in vitro and following direct application to rat mammary glands in vivo. *Canc. Res.*, 45, 4167-4174.
- Pietropaolo C. and Weinstein I.B. (1975) Binding of [^3H]-benzo(a)pyrene to natural and synthetic nucleic acids in subcellular microsomal system. *ibid*, 35, 2191-2198.

Pitts J.N., Lokensgard D.M., Ripley P.S., vanCauvenberghe K.A., Vaeck L.V. and Shaeffer T.B. (1980) "Atmospheric" epoxidation of benzo(a)pyrene by ozone: Formation of metabolite benzo(a)pyrene 4,5-oxide, *Science*, 210, 1347-1349.

Plowman J., Narayanan V.L., Dykes D., Szarvasi E., Briet P., Yoder O.C. and Paull K.D. (1986) Flavone acetic acid: A novel agent with preclinical antitumor activity against colon adenocarcinoma/38 in mice. *Canc. Trtmt. Rep.*, 70, 631-635.

Pong R.S. and Wogan G.N. (1970) Time-course and dose-response characteristics of aflatoxin B₁ effects on rat liver RNA polymerase and ultrastructure. *Canc. Res.*, 30, 294-304.

Post G.B., Snyder R. and Kalf G.F. (1984) Inhibition of mRNA synthesis in rabbit bone marrow nuclei in vitro by quinone metabolites of benzene. *Chem.-Biol. Int.*, 50, 203-211.

Preussmann R. (1976) Chemical carcinogens in the human environment: Problems and quantitative aspects. *Oncology*, 33, 51-57.

Prodi G., Rocchi P. and Grilli S. (1970) Binding of 7,12-dimethylbenzanthracene and benzo(a)pyrene to nucleic acids and proteins of organs in rats. *Canc. Res.*, 30, 1020-1023.

Pruess-Schwartz D. and Baird W.M. (1986) Benzo(a)pyrene:DNA adduct formation in early-passage Wistar rat embryo cell cultures: Evidence for multiple pathways of activation of benzo(a)pyrene. *ibid*, 46, 545-552.

Pruess-Schwartz D., Baird W.M., Nikbakht A., Merrick A. and Selkirk J.K. (1986) Benzo(a)pyrene:DNA adduct formation in normal human mammary epithelial cell cultures and the human mammary carcinoma T47D cell line. *ibid*, 46, 2697-2702.

Pullman A. and Pullman B. (1955) Electronic structure and carcinogenic activity of aromatic molecules. *Adv. Canc. Res.*, 3, 117-169.

Raj A.S. and Morris K. (1984) Inhibitory effects of α -naphthoflavone and β -naphthoflavone on 7,12-dimethylbenzanthracene induced anomalies in germ cells assessed by sperm abnormality assay. *Mut. Res.*, 136, 81-84.

Reeve V.E. and Gallagher C.H. (1981) Metabolism of (¹⁴C)-benzo(a)pyrene in vivo in the rat. *Canc. Lett.*, 13, 79-86.

Ridler P.J., Jennings B.R., Osborne M. and Brookes P. (1986) Electrofluorescence study of polycyclic hydrocarbon diol-epoxide binding to DNA. *Proc. R. Soc. Lond.*, B, 227, 441-454.

- Rinaldy A., Lockhart M.L., Deutsch J.F., Ungers G.E. and Rosenberg B.H. (1982) Effects of benzo(a)pyrene diol epoxide adducts on DNA synthesis in mammalian cells. *Mut. Res.*, 94, 383-391.
- Ringuette M.J., Gordon K., Szyszko J. and Krause M.O. (1982) Specific small nuclear RNAs from SV40-transformed cells stimulate transcription initiation in non-transformed isolated nuclei. *Can. J. Biochem.*, 60, 252-262.
- Roche C.J., Zinger D., Geacintov N.E. and Harvey R.G. (1985) Enhancement of stability of 7 β , 8 α -dihydroxy-9 α ,10 α -epoxybenzo (a)pyrene by complex formation with serum albumin. *Canc. Biochem. Biophys.*, 8, 35-40.
- Roe F.J.C. and Waters M.A. (1967) Induction of hepatoma in mice by carcinogen of the polycyclic hydrocarbon type. *Nature*, 214, 299-300.
- Roeder R.G. and Rutter W.J. (1969) Multiple forms of DNA-dependent RNA polymerase in eukaryotic organisms. *ibid*, 224, 234-237.
- Roeder R.G. and Rutter W.J. (1970) Specific nucleolar and nucleoplasmic RNA polymerases. *Proc. Natl. Acad. Sci.*, 65, 675-682.
- Roeder R.G. (1976) Eukaryotic nuclear RNA polymerases. In: *RNA polymerase* (Ed. Losick R. and Chamberlin M.) pp 285-329, Cold Spr. Harb. Lab.
- Rogan E.G. and Cavalieri E. (1974) 3-Methylcholanthrene-inducible binding of aromatic hydrocarbons to DNA in purified rat liver nuclei. *Biochem. Biophys. Res. Comm.*, 58, 1119-1126.
- Rogan E.G., Mailander P. and Cavalieri E. (1976) Metabolic activation of aromatic hydrocarbons in purified rat liver nuclei: Induction of enzyme activities and binding to DNA with and without monooxygenase-catalyzed formation of active oxygen. *Proc. Natl. Acad. Sci.*, 73, 457-461.
- Salazar I., Tarrago-Litvak L., Gil L. and Litvak S. (1982) The effect of benzo(a)pyrene on DNA synthesis and DNA polymerase activity of rat liver mitochondria. *FEBS Lett.*, 138, 45-49.
- Santella R.M., Hsieh L.L. and Perera F. (1986) Immunologic quantification of carcinogen-DNA adducts. In: *Mechanisms of DNA Damage and Repair* (Ed. Simic M.G., Grossman L. and Upton A.C.) pp. 509-517, Plenum Publ. Grp.

Sarma D.S.R., Rajalakshmi S. and Farber E. (1975) Chemical carcinogenesis: Interaction of carcinogens with nucleic acids. In: Cancer: A Comprehensive Treatise, Vol. 1: Etiology: Chemical and Physical Carcinogenesis (Ed. Becker F.F.), pp 235-287, Plenum Press, New York.

Sarraf A.M., Danenberg P.V., Heidelberger C. and Ketterer B. (1976) Separate identities of ligandin and the h-protein, a major protein to which hydrocarbons are covalently bound. *Biochem. Biophys. Res. Comm.*, 70, 869-877.

Saunders F.C., Barker E.A. and Smuckler E.A. (1972) Selective inhibition of nucleoplasmic rat liver DNA-dependent RNA polymerase by aflatoxin B₁. *Canc. Res.*, 32, 2487-2494.

Sayer J.M., Yagi H., Wood A.W., Conney A.H. and Jerina D.M. (1982) Extremely facile reaction between ultimate carcinogen benzo(a)pyrene 7,8-diol 9,10-epoxide and ellagic acid. *J. Amer. Chem. Soc.*, 104, 5562-5564.

Schelin C., Wallin H., Halpert J. and Jergil B. (1984) Covalent binding of benzo(a)pyrene to cytochrome P-450 / β NF-B₂ and other proteins in reconstituted mixed-function oxidase systems. *Chem.-Biol. Int.*, 49, 269-281.

Schlegel J.U., Pipkin G.E., Nishimura R. and Schultz G.N. (1970) The role of ascorbic acid in the prevention of bladder tumor formation. *J. Urol.*, 103, 155-159.

Schultz G., Gallwitz D. and Sekeris C.E. (1968) Rapidly labelled high molecular weight RNA from rat liver. *Eur. J. Biochem.*, 4, 149-156.

Sebti S.M., Pruess-Schwartz D. and Baird W.M. (1985) Species- and length of exposure-dependent differences in the benzo(a)pyrene:DNA adducts formed in embryo cell cultures from mice, rats and hamsters. *Canc. Res.*, 45, 1594-1600.

Selkirk J.K., Croy R.G. and Gelboin H.V. (1974) Benzo(a)pyrene metabolites: Efficient and rapid separation by High-pressure liquid chromatography. *Science*, 184, 169-171.

Selkirk J.K., Yang S.K. and Gelboin H.V. (1976) Analysis of benzo(a)pyrene metabolism in human liver and lymphocytes and kinetic analysis of benzo(a)pyrene in rat liver microsomes. In: Carcinogenesis, vol 1, Polynuclear Aromatic Hydrocarbons: Chemistry, Metabolism and Carcinogenesis (Ed. Freudenthal R.I. and Jones P.W.), pp 153-159, N.Y. Press.

Selkirk J.K. (1980) Comparison of epoxide and free-radical mechanisms for activation of benzo(a)pyrene by Sprague-Dawley rat liver microsomes. *J. Natl. Canc. Inst.*, 64, 771-774.

- Sentenac A. (1985) Eukaryotic RNA polymerases. CRC Crit. Rev. Biochem., 18, 31-91.
- Sestili M.A. (1983) Possible adverse health effects of vitamin C and ascorbic acid. Seminars in Oncol., X, 299-304.
- Shamberger R.J. (1972) Increase of peroxidation in carcinogenesis. J. Natl. Canc. Inst., 48, 1491-1497.
- Shamberger R.J., Baughman F.F., Kalchert S.L., Willis C.E. and Hoffman G.C. (1973) Carcinogen-induced chromosomal breakage decreased by antioxidants. Proc. Natl. Acad. Sci., 70, 1461-1463.
- Shen A.L., Fahl W.E., Wrighton S.A. and Jefcoate C.R. (1979) Inhibition of benzo(a)pyrene and benzo(a)pyrene 7,8-dihydrodiol metabolism by benzo(a)pyrene-quinones. Canc. Res., 39, 4123-4129.
- Shields D. and Tata J.R. (1976) Variable stabilities and recoveries of rat-liver RNA polymerase A and B according to growth status of the tissue. Eur. J. Biochem., 64, 471-480.
- Shinohara K. and Cerutti P.A. (1977) Excision repair of benzo(a)pyrene-deoxyguanosine adducts in baby hamster kidney 21/C13 cells and in secondary mouse embryo fibroblasts C57BL/6J. Proc. Natl. Acad. Sci., 74, 979-983.
- Shipman L.L. (1978) Ab initio quantum mechanical characterization of the ground electronic state of benzo(a)pyrene. Implications for the mechanism of polynuclear aromatic hydrocarbon oxidation to epoxides by cytochrome P-450. In: Carcinogenesis, vol 3: Polynuclear Aromatic Hydrocarbons (Ed. Jones P.W. and Freudenthal R.I.), pp 139-144, Raven Press, New York.
- Shirai T., Masuda A., Fukushima S., Hosoda K. and Ito N. (1985) Effects of sodium L-ascorbate and related compounds on rat stomach carcinogenesis initiated by N-methyl-N'-nitro-N-nitrosoguanidine. Canc. Lett., 29, 283-288.
- Shugart L., Holland J.M. and Rahn R.O. (1983) Dosimetry of PAH skin carcinogenesis; covalent binding of benzo(a)pyrene to mouse epidermal DNA. Carcinogenesis, 4, 195-198.
- Shugart L. and Kao J. (1984) Effect of ellagic acid and caffeic acid on covalent binding of benzo(a)pyrene to epidermal DNA of mouse skin in organ culture. Int. J. Biochem., 16, 571-573.
- Shugart L. and Matsunami R. (1985) Adduct formation in hemoglobin of the newborn mouse exposed in utero to benzo(a)pyrene. Toxicology, 37, 241-245.

Sikis B.I., Mimnaugh E.G. and Gram T.E. (1977) Effects of dietary ascorbic acid supplementation on hepatic drug-metabolizing enzymes in the guinea pigs. *Biochem. Pharmacol.*, 26, 2037-2041.

Silverman B.D. (1981) Carcinogenicity of methylated benzo(a)pyrene: Calculated ease of formation of the bay-region carbonium ion. *Canc. Biochem. Biophys.*, 5, 207-212.

Sims P. (1967) The metabolism of benzo(a)pyrene by rat liver homogenates. *Biochem. Pharmacol.*, 16, 613-618.

Sims P. (1970) Qualitative and quantitative studies on the metabolism of a series of aromatic hydrocarbons by rat liver preparations. *ibid.*, 19, 795-818.

Sims P., Grover P.L., Swaisland A., Pal K. and Hewer A. (1974) Metabolic activation of benzo(a)pyrene proceeds by a diol-epoxide. *Nature*, 252, 326-328.

Singer B. and Grunberger D. (1983) Molecular biology of mutagens and carcinogens. Ch.V: Metabolic activation of carcinogens and mutagens, pp. 97-141. Ch.VI: Reactions of metabolically activated carcinogens with nucleic acids, pp. 193-197. Ch. VII: Conformation of DNA modified by bulky aromatic carcinogens, pp 199-219. Plenum Press, New York.

Singh S.V., Srivastava S.K. and Awasthi Y.C. (1985) Binding of benzo(a)pyrene to rat lung glutathione S-transferases in vivo. *FEBS Lett.*, 179, 111-114.

Slaga T.J., Viage A., Berry D.L., Bracken W., Buty S.G. and Scribner J.D. (1976) Skin tumor-initiating activity of benzo(a)pyrene 4,5- and 7,8-diol- and 7,8-diol 9,10-epoxides. *Canc. Lett.*, 2, 115-122.

Slaga T.J. and Bracken W.M. (1977) The effects of antioxidants on skin tumor initiation and aryl hydrocarbon hydroxylase. *Canc. Res.*, 37, 1631-1635.

Slaga T.J., Bracken W.M., Dresner S., Levin W., Yagi H., Jerina D.M. and Conney A.H. (1978) Skin tumor initiating activities of the 12 isomeric phenols of benzo(a)pyrene. *ibid.*, 38, 678-681.

Slaga T.J. (1981) Food additives and contaminants as modifying factors in cancer induction. In: *Nutrition and Cancer: Etiology and Treatment* (Ed. Newell G.R. and Ellison N.M.), pp. 279-290, Raven Press, New York.

Slor H., Mizusawa H., Neihart N., Kakefuda T., Day R.S. III and Bustin M. (1981) Immunochemical visualization of binding of the chemical carcinogen benzo(a)pyrene diol-epoxide 1 to the genome. *Canc. Res.*, 41, 3111-3117.

Soderkvist P., Poellinger L. and Gustafsson J.-A. (1986) Carcinogen-binding proteins in the rat ventral prostate: Specific and nonspecific high-affinity binding sites for benzo(a)pyrene, 3-methylcholanthrene and 2,3,7,8-tetrachlorodibenzo-p-dioxin. *ibid*, 46, 651-657.

Soeiro R., Vaughan M.H., Warner J.R. and Darnell J.E. Jr. (1986) The turnover of nuclear DNA-like RNA in Hela cells. *J. Cell Biol.*, 39, 112-118.

Spelsberg T.C., Zytkevich T.H. and Moses H.L. (1977) Effects of metabolism on the binding of polycyclic hydrocarbons to nuclear subfractions of cultured AKR mouse embryo cells. *Canc. Res.*, 37, 1490-1496.

Sporn M.B. and Newton D.L. (1981) Retinoids and chemoprevention of cancer. In: *Inhibition of Tumor Induction and Development* (Ed. Zadek M.S. and Lipkin M.), pp 71-100, Plenum, New York.

Stairs P.W., Guzelian P.S. and vanTuyle G.C. (1983) Benzo(a)pyrene differentially alters mitochondrial and nuclear DNA synthesis in primary hepatocyte cultures. *Res. Comm. Chem. Pathol. Pharmacol.*, 42, 95-106.

Stampfer M.R., Bartholomew J.C., Smith H.S. and Bartley J.C. (1981) Metabolism of benzo(a)pyrene by human mammary epithelial cells: Toxicity and DNA adduct formation. *Proc. Natl. Acad. Sci.*, 78, 6251-6255.

Steel W.J. (1968) Localization of deoxyribonucleic acid complementary to ribosomal ribonucleic acid and preribosomal ribonucleic acid in the nucleolus of rat liver. *J. Biol. Chem.*, 243, 3333-3341.

Stein G.S., Stein J.L. and Thomson J.A. (1978) Chromosomal proteins in transformed and neoplastic cells: A review. *Canc. Res.*, 38, 1181-1201.

Stevens C.W., Bouck N., Burgess J.A. and Fahl W.E. (1985) Benzo(a)pyrene diol epoxides: different mutagenic efficiency in human and bacterial cells. *Mut. Res.*, 152, 5-14.

Stewart B.W. and Magee P.N. (1971) Effect of a single dose of dimethylnitrosamine on biosynthesis of nucleic acid and protein in rat liver and kidney. *Biochem. J.*, 125, 943-952.

- Stich H.F., Karim J., Koropatnick J. and Lo L. (1976) Mutagenic action of ascorbic acid. *Nature*, 260, 722-724.
- Stohs S.J., Grafstrom R.C., Burke M.D., Moldeus P.W. and Orrenius S.G. (1976) The isolation of rat intestinal microsomes with stable cytochrome P-450 and their metabolism of benzo(a)pyrene. *Arch. Biochem. Biophys.* 177, 105-116.
- Stowers S.J. and Anderson M.W. (1984) Ubiquitous binding of benzo(a)pyrene metabolites to DNA and protein in tissues of the mouse and rabbit. *Chem.-Biol. Int.*, 51, 151-166.
- Stowers S.J. and Anderson M.W. (1985) Formation and persistence of benzo(a)pyrene metabolite-DNA adducts. *Environ. Hlth. Persp.* 62, 31-39.
- Sullivan P.D. (1985) Free radicals of benzo(a)pyrene and derivatives. *ibid*, 64, 283-295.
- Supowit S.C. and Rosen J.M. (1980) Gene expression in normal and neoplastic mammary tissue. *Biochemistry*, 15, 3452-3460.
- Supowit S.C. and Rosen J.M. (1981) Tumor-specific polyadenylated RNA's from 7,12-dimethylbenz(a)anthracene-induced mammary tumors revealed through hybridization with fractionated single-copy DNA. *Canc. Res.*, 41, 3827-3834.
- Swann P.F., Peacock A.C. and Bunting S.C. (1975) Carcinogenesis and cellular injury: The effect of ethionine on RNA synthesis in rat liver. *Biochem. J.*, 150, 335-349.
- Swenberg J.A., Richardson F.C., Boucheron J.A. and Dyroff M.C. (1985) Relationships between DNA adduct formation and carcinogenesis. *Environ. Hlth. Persp.*, 62, 177-183.
- Szeberenyi J., Kleeberg U. and Tigyi A. (1982) Effect of 3-methylcholanthrene on RNA polymerase and protein kinase activities of transcriptionally active chromatin subfractions in rat liver. *Experientia*, 38, 687-688.
- Takeda N. and Teraniski K. (1986) Effects of environmental conditions on polycyclic aromatic hydrocarbons and mutagenicity of airborne particulate matter. *Bull. Environ. Contam. Toxicol.*, 36, 685-692.
- Tan K.B., Borun T.W., Charpentier R., Cristofalo V.J. and Croce C.M. (1982) Normal and neoplastic human cells have different histone H1 compositions. *J. Biol. Chem.*, 257, 5337-5338.

Thakker D.R., Yagi H., Levin W., Wood A.W., Conney A.H. and Jerina D.M. (1985) Polycyclic aromatic hydrocarbons: metabolic activation to ultimate carcinogens. In: Bioactivation of Foreign Compounds (Ed. Anders M.W.), pp. 177-242, Academic Press, New York.

Thomale J. and Nass G. (1982) Elevated urinary excretion of RNA catabolites as an early signal of tumor development in mice. *Canc. Lett.*, 15, 149-159.

Thompson A., Biggley W.H., Posner G.H., Lever J.R. and Seliger H.H. (1986) Microsomal chemiluminescence of benzo(a)pyrene-7,8-dihydrodiol and its synthetic analogues trans- and cis-1-methoxyvinylpyrene. *Biochem. Biophys. Acta*, 882, 210-219.

Thompson M.H., King H.W.S., Osborne M.R. and Brookes P. (1976) Rat liver microsome-mediated binding of benzo(a)pyrene metabolites to DNA. *Int. J. Canc.*, 17, 270-274.

Törnquist S., Wiklund L. and Toftgard R. (1985) Investigation of absorption, metabolism kinetics and DNA-binding of intra-tracheally administered benzo(a)pyrene in the isolated, perfused rat lung: A comparative study between microcrystalline and particulate adsorbed benzo(a)pyrene. *Chem.-Biol. Int.*, 54, 185-198.

Troll W., Belman S., Berkowitz E., Chmielewicz Z.F., Ambrus J.L. and Bardos T.J. (1968) Differential responses of DNA and RNA polymerase to modifications of the template rat liver DNA caused by action of the carcinogen acetylaminofluorene, in vivo and in vitro. *Biochem. Biophys. Acta*, 157, 16-24.

Turkington R.W. (1971) Changes in hybridizable nuclear RNA during the neoplastic development of mouse mammary cells. *Canc. Res.*, 31, 427-432.

Ueyama H., Matsuura T., Nomi S., Nakayasu H. and Ueda K. (1981) Binding of benzo(a)pyrene to rat liver nuclear matrix. *Life Sci.*, 29, 655-661.

Vadi H., Moldeus P., Capdevila J. and Orrenius S. (1975) Metabolism of benzo(a)pyrene in isolated rat liver cells. *Canc. Res.*, 35, 2083-2091.

Vahakangas K., Nebert D.W. and Pelkonen O. (1979) The DNA binding of benzo(a)pyrene metabolites catalyzed by rat lung microsomes in vitro and in isolated perfused rat lung. *Chem.-Biol. Int.*, 24, 167-176.

- Vahakangas K., Haugen A. and Harris C.C. (1985) An applied synchronous fluorescence spectrophotometric assay to study benzo(a)pyrene-diolepoxide-DNA adducts. *Carcinogenesis*, 6, 1109-1116.
- van der Hoeven J.C.M., Bruggeman I.M. and Debetes F.M.H. (1984) Genotoxicity of quercetin in cultured mammalian cells. *Mut. Res.*, 136, 9-21.
- vanDuuren B.L., Blazej T., Goldschmidt B.M., Katz C., Melchionne S. and Sivak A. (1971) Cocarcinogenesis studies on mouse skin and inhibition of tumor induction. *J. Natl. Canc. Inst.*, 46, 1039-1044.
- vanDuuren B.L. and Goldschmidt B.M. (1976) Cocarcinogenic and tumor-promoting agents in tobacco carcinogenesis. *ibid.*, 56, 1237-1242.
- Varanasi U., Nishimoto M. and Stover J. (1983) Analyses of biliary conjugates and hepatic DNA binding in benzo(a)pyrene exposed English sole. In: *Polynuclear Aromatic Hydrocarbons: Eighth International Symposium on Mechanisms, Methods and Metabolism* (Ed. Cooke M.W. and Dennis A.J.) pp 1315-1328, Battelle Press, Ohio.
- Vaught J. and Bresnick E. (1976) Binding of polycyclic hydrocarbons to nuclear components "in vitro". *Biochem. Biophys. Res. Comm.*, 69, 587-591.
- Vigny P. and Duquesne M. (1979) Fluorimetric detection of DNA-carcinogen complexes. In: *Chemical Carcinogens and DNA* (Ed. Grover P.L.), Vol I, pp 85-100, CRC Press Inc., Florida.
- Vinocour M. and Lewis R.A. (1985) Initiation of transcription in nuclei is inhibited by captan. *Chem.-Biol. Int.*, 56, 289-301.
- Viviani A. and Lutz W.K. (1978) Modulation of the binding of the carcinogen benzo(a)pyrene to rat liver DNA in vivo by selective induction of microsomal and nuclear aryl hydrocarbon hydroxylase activity. *Canc. Res.*, 38, 4640-4644.
- Wade C.G., Baker D.E. and Bartholomew J.C. (1978) Selective fluorescence quenching of benzo(a)pyrene and a mutagenic diol epoxide derivative in mouse cells. *Biochemistry*, 17, 4332-4337.
- Wang I.Y., Rasmussen R.E. and Crocker T.T. (1972) Isolation and characterization of an active DNA-binding metabolite of benzo(a)pyrene from hamster liver microsomal incubation systems. *Biochem. Biophys. Res. Comm.*, 49, 1142-1149.

- Waterfall J.F. and Sims P. (1972) Epoxy derivatives of aromatic polycyclic hydrocarbons. *Biochem. J.*, 128, 265-277.
- Wattenberg L.W., Page M.A. and Leong J.L. (1968) Induction of increased benzpyrene hydroxylase activity by flavones and related compounds. *Canc. Res.*, 28, 934-937.
- Wattenberg L.W. and Leong J.L. (1970) Inhibition of carcinogenic action of benzo(a)pyrene by flavones. *ibid*, 30, 1922-1925.
- Wattenberg L.W. (1978) Inhibition of chemical carcinogenesis. *J. Natl. Canc. Inst.*, 60, 11-18.
- Wattenberg L.W. (1981) Inhibitors of chemical carcinogens. In: *Cancer: Achievements, Challenges and Prospects for the 1980s* (Ed. Burchenal J.H. and Oettgen H.F.), vol 1, pp 517-539. Grune and Stratton, New York.
- Wattenberg L.W. (1985) Chemoprevention of cancer. *Canc. Res.*, 45, 1-8.
- Weibel F.J., Waters H.L., Elliot S. and Gelboin H.V. (1976) Effect of 7,8-benzoflavone on the duration of zoxazolamine paralysis and hexabarbital hypnosis in rats. *Biochem. Pharmacol.*, 25, 1431-1432.
- Weiland T. (1968) Poisonous principles of mushrooms of the genus *Amanita*. *Science*, 159, 946-952.
- Weinstein I.B., Jeffrey A.M., Jennette K.W., Blobstein S.H., Harvey R.G., Harris C., Autrup H., Kasai H. and Nakanishi K. (1976) Benzo(a)pyrene diol epoxides as intermediates in nucleic acid binding in vitro and in vivo. *Science*, 193, 592-595.
- West C.E. and Horton B.J. (1976) Transfer of polycyclic hydrocarbons from diet to milk in rats, rabbits and sheep. *Life Sci.*, 19, 1543-1552.
- Whalen D.L., Ross A.M., Montemarono J.A., Thakkar D.R., Yagi, H. and Jerina D.M. (1979) General acid catalysis in the hydrolysis of BP-7,8-diol 9,10-epoxides. *J. Amer. Chem. Soc.*, 101, 5086-5088.
- White K.L. and Holsapple M.P. (1984) Direct suppression of in vitro antibody production by mouse spleen cells by the carcinogen benzo(a)pyrene, but not by the noncarcinogenic congener benzo(e)pyrene. *Canc. Res.*, 44, 3388-3393.

- Widnell C.C. and Tata J.R. (1964) Evidence for two DNA-dependent RNA polymerase activities in isolated ~~rat~~ liver nuclei. *Biochem. Biophys. Acta*, 87, 531-533.
- Wilkinson D.S., Cihak A. and Pitot H.C. (1971) Inhibition of ribosomal ribonucleic acid maturation in rat liver by 5-fluoroorotic acid resulting in the selective labeling of cytoplasmic messenger ribonucleic acid. *J. Biol. Chem.*, 246, 6418-6427.
- Winicov I. (1981) Liver and kidney nuclear RNA synthesis and modifications in dimethylnitrosamine-treated rats. *Biochem. Biophys. Acta*, 654, 31-41.
- Wislocki P.G., Chang R.L., Wood A.W., Levin W., Yagi H., Hernandez O., Mah H.D., Dansette P.M., Jerina D.M. and Conney A.H. (1977) High carcinogenicity of 2-hydroxybenzo(a)pyrene on mouse skin. *Canc. Res.*, 37, 2608-2611.
- Wislocki P.G., Kapitulnik J., Levin W., Conney A.H., Yagi H. and Jerina D.M. (1978) Tumorigenicity of benzo(a)pyrene 4,5-, 7,8-, 9,10- and 11,12-oxides in newborn mice. *Canc. Lett.*, 5, 191-197.
- Wojciechowski M.F. and Meehan T. (1984) Inhibition of DNA methyltransferases in vitro by benzo(a)pyrene diol epoxide-modified substrates. *J. Biol. Chem.*, 259, 9711-9716.
- Wollenweber E. and Dietz V.H. (1981) Occurrence and distribution of free flavonoid aglycones in plants. *Phytochemistry*, 20, 869-932.
- Wood A.W., Levin W., Lu A.Y.H., Yagi H., Hernandez O., Jerina D.M. and Conney A.H. (1976) Metabolism of benzo(a)pyrene and benzo(a)pyrene derivatives to mutagenic products by highly purified microsomal enzymes. *J. Biol. Chem.*, 251, 4882-4890.
- Wood A.W., Sayer J.M., Newmark H.L., Yagi H., Michaud D.P., Jerina D.M. and Conney A.H. (1982a) Mechanism of the inhibition of mutagenicity of a benzo(a)pyrene 7,8-diol 9,10-epoxide by riboflavin 5'-phosphate. *Proc. Natl. Acad. Sci.*, 79, 5122-5126.
- Wood A.W., Huang M.-T., Chang R.L., Newmark H.L., Lehr R.E., Yagi H., Sayer J.M., Jerina D.M. and Conney A.H. (1982b) Inhibition of the mutagenicity of bay-region diol epoxides of polycyclic aromatic hydrocarbons by naturally occurring plant phenols: Exceptional activity of ellagic acid. *ibid*, 79, 5513-5517.

Yamano H., Sawai Y., Thung W.L., Sato F. and Tsukada K. (1982) Alterations in DNA-dependent RNA polymerase I from rat liver accompanying ethionine administration. *Biochem. Biophys. Res. Comm.*, 105, 799-805.

Yamaura I., Marquardt H. and Cavalieri L.F. (1978)= Effects of benzo(a)pyrene adducts on DNA synthesis in vitro. *Chem.-Biol. Int.*, 23, 399-407.

Yamaura I., Rosenberg B.H. and Cavlieri L.F. (1981) The major adducts of cis and trans benzo(a)pyrene diol epoxides cause chain termination during DNA synthesis in vitro. *ibid*, 37, 171-180.

Yan L.S. (1985) Study of carcinogenic mechanism for polycyclic aromatic hydrocarbons-extended bay region theory and its quantitative model. *Carcinogenesis*, 6, 1-6.

Yang N-c.C., Hrinyo T.P., Petrich J.W. and Yang D-D.H. (1983) Base sequence selectivity in binding of aromatic hydrocarbons with synthetic polynucleotides. *Biochem. Biophys. Res. Comm.*, 114, 8-13.

Yang S.K., Gelboin H.V., Trump B.F., Autrop H. and Harris C.C. (1977) Metabolic activation of benzo(a)pyrene and binding to DNA in cultured human bronchus. *Canc. Res.*, 37, 1210-1215.

Yoda K., Sakiyama S. and Fujimura S. (1978) Modification of rat liver chromatin by N-methyl-N'-Nitro-N-nitrosoguanidine or N-ethyl-N'-nitro-N-nitrosoguanidine and template activity for RNA synthesis by *Escherichia coli* RNA polymerase after reconstitution. *Biochem. Biophys. Acta*, 521, 677-688.

Yu F.-L. and Feigelson P. (1971) Cortisone stimulation of nucleolar RNA polymerase activity. *Proc. Natl. Acad. Sci.*, 68, 2177-2180.

Yu F.-L. (1974) Two functional state of the RNA polymerases in the rat hepatic nuclear and nucleolar frctions. *Nature*, 251, 344-346.

Yu F.-L. (1975) An improved method for the quantitative isolation of rat liver nuclear RNA polymerases. *Biochem. Biophys. Acta*, 395, 329-336.

Yu F.-L. and Grunberger D. (1976) Multiple sites of action of N-hydroxy-2-acetyl-amino-fluorene in rat hepatic nuclear transcription. *Canc. Res.*, 36, 3629-3633.

Yu F.-L. (1977) Mechanism of aflatoxin B₁ inhibition of rat hepatic nuclear RNA synthesis. *J. Biol. Chem.*, 252, 3245-3251.

Yu F.-L. (1980) Selective inhibition of rat liver nuclear RNA polymerase II by actinomycin D in vivo. *Carcinogenesis*, 1, 577-581.

Yu F.-L., Hunter G., Cass M., Barrett A., Yap T.G., Nowicki D. and Novak R. (1983) Effect of methylazoxymethanol acetate on rat liver nuclear and nucleolar RNA synthesis. *Canc. Res.* 43, 287-294.

Yuspa S.H. (1983) Retinoids and tumor promotion. In: *Diet, Nutrition and Cancer: From Basic Research to Policy Implications* (Ed. Roe D.A.), pp 95-109, Alan R. Liss. Inc., New York.

Zannoni V.G., Sato P.H. and Rikans L.E. (1978) Ascorbic acid and drug metabolism. In: *Nutrition and Drug Interactions* (Ed. Hathcock J.N. and Coon J.), pp 347-370, Academic Press, New York.

Zieve F.J. (1972) Inhibition of rat liver ribonucleic acid polymerase by the carcinogen N-hydroxy-2-fluorenylacetamide. *J. Biol. Chem.*, 247, 5987-5995.

Zytkowicz T.H., Moses H.L. and Spelsberg T. (1981) Covalent binding of benzo(a)pyrene metabolite to DNA, RNA and chromatin proteins in AKR mouse embryo cell line. *Chem.-Biol. Int.*, 35, 39-54.