List of Figures

	Fig. 1.1: Phylogenetic distribution of radiation-resistant <i>Eubacteria</i>
	Fig. 1.2 Types of ionising radiation and their effect of DNA and survival of organisms
	Fig. 1.3 Scanning electron micrograph of <i>Deinococcus radiodurans</i> R19
	Fig. 1.4 Carbonylation process in protein12
	Fig. 1.5 Kinetics and repair of DSB in D. radiodurans15
	Fig. 1.6 Regulation of ROS combating enzyme in <i>D. radiodurans</i> R121
	Fig. 1.7 XRF image analysis demonstrating the distribution of Mn^{2+} and Fe^{2+} in D. radiodurans R1
	Fig. 2.1: Schematic representation of the <i>Deinococcus</i> specific heminested PCR protocol and culture-independent DGGE analysis of deinococcal diversity
	Fig. 2.2: Optimization of the <i>Deinococcus</i> specific hemi-nested PCR42
	Fig. 2.3: Specificity of the <i>Deinococcus</i> specific hemi-nested PCR42
·	Fig. 2.4: Sensitivity of the hemi-nested PCR amplification for the amplification of deinococcal 16S rRNA gene fragments
	Fig. 2.5: Map showing the location of soil sampling sites
•	Fig. 2.6: <i>Deinococcus</i> specific nested PCR amplification from representative soil samples used in the study
	Fig. 2.7: Neighbor-joining tree based on 16S rRNA gene sequences showing the phylogenetic affiliation with type strains of <i>Deinococcus</i> species as listed in RDP49
	Fig. 2.8: PCR-DGGE analysis of deinococcal diversity from environmental samples.
	Fig. 2.9: Dendrogram for the DGGE profile from soil bacterial communities51
	Fig. 2.10: UV tolerance of the radiation-resistant bacterial isolates
	Fig. 2.11: Growth of radiation resistant bacterial isolates at different temperatures58
	Fig. 3.1: D50 determination of deinococcal strains to Hg ²⁺ 73
	Fig. 3.2 D50 determination of deinococcal strains to Cr ⁶⁺ 73
	Fig. 3.3 D50 determination of deinococcal strains to Cd ²⁺ 74

.

Fig. 3.4 Effect of growth phase on Cd ²⁺ tolerance in DR176
Fig. 3.5 Effect of exogenous Mn ²⁺ addition to Cd ²⁺ toxicity to stationary phase cultures in DR1
Fig. 3.6 Cd^{2+} induced stasis of the stationary phase culture of DR177
Fig. 3.7: Micrographs of the Cd ²⁺ treated DR1
Fig. 3.8 Fluoresence micrograph of Cd^{2+} exposed DR1 cells stained with H ₂ DCFDA.
Fig. 3.9 Effect of Cd^{2+} and H_2O_2 on ROS generation in DR181
Fig. 3.10 Effect of Cd ²⁺ on carbonylated proteins in DR1
Fig. 3.11 Detection of carbonylated proteins in DR1 exposed to Cd ²⁺ 82
Fig. 3.12 Cd ²⁺ induced lipid peroxidation in DR1
Fig. 3.13 Effect of Mn^{2+} and Cd^{2+} on recovery of DR1 from H_2O_2
Fig. 3.14 Effect of Cd ²⁺ on DR1 sod A ⁻ mutant
Fig. 3.15 A comparative analysis of Cd ²⁺ D50 for the wild type DR1 grown under different growth phases and the DR1 sod A- mutant
Fig. 3.16 Growth dependent expression of anti-oxidant enzymes in DR1
Fig. 3.17 Effect of Cd ²⁺ on the catalase in DR1
Fig. 3.17 Effect of Cd2+on the catalase in DR187Fig. 3.18 Effect of Cd2+on the SOD in DR188
Fig. 3.17 Effect of Cd^{2+} on the catalase in DR187Fig. 3.18 Effect of Cd^{2+} on the SOD in DR188Fig. 3.19 Effect of Cd^{2+} and Mn^{2+} on recovery of DR1 from UV radiation89
Fig. 3.17 Effect of Cd^{2+} on the catalase in DR187Fig. 3.18 Effect of Cd^{2+} on the SOD in DR188Fig. 3.19 Effect of Cd^{2+} and Mn^{2+} on recovery of DR1 from UV radiation89Fig. 3.20 Effect of Cd^{2+} on rec A ⁻ mutant of DR189
Fig. 3.17 Effect of Cd^{2+} on the catalase in DR187Fig. 3.18 Effect of Cd^{2+} on the SOD in DR188Fig. 3.19 Effect of Cd^{2+} and Mn^{2+} on recovery of DR1 from UV radiation89Fig. 3.20 Effect of Cd^{2+} on rec A ⁻ mutant of DR189Fig. 3.21 A comparative analysis of Cd^{2+} D50 for the wild type DR1 grown under different growth phases and the rec A- mutant.90
Fig. 3.17 Effect of Cd^{2+} on the catalase in DR187Fig. 3.18 Effect of Cd^{2+} on the SOD in DR188Fig. 3.19 Effect of Cd^{2+} and Mn^{2+} on recovery of DR1 from UV radiation89Fig. 3.20 Effect of Cd^{2+} on rec A ⁻ mutant of DR189Fig. 3.21 A comparative analysis of Cd^{2+} D50 for the wild type DR1 grown under different growth phases and the rec A- mutant.90Fig. 3.22 Amplification and clone confirmation of the pTZ57R/T-90
Fig. 3.17 Effect of Cd^{2+} on the catalase in DR187Fig. 3.18 Effect of Cd^{2+} on the SOD in DR188Fig. 3.19 Effect of Cd^{2+} and Mn^{2+} on recovery of DR1 from UV radiation89Fig. 3.20 Effect of Cd^{2+} on $rec A^{-}$ mutant of DR189Fig. 3.21 A comparative analysis of Cd^{2+} D50 for the wild type DR1 grown under different growth phases and the rec A- mutant.90Fig. 3.22 Amplification and clone confirmation of the pTZ57R/T-90Fig. 3.23 Clone confirmation of the Prec A - pRAD Z3clones using.91
Fig. 3.17 Effect of Cd^{2+} on the catalase in DR187Fig. 3.18 Effect of Cd^{2+} on the SOD in DR188Fig. 3.19 Effect of Cd^{2+} and Mn^{2+} on recovery of DR1 from UV radiation89Fig. 3.20 Effect of Cd^{2+} on rec A ⁻ mutant of DR189Fig. 3.21 A comparative analysis of Cd^{2+} D50 for the wild type DR1 grown under different growth phases and the rec A- mutant.90Fig. 3.22 Amplification and clone confirmation of the pTZ57R/T-90Fig. 3.23 Clone confirmation of the Prec A - pRAD Z3clones using.91Fig. 3.24 Vector map of Prec A -pRAD Z3.91
Fig. 3.17 Effect of Cd^{2+} on the catalase in DR187Fig. 3.18 Effect of Cd^{2+} on the SOD in DR188Fig. 3.19 Effect of Cd^{2+} and Mn^{2+} on recovery of DR1 from UV radiation89Fig. 3.20 Effect of Cd^{2+} on rec A ⁻ mutant of DR189Fig. 3.21 A comparative analysis of Cd^{2+} D50 for the wild type DR1 grown under different growth phases and the rec A- mutant.90Fig. 3.22 Amplification and clone confirmation of the pTZ57R/T-90Fig. 3.23 Clone confirmation of the Prec A - pRAD Z3clones using.91Fig. 3.24 Vector map of Prec A -pRAD Z3.91Fig. 3.25 Effect of Cd^{2+} on rec A promoter. Activity was assayed by using rec A- lac Z reporter in DR1.92
Fig. 3.17 Effect of Cd^{2+} on the catalase in DR187Fig. 3.18 Effect of Cd^{2+} on the SOD in DR188Fig. 3.19 Effect of Cd^{2+} and Mn^{2+} on recovery of DR1 from UV radiation89Fig. 3.20 Effect of Cd^{2+} on $rec A^-$ mutant of DR189Fig. 3.21 A comparative analysis of Cd^{2+} D50 for the wild type DR1 grown under different growth phases and the rec A- mutant.90Fig. 3.22 Amplification and clone confirmation of the pTZ57R/T-90Fig. 3.23 Clone confirmation of the Prec A - pRAD Z3clones using.91Fig. 3.24 Vector map of Prec A -pRAD Z3.91Fig. 3.25 Effect of Cd^{2+} on rec A promoter. Activity was assayed by using rec A- lac Z reporter in DR1.92Fig. 3.27 Cd^{2+} toxicity in PprI and PprM mutant of DR1.93

Fig. 3.28 SDS-PAGE analysis of total cellular proteins from DR1 treated with Cd ²⁻ under different growth conditions	+ 5
Fig. 3.29 2D gel electrophoresis of DR1 under different growth conditions	5
Fig. 3.30 Proteins differentially expressed under different growth conditions in DR1	l 7
Fig. 3.31 SDS PAGE of Mn ²⁺ and Cd ²⁺ binding proteins in DR1 obtained by IMAC	3
Fig. 3.32 Dendrogram obtained from In silico analysis of the proteins)
Fig. 3.33 IMAC purification)
Fig. 3.34 Effect of Cd ²⁺ on different ROS pathways in DR1100)
Fig. 4.1 Scheme for synthesis of <i>ec</i> 20111	t
Fig. 4.2 Synthesis and PCR amplification of ec20111	L.
M: ϕ X 174 DNA <i>Hae</i> III digest marker; 1: amplified <i>ec</i> 20111	ł
Fig. 4.3 T-vector clone confirmation of <i>ec-</i> 20112	2
Fig. 4.4 Vector map of pTZ57R-6ST112	2
Fig. 4.5 Cloning strategy of <i>ec</i> 20 in pRADZ3 and representative map of pRAD-EC clone	3
Fig. 4.6 Clone confirmation of ec20 obtained in pRADZ3113	
Fig. 4.7 Confirmation of ec 20 clones obtained in pRADZ3	
Fig. 4.8 PCR amplification of smt A from pMHNR1114	1
Fig. 4.9 Strategy employed for cloning smt A and representative map for the pRAD smtA clones	- 5
Fig. 4.10 Clone confirmation of pRAD-smtA using Spe I and BamHI115	5
Fig. 4.11 Clone confirmation of pRAD-smtA by PCR Amplification	5
Fig. 4.12 Expression analysis of the DR1 transformants carrying MT117	7
Fig. 4.13 Growth kinetics of DR1 (pRAD-EC) and DR1 (pRAD- smtA)118	3
Fig. 4.14 Cd ²⁺ tolerance of MT expressing transformants of DR1118	3
Fig. 4.15 Bioaccumulation of Cd ²⁺ by MT transformants of DR1119)
iz	ĸ

Fig. 4.16 Effect of cysteine on Cd ²⁺ tolerance on MT expressing transformants of DR1
Fig. 4.17: Cd ²⁺ tolerance of DR1 expressing MT in presence and absence of 0.4 mM cysteine
Fig. 4.18 Bioaccumulation of Cd ²⁺ by MT transformants of DR1 in presence of cysteine
Fig. 4.19 Sulfur recycling in DR1.Black arrows indicate the pathway identified in DR1

-

х