

LIST OF FIGURES

Table No.	Title	Page No.
	Review of Literature	
2.1	A conceptual framework of nutritional problems and causal factors in adolescence	6
2.2	Trends in Obesity among children and adolescents: United States, 1963-2008	15
2.3	Stages of nutrition transition	22
2.4	Recommended constituents of day's food intake	30
2.5	Iron needs during adolescence	37
2.6	Health promotion encompasses a range of activities	42
2.7	A framework for health promotion	43
	Methods & Materials	
3.1	Study design	84
	Results	
	PHASE I	
4.1.1	Sales trend of popular snacks (average pieces/day) at baseline in experimental school	136
4.1.2	Daily sales trend of beverages (glasses or tetra packs) at baseline in experimental school	137
4.1.3	Percent prevalence of nutritional status of the study subjects from the selected two schools	139
4.1.4	Prevalence of anemia, malnutrition, sub optimal blood pressure and central obesity among the study subjects from the selected two schools	140

4.1.5	Percent prevalence of Dyslipidemia (ATP III & AHA classifications) and Glucose Intolerance (IDF Classifications) in over weight and obese subjects from the selected two schools	141
4.1.6	Percent prevalence of modifiable risk factors among the study subjects from the selected two schools	141
	PHASE II	
4.2.1	Sale trend of proposed milk beverages and fruit juices in the five days trial period	168
	PHASE III	
4.3.1	Compliance of IFA supplementation among study subjects ($\geq 75\%$)	177
4.3.2	BMI chart plotting by the students after capacity building session	186
	Summary and Conclusion	
5.1	Percent prevalence of healthy and malnourished study subjects in the selected two schools	219
5.2	Percent prevalence of malnutrition, central obesity and high blood pressure among the study subjects	220
5.3	Mean number of products sold in five days pilot trial	220
5.4	Changes in prevalence of anaemia in the study subjects before and after intervention in relation to compliance of IFA tablets	221
5.5	Percent change in prevalence of anaemia in relation to compliance of IFA tablets (11-12 tablets) among the study subjects (N=197)	221
5.6	Percent change in prevalence of anaemia in relation to compliance of IFA tablets (9-10 tablets) among the study subjects (N=197)	222

5.7	Percent change in prevalence of anaemia in relation to compliance of IFA tablets (8 tablets) among the study subjects (N=197)	222
5.8	Percent change in prevalence of anaemia in relation to compliance of IFA tablets (< 8 tablets) among the study subjects (N=197)	223