

BIBLIOGRAPHY

- Abdal-Kader, M.M. and Barak, A.V. 1979. Evidence for a sex pheromone in the Hide beetle, Dermestes maculatus DeGeer (Coleoptera: Dermestidae). J. Chem. Ecol., 5 : 805 - 813.
- Averill, A.L. and Prokopy, R.J. 1981. Oviposition deterring fruit marking pheromone in Rhagoletis basiola. FLA Entomol., 64(2) : 222 - 226.
- Averill, A.L. and Prokopy, R.J. 1982. Oviposition deterring fruit marking pheromone in Rhagoletis zephyria. J. Ga. Entomol. Soc., 17 (3) : 315 - 319.
- Averill, A.L. and Prokopy, R.J. 1987 a. Fruit flies : Their Biology, Natural Enemies and Control; In : Host marking pheromones. eds. A.S. Robin and G.H.S. Hooper, Elsevier Press, New York.
- Averill, A.L. and Prokopy, R.J. 1987b. Residual activity of oviposition deterring pheromone in Rhagoletis pomonella (Diptera : Tephritidae) and female response to infested fruit. J. Chem. Ecol., 13 (1) : 167 - 177.
- Avidov, Z., Applebaum, S.W. and Berlinger, M.J. 1965. Physiological aspects of host specificity in the Bruchidae : ii. ovipositional preference and behaviour of Callosobruchus chinensis (L.). Entomol. Exp. Appl., 8 : 96 - 106.

- Barak, A.V. and Burkholder, W.E. 1977. Behaviour and pheromone studies with Attagenus elongatulus Casey (Coleoptera : Dermestidae). J. Chem. Ecol., 3 : 219 - 237.
- Behan, M. and Schoonhoven, L.M. 1978. Chemoreception of an oviposition deterrent associated with eggs in Pieris brassicae. Entomol. Exp. Appl., 24 : 163 - 179.
- Bentley, M.D., Mondaniel, I.N., Lee, H.P. Stiehl, B. and Yatagai, M. 1976. Studies of Aedes triseriatus, oviposition attractants produced by larvae of A. triseriatus and A. atropalpus (Diptera : Culicidae). J. Med. Ent., 13 (1) : 112 - 115.
- Bhalodia, N.K. 1974. Dissertation - "Biology of cigarette beetle, Lasioderma serricorne (F.) (Coleoptera : Anobiidae) and its feeding response to different types of Tobacco". B.A. College of Agriculture, Gujarat Agricultural University, Anand Campus, India.
- Bishop, O.N. 1983. Statistics for Biology, Micro-computer edition. Printed in Hong Kong by Wilture Printing Co. Ltd.
- Burkholder, W.E. 1970. Pheromone research with stored-product Coleoptera; pp. 1 - 20. In : Control of Insect Behaviour by Natural products, eds. D. Wood, R. Silverstein, M. Nakajima, New York : Academic Press.

- Carson, R. 1962. Silent Spring. Houghton Mifflin, Boston.
- Chadha, G.K. and Roome, R.E. 1980. Oviposition behaviour and the sensilla of the ovipositor of Chilo partellus and Spodoptera littoralis (Lepidoptera : Noctuidae). J. Zool., Lond., 192 : 169 - 178.
- Chapman, R.N. 1926. Inhibiting the process of metamorphosis in the confused flour beetle, Tribolium confusum Duval. J. Expt. Zool., 45 : 293 - 299.
- Chen, P.S., Stumm-zollinger, E., Aigaki, T., Balmer, J., Bienz, M. and Bohlen, P. 1988. A male accessory gland peptide that regulate reproductive behaviour of female Drosophila melanogaster. Cell., 54(3): 291 - 298.
- Chuman, T., Kohno, M., Kato, K. and Noguchi, M. 1979a. 4,6-Dimethyl-7-hydroxynonan-3-one, a sex pheromone of cigarette beetle (Lasioderma serricorne F.). Tetrahedron Lett., 25 : 2361 - 2364.
- Chuman, T., Kato, K. and Noguchi, M. 1979b. Synthesis of (+)-serricornin, 4,6-dimethyl-7-hydroxynonan-3-one, a sex pheromone of cigarette beetle (Lasioderma serricorne E.). Agric. Biol. Chem., 43 : 2005 - 2006.
- Chuman, T., Kohno, M., Kato, K., Noguchi, M., Nomi, H. and Mori, K. 1981. Stereoselective synthesis of erythro-serricornin (4R, 6R, 7S)-and (4S, 6R, 7S)-4, 6-dimethyl-7-hydroxynonan-3-one, stereoisomers of the sex pheromone of the cigarette beetle. Agric. Biol. Chem., 45 : 2019 - 2023.
- Chuman, T., Mochizuki, K., Mori, M., Kohno, M., Ono, M., Ohnishi, T. and Kato, K. 1982a. The pheromone activity of (+)-

- serricornin for male cigarette beetle, Lasioderma serricorne (F.). Agric. Biol. Chem., 46 : 593 - 595.
- Chuman, T., Mochizuki, K., Mori, M., Kohno, M., Kato, K., Nami, H. and Mori, K. 1982b. Behavioural and electroantennogram responses of male cigarette beetle (Lasioderma serricorne F.) to optically active serricornins. Agric. Biol. Chem., 46 : 3109 - 3112.
- Chuman, T., Mochizuki, K., Mori, M., Kohno, M., Kato, K. and Moguchi, M. 1985. Lasioderma chemistry, Sex pheromone of cigarette beetle (Lasioderma serricorne F.). J. Chem. Ecol., 11: 417-434.
- Coffelt, J.K. and Burkholder, W.E. 1972. Reproductive biology of cigarette beetle, Lasioderma serricorne. 1. Quantitative laboratory bioassay of the female sex pheromone from females of different ages. Ann. Entomol. Soc. Am., 65: 447 - 450.
- Colwell, A.E., Shorey, H.H., Bäumer, P. and Van Vorhisey, S.E. 1978. Sex pheromone scent marking by females of Pectinophora gossypiella (Lepidoptera : Gelechiidae). J. Chem. Ecol., 4 (6): 717 - 722.
- Corbet, S.A. 1971. Mandibular gland secretion of larvae of the flour moth, Anagasta kuehniella, contains an epideictic pheromone and elicits oviposition movements in a hymenopteran parasite. Nature, 332 : 481 - 484.
- Crajinar, R.M. and Prokopy, R.J. 1982. Morphological and electrophysiological mapping of tarsal chemoreceptors of oviposition deterring pheromone in Rhagoletis pomonella flies. J. Insect Physiology, 28 : 393 - 400.

- Cross, W.H., Hardee, D.D., Nicholas, F., Mitchell, H.C.,
Mitchell, E.B., Huddleston, P.M. and Tumlinson, J.H.
1969. Attraction of female boll weevils of traps
baited with males. J. Econ. Ent., 62 (2) : 159 - 160.
- Dick, J. 1937. Oviposition in certain coleoptera.
Ann. Appl. Biol., 24 : 762 - 796.
- Dittrick, L.E., Jones, R.L. and Chiang, H.C. 1983. An
oviposition deterrent for the European corn borer,
Ostrinia nubilalis (Lepidoptera : Pyralidae),
extracted from larval frass. J. Insect Physiol., 29 (1)
119 - 121.
- Faustini, D.L., Burkholder, W.E. and Laub, R.J. 1981.
Sexually dimorphic setiferous sex patch in the
male red flour beetle, Tribolium castaneum (Hbst.)
(Coleoptera : Tenebrionidae) : Site of aggregation
pheromone production. J. Chem. Ecol., 7 : 465 - 480.
- Faustini, D.L., Rowe, J.R. and Burkholder, W.E. 1982a.
A male produced aggregation pheromone in Tribolium
brevicornis Leconte (Coleoptera : Tenebrionidae)
and interspecific responses of several Tribolium
species. J. Stored Prod. Res., 18 : 153 - 158.
- Faustini, D.L., Giese, W.L., Philips, J.K. and Burkholder,
W.E. 1982b. Aggregation pheromone in Tribolium
brevicornis Leconte (L.). J. Chem. Ecol., 8 :
679 - 687.

- Fenemore, P.G. 1978. Oviposition of potato tuber moth, Phthorimaea operculella Zell. (Lepidoptera : Gelechiidae). The physical nature of the oviposition substrate. New Zealand J. Zoology, 5 : 591 - 599.
- Fletcher, L.W. and Long, J.S. 1971. Influence of food odours on oviposition by the cigarette beetle on non- food materials. J. Econ. Ent., 64 (4) 770 - 771.
- Ghent, W.E. 1963. Studies of behaviour of the Tribolium flour beetles. 1. Contrasting responses of T. castaneum and T. confusum to fresh and conditioned flours. Ecology, 44 : 269 - 283.
- Ghent, A.W. 1966. Studies of behaviour of the Tribolium flour beetles. II. Distributions in depth of T. castaneum and T. confusum in fractionable shell vials. Ecology, 47 : 355 - 367.
- Giesen, H., Khonle, U., Vite, J.P., Pan, M. L. and Francke, W. 1984. Investigation on the aggregation pheromone of a mediterranean engraver beetle, Ips erosus. Z. Angew. Entomol., 98 (1) : 95 - 97.
- Giga, D.P. and Smith, R.H. 1985. Oviposition markers in Callosobruchus maculatus and Callosobruchus rhodesianus (Coleoptera : Bruchidae). Asymmetry of interspecific responses. Agric. Ecosyst. Environ., 12 : 229 - 234.

- Gomez, K.A. and Gomez, A.A. 1976. Statistical procedure for agricultural research. 6th edition. Academic Press.
- Halstead, D.G.H. 1963. External sex differences in stored product Coleoptera. Bull. Ent. Res., 54 : 119 - 135.
- Hiroshi, N. 1978. Oviposition preference of Culex pipiens molestus and Cuex tritaenorrhynchus sumnorosus on to the water conditioned by the egg rafts or larvae. Jpn. J. Sanit. Zool., 29 (2) : 117 - 124.
- Howe, R.W. 1957. A study of the cigarette beetle, Lasiderma serricorne (F.) Coleoptera : Anobiidae with a critical review of the literature on its biology. Bull. Ent. Res., 48 : 9 - 56.
- Hughes, A.L. 1982. Attraction of adult T. confusum to flour conditioned by male conspecifics. Behav. Processes, 7 : 247 - 253.
- Hwang, Y.S., Schultz, G.W., Axelrod, H., Kramer, W.L. and Mulla, M.S. 1982. Oviposition repellency of fatty acids and their derivatives against Culex and Aedes mosquitoes. Environ. Entomol., 611 (1) : 223 - 226.
- Jacob, S.A. and Mohan, M.S. 1973. Predation on certain stored product insects by red flour beetle. Indian J. Ent., 35 : 95 - 98.
- Jacobson, M. 1975. Insecticides of the future. Marcel Dekker, Inc. New York.

- Kabir, S.M.H. 1970. Interactions of lesser grain borer, Rhyzopertha dominica (Fab.) and red flour beetle, Tribolium castaneum (Herbst) with developmental stages of Rice weevil, Sitophilus zeamais (Mots.). Scientific Researches, B.C.S.I.R. Bangladesh, 7 : 67 - 71.
- Katsoyannos, B.I. 1975. Oviposition deterring, male arresting, fruit marking pheromones in Rhagoletis cerasi. Environ. Entomol., 4 : 801 - 807.
- Khorramshahi, A. and Burkholder, W.E. 1981. Behaviour of the lesser grain borer, Rhyzopertha dominica (Coleoptera : Bostrichidae); male produced aggregation pheromone attracts both sexes. J. Chem. Ecol., 2 : 33 - 38.
- Klijnstra, J.W. 1982. Perception of the oviposition deterrent pheromone in Pieris brassicae. In : J.H.Visser and A.K. Minks (eds.), Proc. 5th Int. Symp. Insect - Plant Relationships. Pudoc, Wageningen : 145 - 151.
- Klijnstra, J.W. 1985. Interspecific egg load assessment of host plants by Pieris rapae butterflies. Entomol. Exp. Appl., 38 : 227 - 231.
- Klijnstra, J. W. 1986. The effect of an oviposition deterring pheromone on egg laying in Pieris brassicae. Entomol. Exp. Appl., 41 : 139 - 146.

- Klijnstra, J.W. and Roessingh, P. 1986. Perception of the oviposition deterring pheromone by tarsal and abdominal contact chemoreceptors in Pieris brassicae. Entomol. Exp. Appl., 40 : 71 - 79.
- Klijnstra, J.W. and Schoonhoven, L.M. 1987. Effectiveness and persistence of the oviposition deterring pheromone of Pieris brassicae in the field. Entomol. Exp. Appl., 45 (3) : 227 - 236.
- Kohno, M., Chuman, T., Kato, K. and Noguchi, M. 1983. The olfactory response of the cigarette beetle, Lasioderma serricorne, to various host foods and cured tobacco extracts. Appl. Entomol. Zool., 18 (3): 401 - 406.
- Kohno, M. and Ohnishi, A. 1986. Some factors affecting the oviposition behaviour of cigarette beetle (Coleoptera: Anobiidae). Kontyu, 54 : 29 - 32.
- Lefkovitch, L.P. and Currie, J.E. 1967. Factors affecting adult survival and fecundity in Lasioderma serricorne (F.) with special reference to its life history and control. Ecol. Monogr., 1 : 333 - 393.
- Levinson, H.Z., Levinson, A.R., Francke, W., Mackenroth, W. and Heeman, V. 1981. The pheromone activity of anhydroserriornin and serriornin for male cigarette beetle (Lasioderma serricorne F.). Naturwissenschaften, 68 : 148 - 149.

- Levinson, H.Z., Levinson, A.R., Kahn, G.E. and Schafer, K.
1983. Occurrence of a pheromone - producing gland
in female tobacco beetles. Experientia, 39 : 1095 -
1097.
- Levinson, H.Z., Levinson, A.R., Kahn, G.E. and Schafer, K.
1986. Antagonized pheromone responses of male
tobacco beetles (Lasioderma serricornne F.) due to
erythro-diastereoisomers of 4S, 6S, 7S - serricornin.
Naturwissenschaften, 73 : 36 - 37.
- Levinson, H.Z. and Levinson, A.R. 1987. Pheromone biology
of the tobacco beetle (Lasioderma serricornne F.,
Anobiidae) with notes on the pheromone antagonism
between 4S, 6S, 7S- and 4S, 6S, 7R - serricornin.
J. Appl. Ent., 103 : 217 - 240.
- Loconti, J.D. and Roth, L.M. 1953. Composition of the odorous
secretion of Tribolium castaneum. Ann. Ent. Soc. Am.,
46 : 281 - 289.
- Lundgren, L. 1975. Natural plant chemicals acting as ovi-
position deterrents on Cabbage butterflies, Pieris
brassicae (L.); P. rapae (L.) and P. napi (L.).
Zool. Scripta, 4 : 253 - 258.
- Ma, W.C. and Schoonhoven, L.M. 1973. Tarsal contact chemo-
sensory hairs of the large white butterfly Pieris
brassicae and their possible role in oviposition
behaviour. Ent. Exp. Appl., 16 : 343 - 357.

- Mallikarjuna Rao, K., Jacob, S.A. and Mohan, M.S. 1972.
Resistance of flexible packaging materials to some important pests of stored products. Indian J. Ent., 34 : 94 - 101.
- Mcneil, J.N. and Quiring, D.T. 1983. Evidence of an oviposition deterring pheromone in the alfalfa blotch leafminer, Agromyza frontella (Rondani) (Diptera : Agromyzidae). Environ. Entomol., 12 : 990 - 992.
- Mertz, D.B. 1969. Age distribution and abundance in populations of flour beetle (1) experimental studies. Ecol. Monogr., 39 : 1 - 31.
- Messina, F.J. and Renwick, J.A.A. 1985a. Ability of ovipositing seed beetles to discriminate between seeds with differing egg loads. Ecol. Entomol., 10 : 225 - 230.
- Messina, F.J. and Renwick, J.A.A. 1985b. Mechanism of egg recognition by the cowpea weevil, Callosobruchus maculatus. Entomol. Exp. Appl., 37 : 241 - 245.
- Messina, F.J., Barmore, J.L. and Renwick, J.A.A. 1987. Oviposition deterrent from eggs of Callosobruchus maculatus : Spacing mechanism or artifact. J. Chem. Ecol., 13 : 219 - 226.
- Millar, J.G., Pierce, H.G., Pierce, A.M., Ochlschlager, A.C., Borden, J. H. and Barak, A.V. 1985. Aggregation pheromone of the flat grain beetle, Cryptolestes

- pusillus (Coleoptera : Cucujidae). J. Chem. Ecol., 11 (8) : 1055 - 1070.
- Millar, J.G., Pierce, Jr. H.D., Pierce, A.M., Oehlschlager, A.C. and Borden, J.H. 1985. Aggregation pheromone of the grain beetle, Cryptolestes turcicus (Coleoptera: Cucujidae). J. Chem. Ecol., 11 (8) 1071 - 1081.
- Mitchell, E.R. 1975. The evolution of oviposition tactics in the bean weevil, Callosobruchus maculatus (F.). Ecology, 56 : 696 - 702.
- Mitchell, E.R. and Heath, R.R. 1985. Influence of allelochemicals on oviposition behaviour of Spodoptera spp. J. Chem. Ecol., 11 : 609 - 618.
- Mondal, K.A.M.S.H. 1983. Response of T. castaneum larvae Herbst to the different components of conditioned medium. Tribolium Inf. Bull., 23 : 104 - 111.
- Mondal, K.A.M.S.H. 1984. A method of determining the larval instars of Tribolium castaneum (Herbst) (Coleoptera: Anobiidae). Lab. Practice, 33 : 10 - 15.
- Mondal, K.A.M.S.H. and Port, G.R. 1984. Response of T. castaneum larvae to synthetic aggregation pheromone. Entomol. Exp. Appl., 36 (1) : 43 - 46.
- Mondal, K.A.M.S.H. 1985. Response of T. castaneum larvae to aggregation pheromone and quinones produced by adult conspecifics. J. International pest control, 27 : 64 - 66.

- Mori, K., Nomi, H., Chuman, T., Kohno, M., Kato, K. and Noguchi, M. 1981. Determination of the absolute configuration at C-6 and C-7 of serricornin (4,6-dimethyl-7-hydroxy-3-nonanone), the sex pheromone of the cigarette beetle. Tetrahedron Lett., 22 : 1127-1130.
- Mori, K., Nomi, H., Chuman, T., Kohno, M., Kato, K. and Noguchi, M. 1982. Synthesis and absolute stereochemistry of serricornin (4S, 6S, 7S) -4, 6- dimethyl-7-hydroxy - 3- nonanone, the sex pheromone of the cigarette beetle. Tetrahedron Lett., 38 : 3705 - 3711.
- Mori, M., Chuman, T., Kohno, M., Kato, K., Noguchi, M., Nomi, H., and Mori, K. 1982a. Absolute stereochemistry of serricornin, the sex pheromone of the cigarette beetle, as determined by the synthesis of its (4S, 6R, 7R) isomer. Tetrahedron Lett., 23 : 667 - 670.
- Mori, M., Chuman, T., Kato, K. and Mori, K. 1982b. A stereoselective synthesis of "natural" (4S, 6S, 7S)-serricornin, the sex pheromone of the cigarette beetle, from levoglucosenone. Tetrahedron Lett., 23 : 4593 - 4596.
- Mumtaz, M.M. and Aliniaze, M.T. 1983. The oviposition deterring pheromone in the western cherry fruit fly, Rhagoletis indifferens (Diptera : Tephritidae) : Biological properties. Z. Angew. Entomol., 96 (1): 83 - 93.
- Naylor, A.F. 1959. An experimental analysis of dispersal

- in the flour beetle, Tribolium confusum. Ecology,
40 : 453 - 465.
- O'Ceallachain, D.P. and Ryan, M.F. 1977. Production and
 perception of pheromones by the beetles T. confusum.
J. Insect Physiol., 23 : 1303 - 1309.
- Ogden, J.C. 1969. Effects of components of conditioned
 medium on behaviour in T. confusum. Physiol. Zool.,
42 : 266 - 274.
- Osgood, C.E. 1971. An oviposition attractant pheromone
 associated with the egg rafts of Culex tarsalis.
J. Econ. Ent., 64 (5) : 1038 - 1041.
- Oshima, K., Honda, H. and Yamamota, I. 1973. Isolation of
 an oviposition marker from azuki bean weevil,
Callosobruchus chinensis (L.). Agric. Biol. Chem.,
37 : 2679 - 2680.
- Park, T. 1937. Experimental studies on insect populations.
Am. Nat., 71 : 21 - 33.
- Park, T. 1962. Beetles, competition, and population. Science,
138 : 1369 - 1375.
- Park, T. and Frank, M.B. 1948. The fecundity and development
 of the flour beetles, Tribolium confusum and Tribolium
castaneum at three constant temperatures. Ecology,
29 : 368 - 375.
- Park, T., Mertz, D.B., Grodzinski, W. and Prus, T. 1965.
 Cannibalistic predation in populations of flour

- beetles. Physiol. Zool., 38 : 289 - 321.
- Phillips, W.M. 1978. Sensilla types from the ovipositor of the flea beetle, Attica lythri (Coleoptera : Chrysomelidae). Entomol. Exp. Appl., 24 : 599 - 601.
- Phillips, J.K. and Burkholder, W.E. 1981. Evidence of male produced aggregation pheromone in rice weevil, S. oryzae. J. Econ. Ent., 74 (5) : 539 - 542.
- Phillips, T.W., West, J.R., Foltz, J.L., Silverstein, R.M. and Lanier, G.N. 1984. Aggregation pheromone of the Deodar weevil, Pissodes nemorensis (Coleoptera : Curculionidae); Isolation and activity of grandisol and grandisal. J. Chem. Ecol., 10 (10) : 1417 - 1424.
- Polnik, A. 1960. Effects of some intra-species processes on competition between two species of flour beetles, Latheticus oryzae and Tribolium confusum. Physiol. Zool., 33 : 42 - 57.
- Pouzat, J. 1978. Host plant chemosensory influence on oogenesis in the bean weevil, Acanthoscelides obtectus (Coleoptera : Bruchidae). Entomol. Exp. Appl., 24 : 401 - 408.
- Powell, J.E., Jr. 1931. An ecological study of the tobacco beetle, Lasioderma serricorne (F.) with special reference to its life history and control. Ecol. Monogr., 1 (3) : 333 - 339.
- Prokopy, R.J. 1972. Evidence for a marking pheromone deterring

- repeated oviposition in apple maggot flies. Environ. Ent., 1 : 326 - 332.
- Prokopy, R.J. 1975. Oviposition deterring pheromone in Rhagoletis fausta. Environ. Entomol., 4 : 298 - 230.
- Prokopy, R.J., Reissing, W.H. and Moericke, V. 1976. Marking pheromones deterring repeated oviposition in Rhagoletis flies. Entomol. Exp. Appl., 20 (2) : 170 - 178.
- Prokopy, R.J., Greany, P.D. and Chambers, D.L. 1977. Oviposition deterring pheromone in Anastrepha suspensa. Environ. Entomol., 6 : 463 - 465.
- Prokopy, R.J., Ziegler, J.R. and Wong, T.Y. 1978. Deterrence of repeated oviposition by fruit marking pheromone in Ceratitidis capitata (Diptera : Tephritidae). J. Chem. Ecol., 4 : 55 - 65.
- Prokopy, R.J. 1981a. Oviposition deterring pheromone system of apple maggot flies; pp. 477 - 494. In : Management of Insect pest with Semiochemicals. eds. E.R. Mitchell, Plenum Press, New York.
- Prokopy, R.J. 1981b. Epideictic pheromones that influence spacing patterns of phytophagous insects; pp.181 -.213. In : Semiochemicals, Their role in pest control, eds. D.A. Nordlund., R. L. Jones and W. J. Lewis, Wiley Press, New York.
- Prokopy, R.J., Averill, A.L., Berdinelli, C.M., Bowdan, E.S., Cooley, S.S., Cranjar, R.M., Dundulis, E.A., Roitberg,

- C.A., Spalchar, P.J., Tunlinson, J.H. and Weeks, B.L.
1982. Site of production of an oviposition deterring
pheromone component in Rhagoletis pomonella flies. J.
Insect Physiol., 28 (1) : 1 - 10.
- Prokopy, R.J., Malavasi, A. and Morgante, J.S. 1982.
Oviposition deterring pheromone in Anastrepha
fraterculus flies. J. Chem. Ecol., 8 : 763 - 772.
- Prokopy, R.J., Roitberg, B.D. and Averill, A.L. 1984. Resource
partitioning; pp.301 - 330. In : Chemical Ecology of
Insects, eds. W.J. Bell and R.T. Carde, Sinauer
Associates, Massachusetts.
- Raina, A.K. 1981. Deterrence of repeated oviposition in
sorghum shootfly, Atherigona soccata. J. Chem. Ecol.,
7 (5) : 785 - 790.
- Rausher, M.D. 1979. Egg recognition : Its advantage to a
butterfly. Anim. Behav., 27 : 1034 - 1040.
- Renwick, J.A.A. and Radke, C.D. 1980. An oviposition deterrent
associated with the frass from feeding larvae of
cabbage looper, T. ni. (Lepidoptera : Noctuidae).
Environ. Entomol., 9 (3) : 318 - 320.
- Renwick, J.A.A. and Radke, C.D. 1981. Host plant constituents
as oviposition deterrents for the cabbage looper,
Trichoplusia ni. Entomol. Exp. Appl., 30 : 201 - 204.
- Reville, R. and Kannowski, P.B. 1975. Sexual excitation by
pheromones of the confused flour beetle. J. Insect

- Physiol., 21 : 81 - 84.
- Rice, M.J. 1976. Contact chemoreceptors on the ovipositor of Lucilia cuprina. Aust. J. Zool., 24 : 353 - 360.
- Rice, M.J. and Mcrae, T.M. 1976. Contact chemoreceptors on the ovipositor of Locusta migratoria (L.). J. Aust. Ent. Soc., 15 : 364 - 368.
- Rich, E.R. 1956. Egg cannibalism and fecundity in Tribolium. Ecology, 37 : 109 - 120.
- Rothschild, M. and Schoonhoven, L.M. 1977. Assessment of egg load by Pieris brassicae. Nature, London, 226 : 352 - 355.
- Ryan, M.F. and O'Ceallachain, D.P. 1976. Aggregation and sex pheromone in the beetle T. confusum. J. Insect. Physiol., 22 (11) : 1501 - 1504.
- Sakai, A., Honda, H., Oshima, K. and Yamamota, I. 1986. Oviposition marking pheromone of two bean weevils, Callosobruchus chinensis and Callosobruchus maculatus. J. Pesticide Sci., 11 : 163 - 168.
- Salt, G. 1934. Experimental studies in insect parasitism, Introduction and Technique. In : Chemical Ecology of insects, eds. W.J. Bell and R.T. Carde, Sinauer Associates, Massachusetts.
- Salt, G. 1937. Experimental studies in insect parasitism, the sense used by Trichogramma to distinguish between parasitized and unparasitized hosts. In :

- Chemical Ecology of Insects, eds. W.J. Bell and R.T. Carde, Sinauer Associates, Massachusetts.
- Schoonhoven, L.M., Sparnaay, T., Wissen, W.V. and Meerman, J. 1981. Seven week persistence of an oviposition deterring pheromone. J. Chem. Ecol., 7 : 583 - 588.
- Shaaya, E. 1981. Sex pheromone of Dermestes maculatus (Coleoptera : Dermestidae). J. Stored Prod. Res., 17 : 13 - 16.
- Sokoloff, A. 1972. "The biology of Tribolium". Oxford University Press. Vol.1.
- Sokoloff, A. 1974. "The biology of Tribolium". Oxford University Press, Vol.2.
- Sonleitner, F.J. 1961. Factors affecting egg cannibalism and fecundity in populations of adult Tribolium castaneum (Herbst). Physiol. Zool., 34 : 233 - 255.
- Sridevi, R., Aparna, R. and Ramamurty, P.S. 1987. An oviposition stimulant in the male accessory gland extract of Spodoptera litura (Lepidoptera : Noctuidae). Entomon., 12 (3) : 231 - 234.
- Stadler, E. 1984. Contact Chemoreception; pp. 3 - 55. In : Chemical Ecology of Insects, eds. W. J. Bell and R. T. Carde, Sinauer Associates, Massachusetts.
- Suzuki, T. and Sugawara, R. 1979. Isolation of an aggregation pheromone from the flour beetles, T. castaneum and T. confusum (Coleoptera : Tenebrionidae). Appl.

- Entomol. & Zool., (Jap.). 14 : 228 - 231.
- Suzuki, T. 1980. 4, 8-Dimethyldecanal : The aggregation pheromone of the flour beetles, T. castaneum and T. confusum. (Coleoptera : Tenebrionidae). Agric. & Biol. Chem., 44 : 2519 - 2520.
- Suzuki, T. 1981. Identification of aggregation pheromone of the flour beetles Tribolium castaneum and T. confusum (Coleoptera : Tenebrionidae). Agric. Biol. Chem., 45 (6) : 1357 - 1363.
- Suzuki, T. 1985. Presence of another aggregation substance(s) in the frass of the red flour beetle, Tribolium castaneum (Coleoptera : Tenebrionidae). Appl. Ent. Zool., 20 (1) : 90 - 91.
- Szentesi, A. 1976. The effect of amputation of head appendages on the oviposition of bean weevil, Acanthoscelides obtectus (Coleoptera : Bruchidae). Abst. Symp. Biol. Hung., 16 : 275 - 281.
- Szentesi, A. 1981. Pheromone like substances affecting host related behaviour of larvae and adults in the dry bean weevil, Acanthoscelides obtectus. Entomol. Exp. Appl., 30 : 219 - 226.
- Tabashnik, B.E., Wheelock, H., Rainbolt, J. D. and Watt, W.B. 1981. Individual variation in oviposition preference in the butterfly, Colias eurytheme. Oecologia (Berlin), 50 : 225 - 230.

- Tanaka, K., Ohsawa, K., Honda, H. and Yamamota, I. 1981.
Copulation release pheromone, Erectin, from the
Azuki bean weevil (Callosobruchus chinensis L.).
J. Pesticide Sci., 6 : 75 - 82.
- Tingle, F.C. and Mitchell, E.R. 1984. Aqueous extracts
from indigenous plants as oviposition deterrents for
Heliothis virescens (F.). J. Chem. Ecol., 10 :
101 - 113.
- Touchstone, J.E. and Dobbine, M.F. 1978. Practice of thin .
layer chromatography. pp. 1 - 345. John Wiley and
Sons. Inc. New York.
- Umeya, K. 1966. Studies on the comparative ecology of bean
weevils. 1. On egg distribution and the oviposition
behaviours of three species of bean weevils infest-
ing Azuki bean. Res. Bull. Plant Prot. Service,
Japan, 3 : 1 - 11.
- Valencia, L. and Rice, M.J. 1982. Contact chemoreceptors
on the ovipositor of the potato moth, Phthorimaea
operculella (Lepidoptera : Gelechiidae). Int. J.
Insect Morph. and Embryol., 11 : 121 - 128.
- Waladde, S.M. 1983. Chemoreception of adult stem borers :
Tarsal and ovipositor sensilla on Chilo partellus
and Eldana saccharina. Insect Sci. Appl., 4 : 159-165.
- Wasserman, S. 1981. Host induced oviposition preference
and oviposition marker in Cowpea weevil,

Callosobruchus maculatus. Ann. Ent. Soc. Amer.,

74 : 242 - 245.

Wasserman, S. 1985. Oviposition behaviour and its disruption in the southern cowpea weevil, Callosobruchus maculatus F. (Coleoptera : Bruchidae). J. Econ. Ent.,
78 : 89 - 92.

Wiedner, H. 1979. Der Tabakkafer Lasioderma serricorne F.
Der Praktische Schadlingsbekämpfer Bl. 129 - 132.

Wood, D.L., Silverstein, R.M. and Nakasima, M. 1970. "Control of insect behaviour by natural products". Academic Press, New York.

Yamamoto, R.T. and Fraenkel, G. 1960. The suitability of tobaccos for the growth of the cigarette beetle, Lasioderma serricorne. J. Econ. Ent., 53 (3) : 381-384.

Yoshida, T. 1961. Oviposition behaviour of two species of bean weevil and interspecific competition between them, Mem. Fac. Lib. Arts. Educ. Miyazaki Univ.,
11 : 41 - 66.

Zimmerman, M. 1979. Oviposition behaviour and existence of an oviposition deterring pheromone in Hylema. Environ. Entomol., 8 (2) : 277 - 279.

Zyromska-Rudzka, H. 1966. Abundance and emigrations of Tribolium in a laboratory model. Ekol. Pol. A.,
14 : 491 - 518.