

BIBLIOGRAPHY

BIBLIOGRAPHY

- Ablas Kimyai, 1993. Eocene palynomorphs from the Black diamond mines Regional Preserves, contra costa country California, palynology (AASP) vol.17, : 101-114.
- Adegoke O.S.Chene, R.E., J. Du Agumanu, A.E., 1978. Palynology and age of the Kerri-Kerri formation, Nigeria, Revta, esp. Micropaleont, 10:267-283.
- Agashe, S.N., 1969 studies on the fossil gymnosperms of India Part-I. A new species of *Mesembrioxylon*, M.Mahabalei sp. Nov. paleobotanist, 17(3) 31-316.
- Ambwani K., 1982 Palynology of the Deccan Intertrappean Beds of Rajahmundry District, Andhra Pradesh, Paleobotanist 30(1); 28 -33.
- Anderson, R.Y., 1960. Cretaceous-Tertiary palynology eastern side of the San Juan basin New Mexico Mem. New Mex, Bur.Min Minder resources, 6.
- Archangelsky, S. 1969. Estudio del paleomicroplancton de la Formacion Rio Turbio (Eocene), Provincia de Santa Cruz, Amegbiniada 6(3): 181-218.
- Baksi, S.K., 1962. Palynological investigation of Simsang river Tertiaries, South Shillong Front, Assam. Bull.Geol.Min.Metall.Soc. India, 26: 1-22.
- Baksi, S.K., 1965. Stratigraphy of Barail Series in the southern part of Shillong Plateau, Assam, India. Bull.Amer.Assoc., Petrol.Geol.Vol.49, no.12.
- Baksi, S.K., 1968. Siwalik microflora from Punjab, India. Rev.Palaeobot Palynol 6: 171-176.
- Baksi, S.K., 1972 on the palynological biostratigraphy of Bengal basin. Seminar Paleopalyn. Indian Stratigr., Calcutta, :188-205.
- Baksi, S.K., 1974. Significant pollen taxa in the stratigraphical analysis of the Tertiary sediments of Assam (in Asp. and Appro. of Ind.Palaeob). spl. silv. Jub.Publ.B.SIP, :502-515.
- Baksi, S.K. and Deb, U. 1980. Palynostratigraphic zonation of the Upper Cretaceous Paleogene sequence of Bengal basin. Geophytology, 10:199-224.
- Banerjee, D., 1967. Significance of saccate grains in the Tertiary stratigraphy of Assam. Seminar on Geology of NE India, Shillong (Abstract), p.18.
- Banerjee, D., 1968. Siwalik microflora from Punjab, India. Rev. Palaeobot,Palynol. 6:171-176.
- Banerjee, D. and Mishra, C.M., 1971. Hystrichospherids in the Tertiary formations of Assam and Tripura. Seminar on Palaeopalynology, Calcutta.

- Banarjee, D., 1973. Palynological evidences and the Tertiary sediments of India. Journal of the Paleontological Society of India, Vol.17, 1972, :1-6.
- Bennett, J., 1989. Palynology and paleoecology of the Tertiary weaverville formation, Northwestern California, USA palynology 13:195-246.
- Biswas, B., 1962. Stratigraphy of the Mahadeo, Langpar, Cherra and Tura formations, Assam, India. Bull. Geol.min.metal.Soc.India, 25:1-48.
- Biswas, S.K. & D.S.N.Raju, 1973. The rock stratigraphic classification of the Tertiary sediments of Kutch, Bull., ONGC, Vol.1, No.1-2, :37-45.
- Brosins, M., 1963. Plankton aus dem nordhessischen kasseler Meeressad (ober-oligozan) Z. dt. geol. Ges. 114:32-56.
- Bujak, J.P., 1980. Dinoflagellate cysts and Acritarchs from the Eocene Barton beds of southern England. In Bujak, J.P., Downie C, Eaton, G.L. and Williams, G.L.Dinoflagellate cysts.
- Bujak, J.P.Downie, C. Eaton G.L. & Williams G.L., 1980. Dinoflagellate cysts and acritarchs from the Eocene of southern England. Spec. Publ. Paleont. London 24:100.
- Benedek, P.N., 1972. Phytoplankton aus dem mittal and oberoligozan von tonisberg (Niederrheingebiet), Palaeontographica, Stuttgart B 137 (1-3) : 1-71.
- Bennie, J and Kidston R., 19886 on the occurrence of spores in the carboniferous formation of Scotland. Proc, Roy.Soc.Edinb., 9:82-117.
- Caro, Y., 1973. Contribution a la connaissance des dinoflagelles du Paleocene-Eocene inferieur des Pyrenees Espangnoles. Rev. espanola Micropalaeont. 5(3):329-372.
- Chandra, P.K. & Chaudhary L.R., 1969. Stratigraphy of the Cambay Basin. Bull. ONGC Vol.6 No.2, pp.37-50.
- Chitaley, S.D., 1951. Fossil microflora from the Mohgaonkhan beds of M.P., India. Proc., Natn.Inst., Sci., India, 17(5):373-383.
- Clarke, R.T. & Friederiksen, N.O.(1968). Some new sporonomorphs from the upper tertiary of Nigeria, Grana palynol.8(1):210-224.
- Cookson, I.C., 1947. Plant microfossils from the lignites. Reports, Ser.A.II (8) 17-142; pl.XIII-XVII.
- Cookson, I.C., 1950. Fossil pollen grains of Protensous type from Tertiary deposits in Australia Aust.J. Sci.Res.Ser.Biol,sci 3(2):166-177.
- Cookson, I.C., 1953. Records of the occurrence of Botryococcus braunii, Pediasium and the Hystrichosphaerideae in Cainozoic deposits of Australia, Memoire of the National Mussum, M.labourne, 18:107-123.

- Cookson, I.C. & Pike, K.M., 1954 . Some dicotyledonous pollen types from Cainozoic deposits in the Australian region Aust.J.bot.2:197-219.
- Cookson, I.C., 1965. Cretaceous and tertiary microplankton from south-eastern Australia. Proc.R.Soc.Vict.78:85-93.
- Cookson, I.C. & Eisenack A., 1965. Microplankton from the Palaeogene pebble point formation, south-western Victoria Proc.R.Soc.Vict. 79, 139-146.
- Cookson, I.C. & Eisenack, A. 1967. Some microplankton from the palaeogene river nook bed victoria, Proc.R.Soc.Vict., 80, 247-257
- Cookson, I.C. & Eisenack, A., 1967. Some early Tertiary microplankton and pollen grains from a deposit near Strahan Western Tasmania, Proc.R.Soc. Vict. 80(1) : 131-140.
- Couper, R.A., 1953. Upper Mesozoic and Cainozoic spores and pollen grains from New Zealand, New Zealand Geological Survey Paleontological Bulletin 22:1-77.
- Couper, R.A., 1960. New Zealand Mesozoic and Cainozoic plant microfossils, N.Z.geol.surv.palaeont., Bull.32:1-87.
- Davey, R.J. & Williams, G.L., 1966. The genus *Hystrichosphaeridium* and its allies, in Mesozoic and Cainozoic dinoflagellate cysts by Davey, R.J., Downie, C., Sarjeant, W.A.S. & Williams, G.L., Bull.Br.Mus.nat.Hist. (Geol.), 3:1-248.
- Deb, U., 1972. Some pollen grains from the Neyveli lignite: pp.220-228 in Ghosh A.K. et al. (Editors) proceedings of the seminar on Paleopalynology and Indian stratigraphy, Calcutta, 1971.
- Davey, R.J., Downie, C., Sarjeant, W.A., and Williams, M.S.G.L., 1966. Studies on Mesozoic and Cainozoic dinoflagellate cysts. Bull.Br.Mus.nat.Hist. Suppl.London, 3:1-248.
- De conick, J., 1965. Microfossils planktoniques du sable Ypresien à merelbeke inophyceae et Acritarcha Mem. Acad r. Belg. 36(2) : 1-54
- De conick, J., 1967 in Moorkens T.L. et al., 1967. Het fossiel bounded ieperiaan van merelbeke matuurw Tijdechr 48: 202-227
- De connick, J., 1968. Dinophyceae et Acritarcha de Iypresien du sondage de kallo, Roy sci Not mem Belg 161i:1-67.
- Deb U., Baksi, S.K. & Ghosh, A.K., 1973 on the age of Neyveli lignite a palynological approach, QJI Geol.Min.Metall soc.India 45:23-38.
- Deflandre, G., 1937 . *Phanerodinium*, genera nouveau de dinoflagelle fossile des silex, Bulletin de la societe Francaise de microscopic 6:109-115.
- De conick, J. (1972) Application stratigraphique des microfissiles organiques dans Iypresien du bassin belge. bull. soc. Belg. Geol. paleont Brussels 81 (1,2) : 1-11.

- Deflandre, G. & Cookson, I.C., 1955. Fossil microplankton from Australian Late Mesozoic and Tertiary sediments. Austr.J. marine fresh water Res. 6:242-313.
- Delcourt, A.F. & Sprumont, G., 1955. Les spores et grains de pollen du Wealdien du Hainaut, Mem.Soc.Belge, Geol.4:73.
- Delcourt, A.F.Dettmann, M.E. & Hughes, N.F., 1963. Revision of some Lower Cretaceous microspores from Belgium Palaeontology, 6:82-292.
- Dettman, M.E., 1963. Upper Mesozoic microfossils from south eastern Australia Proc.R.Soc.Vict. 77(1):1-148.
- Dilcher, D.L. 1965. Epiphyllous fungal from Eocene deposits in western Tennessee, U.S.A., Palaeontographica 116B : 1-54.
- Daniel Michoux, 1988. Dinoflagellate cysts of the *Wetzelieilla* complex from Eocene sediments of the Aquitaine Basin south western France palynology (AASP) vol.12, : 11-12.
- Downie, C., Evitt, W.R., and Sarjeant, W.A.S., 1963. Dinoflagellates hystrichospheres and the classification of the acritarchs Stanford Uni.Pubs., Geol.Sci.v.7, :1-16.
- Downie, C. and Sarjeant, W.A.S., 1963. On the interpretation and status of some hystrichophore genera . Paleontology. V.6, :83-96.
- Downie, C, Hussain, M.A. & Williams, G.L., 1971. Dinoflagellate cyst and acritarch associations in the Palaeogene of southeast England. Geosci.Man 3:29-35.
- Drugg, W.S., and Loeblich, A.R.Jr., 1967. Some Eocene and Oligocene phytoplankton from the Gulf Coast, USA, Tulane studies geology, 5:181-94.
- Dutta, S.K. & Sah, S.C.D., 1967. Palynostratigraphy of the sedimentary formation of Assam. 4: Age of the Laitryngew-Mawkma coal-bearing sandstone and their relationship with the Cherra foramtion. Seminar on Geology of NE India, Shillong (Abstract), 10.
- Dutta, S.K., 1974. Palynology of the Tertiary sediments of Palana, Rajasthan -Palaeobotanist, 21:163-188.
- Dutta, S.K. & Jain, K.P., 1980. Geology and palynology of the area around Lumshnong, Jaintia Hills, Meghalaya, India. Biol.Mem.5(1):56-81.
- Dutta, S.K. & Singh, H.P., 1980. Palynostratigraphy of sedimentary foramtion in Arunachal Pradesh.1 Palynology of Siwalik rocks of the lesser Himalaya, Kameng District, pp 617-622 in Bharadwaj, D.C. et al. (Editors - Proc 4th int.Palynol.conf., Lucknow, 1976-77.2 Birbal Sahni Institute of Paleobotany, Lucknow.

- Eaton, G.L., 1971. A morphogenetic series of dinoflagellate cysts from the Bracklesham Beds of the Isle of the wight, Hampshire, England in Farinacci, A (Ed) proceedings secone planktonic conference Rome 1970 Edizioni technoscience, 1: 355-379.
- Eaton G.L. 1976. Dinoflagellate cysts from the Bracklesham beds (Eocene) of the Isle of wight, southern England, Bull.Br.Mus.nat.Hist.(Geol),London, 26(6) : 227-332.
- Erdtman, G. 1952. Pollen morphology and plant taxonomy,Angiosperms, Vol.1.
- Edwards, W. 1922. An Eocene microthyriaceous fungus from mull scotland Trans Br. Mycol Soc. 8:66-72.
- Ehenberg, C.G. (1838) Über das Massenverhaltniss der Jetzt lebenden kiesel infusorien and über ein neues infusorien-conglamerat als polierschiefer von Jastraba in Ungarn Abhandlungen der Preussischen Akademie der Wissenschaften 1836, 109-135.
- Eisenack, A. 1938. Die phosphoritknollen der Bernsteinformation also Oberlief-erer tertiären Planktons. Schrift Physikalisch ökonomischen Gesellschaft Königsber, 70: 181-188.
- Eisenack, A. 1958. Microplankton aus dem Norddeutschen Apt, nobst einigen Bemerkungen liber fossile dinoflagellaten N.Jb Palaont. Abh. 106:383-422.
- Eisenack, A. 1963. Cordosphaeridium n.g. ex. Hystrichosphaeridium, Hystricho-sphaeridea Neues Jb Geol. 118: 260-265.
- Elsik, W.C. 1968. Palynology of Palaeocene Rockdale Lignite, Milam Country, Texas. I Morphology and taxonomy. pollen spores 10(2) : 263-314.
- Erdtman, G. 1947. Suggestions for the classification of fossil and recent pollen grains and spores. Svensk.bot.Tulkr.41(1):104-114.
- Erdtman, G. 1943. An introduction to pollen analysis Waltham, Mass, USA.
- Evitt, W.R. 1963. A discussion and proposal concerning fossil dinoflagellates, Hystrichosphaeres and Acritarchs, Proc. Natu. Sci. U.S.A., 49:158-164.
- Frederiksen, N.O. 1980. Sporomorphs from the Jackson group (Upper Eocene) and adjacent strata of Mississippi and western Alabama. - U.S.Geol. Surv.Prof.Pap. 1084 : 75 p., 16 pl.
- Faunnel, B.M. 1964. The Tertiary period in W.B. Harland, A.Gilbel Smith and B.Wilcock The phenerozoic time scale : A symposium dedicated to professor A. Holmes suppl.Quart.J.Geol.Soc.London 120:179-191.
- GeiLach, E. 1961. Microfossilien aus dem oligozan and Miozan Nordwestdeutschlands, unter besonderer Berucksichtigung der Hystrichosphaerideen and Dinoflagellaten N.Jb geol. Palaont (Abstract) 112:143-228.

- Germeraad T.H., C.A. Hopping and Muller J. 1968. Palynology of Tertiary sediments from tropical area, Bench mark papers in Geology V.4b, palynology Part-I spore and pollen, : 184-220.
- Germeraad, J.M., Hopping C.A. and Muller, J. 1969. Palynology of Tertiary sediments from tropical areas. Rev. Palaeobotan. Palynol. 6:189-348.
- Ghosh, A.K. and Srivastava, S.K. 1962. Microfloristic evidence on the age of Krol beds and associated formations, Proc, Nor, Inst, Sci, India 28A 5:710-717.
- Ghosh, A.K. S.K. Srivastava and J.Sen. 1963. Polycolpate grains in Pre-Miocene horizons of India. Proc. Nat. Inst. Sci Ind, 29B (5):511-519.
- Gruas Cavagnetto 1968. Etude Palynologique des divers gisements du sparacien du bassin de paris. mem soc. geol. fr. (NS) 47 (2) 143-228.
- Gruas Cavagnetto, C. 1970c. Microflore et microplankton des woolwich beds (Swanscombe, Kent) Pollen spores 12(1) : 71-82.
- Gruas Cavagnetto C. 1970b. Dinophyceae, Acritarcha et pollens de la Formation de verengewille (Cuisien, science maritime) Rev. micropalaeon.13(2):30-78
- Gruas Cavagnetta C. 1971. Presence de microplankton et de pollens dans le Lutetien du bassin de paris, C.r.somm, seanc soc. geol.Fr. 1971(3):172-174
- Guzman, A.E.G..1967. A Palynological study on the upp.Los Cuervos and Mirador Formations (Lower and Middle Eocene, Tibue area, Colombia) Akad, Proefs:1-68
- Haseldonkx, P. 1974. A Palynological interpretation of Paleo-Environment in S.E.Asia sains Malaysiana 3(2):119-127.
- Habb, P. 1971. Dinoflagellate stratigraphy across the Miocene-Pliocene boundary Tabiono stratotype section In:A Farinaci (ed) proceedings of the second planktonic conference, Rome 1970. 1. 591-8.
- Hedberg, H.D. 1976. International Stratigraphic Guide, John Wiley and Sons. New York.
- Hekel, H. 1972 - Pollen and spore assemblages from Queensland Tertiary Sediments - Geol.Surv.Qd.Publ. 355 (Palaeontol. Pap.30) : 34 : 6 pl.
- Hopkins W.S, and Rutter and Rouse G.E. 1972. Geology, Paleocology and Palynology of some Oligocene rocks in the rocky mountain Treach of British Columbia canadian journal of earth schience, volume-9, number-4, pp 460-470.
- Ibrahim, A.C. 1933. Sporenformen des Aegirhorizonts des Ruhr-Reviers, Diss Konard Trittsch, Wurzburg 1-47.
- Islam, M.A. 1983. Dinoflagellate cysts from Eocene of the London and Hampshire basins, southern England palynology (AASP) Volume-77 : 71-92.
- Ivevson, J and Troels-Smith J. 1950. Pollen morphologiske definioner og typer Domm. geol Unders IV 3(8) 1-52.

- Jansonius, J. & Hills, L.V. 1976. Genera file of fossile spores spl, pub. dept. geology, univ. calgary Canada 1-3287.
- Jansonius, J.K. & Hills L.V. 1977. Genera file of fossil spores supplement spl.pub.dept geology univ. calagary Canada 3288-3431.
- Jansonius J.K. & Hills L.V. 1978. Genera file of fossil spores supplement spl. pub. dept, geology, univ.calagary, Canada 3432-3520.
- Janasonius, J.K. & Hills L.V. 1979., Genera file of fossil spores supplement spl.pub. dept. geology univ. calagary, Canada 3521-3628.
- Janasonius, J.K. & Hills 1981. Genera file of fossil spores-supplement, spl. pub. dept. geol, univ. Calgary, Canada : 3801-3932.
- Janasonius, J.K. & Hills L.V. 1982. Genera file of spores-supplement spl.. pub. dept. geology univ. Calagary Canada-3933-4056.
- Janasonius, J.K. & Hills L.V. 1987. Genera file of fossil spores-supplement spl. pub. dept. geology univ. calgary, Canada 4361-4676.
- Jain, K.P. & Gupta, R.C. 1970. Some fungal remains from the Tertiaries of Kerala coast, paleobotanist 18(2) : 177-182.
- Jain, K.P. Kar, R.K. and Sah, S.C.D. 1973. A palynological assemblage from Barmer Rajasthan Geophytology 3(2):150-165.
- Jain, K.P., 1977b. Additional dinoflagellates and acritarchs from Grey Shale Member of Dalmiapuram Formation, South India, Palaeobotanist 24(3) : 170-194.
- Jain, K.P. and Kar, R.K.1979. Palynology of Neogene Sediment around Quilon and Varkala, Kerala Coast, South India-1 Fungal remains Palaeobotanist 26(2) : 105-118.
- Jain, K.P. & Tandon, K.K. 1981. Dinoflagellage and acritarch biostratigraphy of the Middle Eocene rocks of a part of south-western Kutch, India, J.palaeont, soc.India. 26:6-21.
- Jain, K.P. and Garg R. 1986b. Upper Paleocene dinoflagellate cysts and acritarch from Virdhachalam, cauvery Basin Southern India. Palaeontographica 190B (4-5) 61-68.
- Jeanie Barnett 1989. Palynology and paleoecology of the Tertiary Weave ville Formation, north western California USA palynology (AASP) vol.13, pp 195-246.

- Kar, R.K., Singh R.Y. and Sah S.C.D. 1972. On some algal and fungal remains from Tura Formation of Garo Hills, Assam, Palaeobotanist, 19(2):146-154.
- Kar, R.K. & Saxena R.K. 1976. Algal and fungal microfossils from Matanomadh Formation (Palaeocene) Kutch, India, Paleobotanist, 23(1):1-15.
- Kar, R.K. 1978. Palynostratigraphy of the Nardi (Lower Eocene and the Harudi (Middle Eocene) formations in the district of Kutch, India, Paleobotanist 26(1):16-49.
- Kar, R.K. 1979. Fossil algae from fulra limestone (Middle Eocene), Kutch, Gujarat Geophytology 9(1): 88-90.
- Kar, R.K. & Jain, K.P. 1981. Palynology of the Neogene sediments around quilon and Varkala Kerala coast, south India-2 spores and pollen grains paleobotanist, 27(2):113-131.
- Kar, R.K. & Saxena, R.K. 1981, Palynological investigations of a bore core near Rataria, southern Kutch, Gujarat Geophytology, 11(2):103-124.
- Kar, R.K. 1985. The fossil floras of Kutch-IV, Tertiary palynostratigraphy paleobotanist 34:1-280.
- Kar, R.K. & Kumar, M. 1986. Paleocene palynostratigraphy of Menghalaya, pollen spores 28:177-217.
- Kapoor, R. & Singh, R.Y. 1987. A note on the geology and distribution of some significant fossils in the Lower Tertiary sediments exposed along Kalka-Kasauli Road section Bull Indian Geologists Assoc.20(11):17-23.
- Khan, A.M. & Martin, A.R.H. 1971. A note on genus polypodiisporites R.Potonie. pollen spores 3:475-480.
- Kedver, M. 1968. Etudes palynologiques des Cruches du tertiaire inferieur de la region parizienne III.pollen & spores, 10:315-334.
- Klement, K.W. 1960. Dinoflagellaten und Hystrichosphaeriddeen aus dem unteren and Mittleren Malm Sudwestdeutschland, Palaeontographica, 114(A) 1-104.
- Klumpp, B. 1953. Beitrag zur kenntnis der microfossilien des mittleren und oberen Eozan Palaeontographica 103(A):377-406.
- Krutzsch, W. 1966. Zur kinntnis der präquartären periporaten pollenformen, Geologie, 55:16-71.
- Koshal V.N. 1980. Palynology of the Tertiary subsurface of Dabka, Gujarat: pp 686-690 in Bharadwaj et al Proc. 4th int. palyno., Lucknow 1976-77, Birbal Sahni Institute of Paleobotany Lucknow.
- Koshal, V.N. 1988. Palynofossils and palaeoenvironment of the Tertiary subsurface sediments in Ahmedabad-Mehsana Block, Cambay Basin, Western India, pp 621-672 in Whyte P. et al (Editors). The Palaeoenvironment of East Asia from Mid-Tertiary-1,centre of Asian. Studies Univ., Hong Kong.

- Kremp, E.M. 1968. Probable angiosperm pollen from British Barremian to Aptian strata palaeontology 11:421-434.
- Krutzsch, W. 1959. Einige neue formgattungen und Arten von sporen und pollen aus der mitteleuropaischen oberkreide und dem Tertiär, palaeontographica, 105 B:125-157.
- Leidelmeyer, P. 1966. The Paleocene and Lower Eocene pollen flora of Guyana, Leidse geol. Meded. 38, 49-70.
- Loeblich, A.R. Jr and Loeblich, A.R.III 1966. Index to Genera, subgenera and sections of the Pyrrhophyta: studies in Tropical oceanography. Miami no.3 x:94
- Luber A.A. 1955. Atlas of spores and pollen of palaeozoic sediments in Kazakhstan, An.kas, SSR Ahma-Atz.
- Luckose, N.G. 1968. Microfossils from the Middle Siwalik, Bihar, India, J.Palyno. 4(2), 107-112.
- Madan Mohan, S.K.Bhattacharya, D.C.Sharma, C.L.Chowdhary and M.Shanmukhappa, 1987. Environmental reconstruction and depositional modelling of the Hazad pay sands in Gandhar field. ONGC Report (unpublished).
- Maier, D. 1959. Planktonuntersuchungen in tertiären und quartären marinen sedimenten Nöuse Jahrbaah fir geologic and Palaontologic, Abhurdlingen 107 : 278-340.
- Mantell, G.A. 1950. A pictorial atlas of fossil remains consisting of coloured illustrations selected from parkinsons "Organic remains of former world" and Artis Antediluvian phytology, London.
- Martin, H.A. and G.E.Rouse 1966. Palynology of late Tertiary sediments from Queen Charlotte Islands, British Columbia; Canadian Jour. Botany.Vol. 44, p. 171-208.
- Mathur, L.P., Rao, K.L.N. and Chaube, A.N. 1968. Tectonic framework of Cambay Basin, Bull. Oil & Natural Gas Commission Vol.5, No.1.
- Mathur, L.P., Rao K.L.N. and Chaube, A.N. 1980. Tectonic framework of Cambay Basin India. Bull. ONGC, Vol.17, No.2.
- Mathur, Y.K. 1963. Studies in the fossil microflora fo Kutch, India-1, on the microflora and the hystrichosphaerids in the gypaeous shale (Eocene) of Western Kutch, India. Proc.Nat.Inst.Sci.Ind.29B(3), 356-371.
- Mathur, Y.K. 1966. On the microflora in the Supra-Trappeans of Western Kutch, India : Quart. Jour. Geol.Min.Met.Sec.Ind.v.38, p.33-51.
- Mathur, Y.K. 1972. Plant fossils from the Kaur bet, Paccham island, Kutch curr. sci., 41(3):488-489.
- Mathur, Y.K. and Pant, J. 1973. Plant microfossile of palaeocene-Eocene age from the Wagad region, Kutch, curr.sci.42(24):857-859.

- Mathur, Y.K. and Choudhary, L.R. 1976. Paleoecology of the Kalol formation, Cambay basin, India. Proc.IV Colloq, Indian Micropal & Strat.Pub. IPE, ONGC, Dehradun.
- Mathur, Y.K. et al., 1977. Paleoecology of the Kalol Formation, Cambay basin, India, Proc.IV Colloq, Indian microplaeontol. & Strat., IPE, ONGC Dehradun.
- Mathur, Y.K. and Mathur, K. 1980. Barail (laisong) palynofossils and late Oligocene nannofossils from the Andman Island, India, Geosci. Jl. 1(2), p 51-65.
- Mathur, Y.K. and Jain, A.K. 1980. Palynology and age of the Dras volcanics near Shergol, Ladakh, Jammu and Kashmir, India. Geoscience Journal, 1(1):55-74.
- Mathur, Y.K. and A.S.Chopra 1982. Palynology, Morphology and Taxonomy of the cryptogamic spores from the post paleogene subsurface sediments of the Bengal Basin, India, Geoscience Journal, Vol.III no:1,pl.3,pp. 51-80.
- Mathur, Y.K. 1986. Dinoflageelate cyst Biostratigraphy and age of the Middle Eocene Kalol Formation in the Kalol 109 north Cambay Basin, Western India, Review of Palaeobotany and Palynology, 47: 193-202.
- Mathur, Y.K. & Chopra, A.S. 1987. Palynofossils from the Cenozoic subsurface sediments of the Bengal Basin, India, Geoscience Journal, Vol.III, : 109-152.
- Mathur, K., Singh, R.N. and Mathur, Y.K. 1991. Standard laboratory techniques and procedures in Geology, Publ.K.D.Malviya Institute of Petroleum Exploration, Dehradun,(Editors Pandey, J. & Rao, V.K.) : 83-90.
- Mehrotra, N.C. 1981. Palynological correlation of Mikir Formation with Lower Palaeogene sediments of shillong Plateau. Geophytology 11(2) : 133-142.
- Mehrotra, N.C. and Sah S.C.D. 1981. Mikir Formation of Mikir-North Cachar Hills, Assam, its stratigraphic status, age and geological evolution of the depositional basin. Bull, Indian Geologist Assoc. 14(1):1-13.
- Mehrotra, N.C. and Sah, S.C.D. 1982. Palynostratigraphy of Mikin Formation in the type area, Geosci.Jl 3(2),p 113-134.
- Mehrotra, N.C. 1983. Palynology of Mikir Formation in the type area. Geoscience Journal, Vol.IV, no.1,:1-34.
- Mehrotra, R.B. and Ramakrishna, V 1981. Stratigraphy of Cambay Basin at a glance. Unpubl. report ONGC.
- Miner, E.L., 1935. Paleobotanical examinations of Cretaceous and Tertiary coals, American Midland Naturalist, 16:585-525.

- Mohan M. 1988. Palaeoenvironment analysis and Source Rock potential of Paleocene -Eocene succession in Well Jambusar-P; ONGC Report (Unpublished).
- Morgenroth, P. 1966a. Microfossilen und konkretioen des nord-westeuropaischen untereoiziins. *Palaeontographica B* 119:1-53.
- Morgenroth, P. 1966b. Neue in organischer substanz erhaltene microfossilen des Oligozons. *N.Jb. Geol.Palaont*, 127(1):1-12.
- Morgenroth, P. 1968. Zur kenntnis der dinoflagellaten and hystrichosphaeridien des Danien. *Geol. Jb.* 86:533-578.
- Muller, J. 1959. Palynology of recent attributable to Alangium Lamarck (Alangiaceae) from the Tertiary of Malesia Rev. *Paleobot.Palynol.* 36:65-94.
- Muller, J. 1968. Palynology of the Padawan and Plateau Sandstone formations (Cretaceous-Eocene) in Sarawak, Malaysia. *Micropalaeontology*, 14:1 37.
- Nandi, B. 1972. Some observations on the microflora of Middle Siwalik sediments of Mohan (East) field, U.P. *Proc. Sem.Palynol.,Ind. Strat. Calcutta* 1971:375-383.
- Nandi, B. 1975. Palynostratigraphy of the Siwalik Group of Punjab, Him. *Geol.* 5: 411-424.
- Nandi, B. 1980. Further contribution on the Palynostratigraphy of Siwalik group. pp 727-734 in Bhardwaj D.C. et al (Editors) *Proc 4th int. Palynol. Conf.* 1976-77.2, Birbal Sahni Institute of Paleobotany,Lucknow.
- Nandi, B. 1981. Microfloral investigation on the Neogene formations of Moren and Nahorkatiya wells, Upper Assam, India, *Geophytology*, 11:27-40.
- Nandi, B & Sharma R. 1984. Palynology and biostratigraphy of the Boldamgiri Formation Gora Hills Meghalaya pp 565-580 in Sharma A.K. et al., proceedings of the symposium on evolution botany and biostratigraphy Calcutta 1979 A.K.Ghosh commemoration volume current trends in life sciences-10.
- Naskar P. & Baksi S.K. 1978. Palynological investigation of Akli lignite, Rajasthan India, *Paleobotanist* 25: 314-329.
- Naumova, S.N. 1939. The spores and pollen of the coals of USSR Rec. 17th Int. *Geol.Congr. Moscow*,1:553-364.
- Navale, G. K.B. & Mishra B.K. 1979. Some new pollen grains from Neyveli lignite Tamilnadu, India *Geophytology*-8(2):226-239.
- Navale, G.K.B. 1961 - Pollen and spores from Neyveli lignite, south India - *Palaeobotanist*, 10:87-90.
- Nichols, D.J. et al., 1973. On arecipes Wodehouse monocolpopollenites Thomson & Pflug and the species Monocolpopollenites tranquillus *Taxon* 22:241-256
- Norris, G. and W.A.S. Sarjeant 1965. A descriptive index of genera of fossils Dinophyceae and Acritarcha New Zealand *Geol. Surv .Paleont.Bull.*40, 72 pages.
- Narendra P., Pandey, A.N., Chaudhary C.L.& Upadhyay H, 1988. Lithofacies, depositional

- environments and reservoir characteristics of Paleogene strata in well Jambusar : ONGC Report (Unpublished).
- Omar A. Colmenares and Lizaveta Teran, 1993. A biostratigraphic study of Paleogene sequence in south western venezuela palynology (AASP) Vol.17, : 67-90.
- Pandey, J., Singh N.P., Krishna B.R., Sharma, O.D.Paraikh, A.K. and Nath, S.S., 1993. Lithostratigraphy of Indian Petroliferous Basins, Cambay Basin. Publ. KDM, IPE, ONGC, Dehradun.
- Pastiels, A. 1948. Contribution in 'l'etude des microfossils de 'l' Eocene Belge Mem, Mus.R. (Not Hist) Belg. 109: 1-71.
- Patel, R.J. & George, I, 1977. Chlorococcales of Gujarat, India-pediastrum Meyen, Sorastrum Kuetzing and Hydrodictyon Roth. J. Indian bot. soc. 56:172-178.
- Pflug, H., 1953. Zur morphologie der sporomorphae (in pollen and sporen des mittaleuropaischen Tertiars by Thomson, P.W. & Pflug, H) Paleontographica 94B:16-18.
- Phadtrae, N.R. & Kulakarni A.R. 1984b. Palynological assemblage of lignite exposure of Ratnagiri district pp 515-531 in Badvae R.M. et al Proc. 10th Indian colloq.micropal.strat, Pune 1982, Maharashtra Association for the Cultivation of Science, Pune.
- Philipose M.T., 1967. Chlorococcales, Indian council agri. Res.New Delhi.
- Pierce, R.L., 1961. Lower Upper Cretaceous plant microfossils from Minnesota, Bull, geol. surv. minn, Univ. 42:1-86.
- Pocock, ., 1962. Microfloral analysis and age determination of strata at the Jurassic-cretaceous boundary in the western Canada plains, plaeontographica vol.111(B) ; 1-95.
- Potonie, R. 1931. Zur mikroskopie der Braunkohlen I-Z Braunkohle, 30:554-556 Halle.
- Potonie, R. 1934. Zur microbotanik des oecanen Humodils des Geiseltals (in zur Mikrobotanik der Kohlen und ihrer verwandten) Preuss Geol, land 4:25-125.
- Potonie, R. 1951. Die bedeutung der sporomorphen furdie Gesellscha ptschichete cr.s.congr. strat.geol.carbon.501-506.
- Potonie, R. 1956. Synopsies der Gattungen der sporae dispersae pt.I.Beihs geol.Jb.31!1.189.
- Potonie, R. 1958. Synopsis der Gattungen der sporae dispersae pt.ll Beihs, geol.Jb. 31.1-144.
- Potonie, R. 1960. Synopsies der Gattungen der sporae dispersae pt.lll Beihs, geol, Jb 39:1-189.

- Potonie, R. 1961. Synopsis der Gattungen der sporae disperse III. Beih, Geol.Jb; Vol.39:1-189.
- Potonie R. 1966 synopsis der Gattungen der sporae dispersae Pt.IV ibid, Vol.72 : 1-244.
- Potonie, r. 1970. Synopsis der Gattungen der sporae dispersae Pt.V Beih geol.Jb.87:1-222.
- Potonie, R. & Gelletich, J. 1933. Ueber pteridophyten-sporen einer eocanen Braunkohle aus Dorog in Ungarn Sber.Ges.Natural freunde Berl : 517-526.
- Potonie R. & Klaus, W. 1954. Einige sporengattungen des alpinen salzgebings Geol.Jb., 68:517-526.
- Potonie, R. & Kremp, G. 1954. Die Gattungender paleozoizchen sporae dispersae and stratigraphic Geol. Jb-69:111-194.
- Potonie, R. and Kremp, G. 1955. Die sporae disperse des Rutira Karbons. Pt.1, Palaeontographica Vol.98(B):1-136.
- Potonie, R., Thompson, P.W. and Thiengart F., 1950. Zur Nomenklatur and klassification due neogenen sporomorphae (pollen and spores) Geol. Jb.Vol.65:29-40.
- Potonie, R. & Sah, S.C.D. 1958. Spores disperse of the lignites from Cannanove beach of the Malbar coast of India, Paleobotanist, 7(2):121-135.
- Potonie, R. & Venitz, H. 1934. Zur Mikrobotanik des miozanen Humodiv/Sder nieder rheinischem Bucht (in Zur Mikrobotanik der kohlen and iherer verwandten, Preuss Geol. land 5:1-54.
- Prasad, B. & Dey, A.K. 1986. The occurrence of Eocene sediments in Arunachal Pradesh : a palynological evidence Bull. Oil & Natural Gas Commission 23(2):67-74.
- Raha, P.K., Rajendran, C.P. and Kar, R.K., 1986a. Occurrence of eocene palynofossils in subsurface Tertiary sediments of Kerala, J.Geol.Soc.India 28:48-50.
- Raha, P.K., Rajendra, C.P. and Kar, R.K. 1986b. Eocene palynofossils from the subcrop of Kerala, Bull.geol.min.metall,soc.India,54:27-232.
- Raha, P.K., Rajendra, C.P. and Kar R.K. 1987. Record of early Tertiary deposit in Kerala, India and its paleogeographic significance. Geophytology 17(2) : 209-218.
- Ramanujam, C.G.K. 1966. Palynology of the Miocene lignite from south Arcot, District Madras, India, Pollen spores, Vol.8,no.1, : 149-203.

- Ramanujam, .G.K. 1967. Myrtaceous pollen from the peaty lignite of Alleppey in Kerala State India. *Sci.Cull.* 33:229-230.
- Ramanujam, C.G.K. 1972. Revision of pteridophytic spores from the Warkalli lignite of south India. *Proc.Sem.Palaeopalynol.Indian.Stratig. Calcutta:* 248-254.
- Ramanujam, C.G.K. & Rao, K.P. 1977. A palynological approach to the study of Warkali deposits of Kerala in South India, *Geophytology*, 7(2): 160-164.
- Ramanujam, C.G.K. and Rao, K.P. 1978. Fungal spores from the Neogene strata of Kerala in South India:29-304 in Bhardwaj D.C. et al (Editors) *Proc.4th int.Palynol.conf.Lucknow 1976-77.* BSIP, Lucknow.
- Ramanujam, C.G.K. and Srisailam K. 1980. Fossil fungal spores from the Neogene beds around canmore in Kerala State, *Batanique* 9(1-4): 119-138.
- Ramanujam, C.G.K. 1987. Palynology of the Neogene Workalli beds of Kerala State in South India in Cenozoic stratigraphy and palynology in India, *Paleont, Soc.India. Spl.Pub.* 1:57-64.
- Rao, K.L.N., 1969. Lithostratigraphy of Paleogene succession of southern Cambay Basin: *Bull.ONGC Vol.6, No.1.*
- Rao, K.P. & Ramanujam, C.G.K. 1975. A palynological approach to the study of Quilon beds Kerala state in South India *curr.sci.* 44:730-732.
- Rao, K.P. & Ramanujam, C.G.K. 1978. Palynology of the Neogene Quilon Beds of Kerala state in South India. Spores of pteridophytes and pollen of monocotyledous. *Paleobotanist* 25:397-427.
- Rao, K.P. & Ramanujam, C.G.K. 1982. Palynology of Quilon Beds of Kerala State in south India-II pollen of dicotyledous and discussion paleobotanist 30(1):68-100.
- Rao, M.R., Saxena, R.K. and Singh, H.P. 1985. Palynology of the Barail(Oligocene) and Surma(Lr.Miocene) sediments exposed along Sonapur-Badarpur Road section, Jaintia Hills (Meghalaya) and Cachar (Assam) Pt.V, Angiospermous pollen grains. *Geophytology* 15(1):7-23.
- Rao, M.R. & Singh, H.P. 1987. Palynology of the Barail (oligocene) and Sumar (Lower Miocene) sediments exposed along Sonapur-Badarpur Road section, Jaintia Hills (Meghalaya) and Cachar (Assam) Pt.III, Pteridophytic spores paleobotanist 35(3):267-280.
- Rawat, M.S., Mukherjee, J.S. and Venkatachala, B.S. 1977. Palynology of Kadi Foramtion, Cambay Basin, India, *Proc.4 colloq.Indian Micropalaeontoal.& strat.* 179-192.

- Raymond, A, Christopher, David, C. Drowell, Jeorgen Reinhardt and Nalaine W. Markewish, 1980. The stratigraphic and structural significance of Paleocene pollen from warm springs, Georgia palynology` (AASP) vol.4:105-124.
- Rossignol, M. 1962. Analyse pdlinique de sediments marins Quarternaries en Israel, 2. sediments Pleistocenes Pollen spores 4(1):121-148.
- Rossignol, M.1964. Hystrichospheres du quaternaire en mediterrenee orientale dans les sediments pliocenes et les boues marines acutuelles, Rev. Micropaleont, 7: 83-99.
- Rouse, G.E. 1962. Plant microfossils from the Burrard Formation of Western British Columbia micropalaeontology, 8:187-218.
- Sah, S.C.D. 1967. Palynology of an Upper Neogene profile from Rusizi valley (Burundi) Annie Mus-R Fr.cent ser, 8, 57:1-173.
- Sah, S.C.D. & Dutta, S.K. 1966. Palynostratigraphy of the sedimentary formations of Assam-1. Stratigraphical position of the Cherra Foramtion. Palaeobotanist. 15:72-86.
- Sah, S.C.D. and Dutta, S.K. 1967. Palynostratigraphy of Tertiary sedimentary formations of Assam-2, stratigraphic significance of spores and pollen in the Tertiary succession of Assam Palaeobotanist 16(2):177-195.
- Sah, S.C.D. & Dutta, S.K. 1968. Paolynostratigraphy of the Tertiary sedimentary formation of Assam-2, srtatigraphic significance of spores and pollen in the tertiary succession of Assam Paleobotanist 16(1):42-47
- Sah, S.C.D. & Dutta S.K., 1969. Palynostratigraphy of the sedimentary formations of Assam-1. Stratigraphical positin of Cheera Formation. Palaeobotanist, Vol.15, No.2:72-86.
- Sah, S.C.D. & Kar R.K. 1969. Pteriodophytic spores from the Laki series of Kutch, Gujarat, India. J.Sen.Mem.Vol:109-121.
- Sah, S.C.D. & R.K.Kar 1970. Palynology of the Laki sediments in Kutch, 3. Pollen from the borehole around Jhulrai, Baranda and Panandhro, Palaeobotanist. 18(2):127-142.
- Sah, S.C.D. & Dutta, S.K. 1974. Palynostratigraphy of the sedimentary formation of Assam:3. Biostratigraphic zonation of the Cherra foramtion . . . of South Shillong Plateau, Palaeobotanist; 21(1):42-47.
- Sah, S.C.D. & Kar R.K. 1972. Palynostratigraphic evaluation of the Lower Eocene sediments of India:pp.255-265 in Ghosh A.K. et al. (Editors) -Proceedings of the Seminar on Paleopolynology and Indian Stratigraphy Calcutta 1971. Bot.Dept.Calcutta Univ.

- Sah, S.C.D. & Kar R.K. 1974. Palynology of the Tertiary sediments of Palana, Rajasthan. *Palaeobotanist* 21(2):163-188.
- Sah, S.C.D. & Singh, R.Y. 1974. Palynological biostratigraphy of the Tura Formation. *Stratigraphical Palynology Volume B.S.I.P. special volume No.3.*
- Sah, S.C.D. & Singh R.Y. 1977a. Status of palynology in the Tertiary stratigraphy of Assam and Gujarat:pp. 134-143 in Venkatachala B.S. & Sastri V.V. (Editors)-*Proc. 4th Colloq. Indian Micropal. Strat.* Dehradun 1974-75, Institute of Petroleum Exploration, Oil Nat.Gas Commission, Dehradun.
- Sah, S.C.D., Singh R.Y. & Singh H.P. 1980. Palynological zonation of the Tipam Group in Nahorkatiya area of Upper Assam:pp.635-642 in Bharadwaj D.C. et al (Editors)-*Proc.4th Int.palynol.Conf.Lucknow 1976-77* 2, Birbal Sahni Institute of Palaeobotany, Lucknow.
- Sahni, B., R.V.Sithole & G.S.Puri, 1947. Palaeobotany in India-6. Correlation of the tertiary succession in Assam by means of microfossils. *J.Ind. Bot.Soc.* Vol.26, No.4: 261-263.
- Salard-Cheboldaeff, M.1975. Grains de pollen Tertiaires du Cameroun rapportes a la marinene sedimenten Nouse Jahrbuch fir geologic and palantologic Abhurdlingen 107 : 278-340.
- Salard Cheboldaeff M. 1977. Paleopalynologie du bassin sedimentative littoral du cameroun dans ses rapports avec la stratigraphic et al paleoecologie These univ. P.M. Curi, Paris 262 : 29
- Salujha, S.K. & Kindra, G.S. 1981. Palynological fossils from the Langpar formation exposed along South Shilong front, Meghalaya. *India-Geosci. J.2(1):43-62*
- Salujha, S.K. & Kindra G.S. 1984. Palynostratigraphy of Atharmura Anticline, Tripura:pp. 391-409 in Badve R.M. et al (Editors)-*Proc. 10th Indian Colloq. Micropol.strat.*Pune 1982, Maharashtra Association for the Cultivation of Science, Pune.
- Salujha, S.K. & Kindra G.S. 1986. Palynostratigraphy of the Silchar-Haflong Road Traverse, District Cachar.*Bull.geol.Min.metall.Soc.India* 54: 238-249.
- Salujha, S.,K., Kindra G.S. & Rehman K. 1972a. Eximispora, a new genus from the Tertiary sediments of Assam. *J.geol.Soc.India* 13(3): 280-282.
- Salujha S.K., Kindra G.S. & Rehman K. 1972b. Palynology of the South Shillong Front-Part I: The Palaeogene of Garo Hills:pp.265-291 in Ghosh A.K. et al. (Editors)-*Proceedings of the Symposium on Poleopollinology and Indian Stratigraphy* Calcutta 1971, Bot.Dept.Calcutta Univ.

- Salujha, S.K., Kindra, G.S. & Rehman, K. 1977. Palynostratigraphy of the Tertiary sediments of the Gojalia anticline, Tripura, pt.1: Proc.IV Colloq. Micropalaeont.Stratigr. : 122-126.
- Salujha, S.K., Kindra G.S. & Rehman K 1979. Palynostratigraphy of Tertiary sediments of the Gojalia Anticline, Tripura Part 2. Systematic Palynology. J. Palynol, 14(1), 71-93.
- Salujha, S.K., Kindra G.S. & Rehman K. 1980. Palynostratigraphy of the Tertiary sediments of the Tulamura Anticline, Tripura:pp. 667-685 in Bharadwaj D.C. et al. (Editors)-Proc. 4th int.palynol.Conf.Lucknow 1976-77. 2, Birbal Sahni Institute of Palaeobotany, Luckno.
- Salujha & Rehman, K. 1972. Palynology of the South Shillong front, part-1: The Palaeocene of Garo Hills, Proc.Sem.Paleopalynol.Ind.Stratig. 1971, Calcutta, :265-291.
- Salujha, S.K., Rehman K. and Kindra, G.S. 1973. Distinction between the Bhuban and Bokabil sediments on the southern edge of Shillong Plateau based on palynofossil assemblages. Bull.ONC Vol.10, :109-117.
- Salujha & Rehman, 1974. Palynology of the South Shillong front Part-II: The Palaeogene of Khasi and Jaintia Hills-Palaeobotanist. 21:267-284; 1972.
- Sharma, D.C. 1991. Environmental reconstruction of subsurface Paleogene sequence in Broach depression, south Cambay basin, Gujarat in Proc. conference on Integrated, Exploration Research, Achievements and perspectives, Pandey, J and Banerjee (Eds), KDMIPE, Dehradun.
- Sharma K.D. 1988. Late Cretaceous-Palaeocene palynological assemblages from the subsurface of Godavary Krishna Basin:pp.65-67 in Maheshwari H.K.(Editor)-Palaeocene of India: Proceddings of the Symposium on Palaeocene of India:limits and subdivision Lucknow 1986, Indian Assocaiton of Palynostratigraphers, Lucknow.
- Sharma, P.S. & Ramanujam C.G.K. 1988. Pteridophytic sporomorphs from the second mine of ther Neyveli lignite deposit in Tamil Nadu, J.Swamy bot.Club 5(3-4):143-149.
- Sharma, P.S., Reddy P.R. & Kalavathi K. 1984. Pollen grains referable to monocotyledons from Neyveli lignite, Tamil Nadu, Indian, J.Bot.7:201-209.
- Sarjeant, W.A.S. 1970. Acritarches and Tasmanitids from the Chhidru Formation. Upper most permian of West Pakistan, in Stratigraphic boundary problems, Permian and Triassic of West Pakistan Univ.Kansas,Dept. of Geol.Spec, publs. : 277-304.

- Sarjeant, W.A.S. & Downie, C. 1974. The classification of dinoflagellate cysts above generic level : a discussion and revisions. Bibl Sahni Inst.Palaeobot Spec. Publ. 3: 9-32.
- Satish, K.Srivastava and Pier L. Binda 1991. Depositional history of Early Eocene Shumaysi Formation Saudi Arabia Palynology (AASP) vol.15 : 47-62.
- Saxena, R.K. 1978. Palynology of the Matanomadh Formation in type area north-western Kutch, India (Part 2). Systematic description of pteridophytic spores. Palaeobotanist, 25 : 448-456.
- Saxena, R.K. 1979. Reworked Cretaceous spores and pollen grains from the Matanomadh Foramtion (Palaeocene) Kutch, India, Palaeobotanist 26(2) : 167-174.
- Saxena, R.K. 1979. Palynology of the Matanomadh Formation in type area, Northwestern Kutch, India (Part-2). Systematic description of Gynospermous and Angiospermous pollen grains. Palaeobotanist, 26:130-143.
- Saxena, 1980. : Palynology of the Matanomadh Formation in typearea, North-western Kutch, India (Part-3) - Discussion - Palaeobotanist,26:279-296
- Saxena, 1982. Taxonomic study of the polycolpate pollen grains from the Indian Tertiary sediments with special reference to nomenclature. Rev.Palaeobot.Palynol.37:283-315.
- Selling, O.M. 1944. Studies in the recent and fossil species of Schizea, with particular reference to their spore characters. Meddn, Gotborg. bot.trad.16:1-112.
- Sen Gupta, B.K. 1964. Tertiary biostratigraphy of a part of north-western Kutch, Jl.Geol.Soc.India. 5:138-158.
- Sein, M.K. and S.C.D., Sah 1974. Palynological demarcation of the Eocene-Oligocene sediments in the Jowai-Badarpur Road Section, Assam. Sym. Strat.Palynol. :99-105.
- Shanmukhappa, M. 1990. Palynostratigraphy and paleoenvironmental analysis of Gandhar area in Cambay basin. ONGC Bull. Integrated exploration Research - achievements and perspective : 299-320.
- Shelf MVK Dilcher, D.L. 1971. Morphology and taxonomy of fungal spores, Palaeontographica B 133; 34-51.
- Siddhanta, B.K. 1986. The age of Neyveli lignite with reference to stratigraphy and palynology Indian minerals 40: 61-82.
- Singh H.P. 1982, Tertiary palynology of the Himalaya a review, Paleobotanist 28-29: 389-401.

- Singh H.P., Khanna A.K. & Sah, S.C.D. 1973. Problems and prospects of Tertiary palynology in northern India. Bull. Indian geologists Assoc. 6(1): 71-77.
- Singh, H.P., Singh R.Y. & Sah S.C.D. 1976. Palynological correlation of the Palaeocene subsurface assemblages from Garo Hills, Meghalaya. J.Palynol. 11:43-64.
- Singh, H.P., Khanna A.K. & Sah S.C.D. 1979. Palynological zonation of the Subathu Formation in the Kalka-Simla area of Himachal Pradesh, Himalayan Geol. 8(1): 33-46.
- Singh H.P. & Sarkar S. 1987a. Palynostratigraphy of Subathu Formation (Eocene in the Banethi-Bagthan area, Sirmaur District, Himachal Pradesh, Palaeobotanist 35(2):204-210.
- Singh, H.P. & Sarkar S. 1987b. Significant palynozones of Subathu Formation (Himachal Pradesh) and their bearing on stratigraphy, Palaeobotanist 35(3) : ;249-257.
- Singh H.P. & Saxena R.K. 1984. Palynology of the Neogene sediments of Jorajan Well-3, Upper Assam:pp. 613-631 in Sharma A,K. et al., (Editors)- Proceedings of the Symposium on Evolutionary and Biostratigraphy Calcutta 1979. A.K.Ghosh Commemoration Volume, Current Trends in Life Sciences 10.
- Singh, H.P. & Tripathi S,K.M. 1983. A comparative study of spores of ceratopteris Thalicoide (L) Brongu and Striatoleites Vander Hammen emend Kar Geophytology '13(2) : 219-226.
- Singh H.P. & Tripathi S,K.M. 1987. Palynology of the Jaintia Group (Palaeocene -Eocene) exposed along Jowai-Sonapur Road, Meghalaya India (Part -II), Data analysis and interpretation, Palaeobotanist 35(3):301-131.
- Singh, H.P., Singh R.Y. & Sah, S.C.D. 1975. Palynostratigraphic correlation of the Palaeocene subsurface assemblages from Garo Hills, Meghalaya -J.Palynology. 11: 43-64.
- Singh R.Y. 1975. Morphological study of the Retialetes complex from Indian tertiaries geophytology 5(1): 98-104.
- Singh, R.Y. 1977a. Stratigraphy and palynology of the Tura Formation in the type area-Part II (Descriptive palynology). Palaeobotanist 23(3) : 189-205.
- Singh, R.Y. 1977b. Stratigraphy and palynology of the Tura Foramtion in type area (Part III): Discussion. Palaeobotanist 24(1):1-12.
- Singh, R.Y. & Dogra N.N. 1988. Palynological zonation of Palaeocene in India with special reference to western Rajasthan. 51-64 in Maheshwari H.K.(Ed) Paleocene in India : Proceeding of the symposium on Paleocene of India : Limits and sub divisions Lucknow 1986, Indian Association of Palynostratigraphers, Lucknow.

- Srivastava, N.C. 1967. A new microfossil genus Oudhkusumites from Tertiary of India. *Rev. de Micropaleontologie* 10(1) : 37-41.
- Srisailam K & Ramanujam, C.G.K. 1982. Fossil pollen grains from the Warkalli sediments of Cannanore area. *Rec. Geol. Surv. India* 114(5): 123-131.
- Staplin F.L. 1960. Upper Mississippian plant spores from the Golata Formation Alberta, Canada. *Palaentolographica* 107B : 1-40.
- Stover, L.E. 1977. Oligocene and Early Miocene Dinoflagellates from Atlantic core hole 5/5B, black-plateau, Am, Assoc. stratgr, Palynol, contr. ser. 5(1): 66-89.
- Stover, L.E. & Evitt, W.R. 1978. Analysis of pre-Pleistocene organic-walled dinoflagellate cysts. stanf.univ.publ.geol.sci.stanford USA 15:300.
- Sudhakar, R. & Basu, D.N. 1973. A Reappraisal of the paleogene stratigraphy of southern Cambay basin. *Bull. of ONGC*, Vol.10, Nos.1-2: 55-76.
- Takahashi, K. 1964. Sporen und pollen der oberkertazeischen Hakobouchi schiehtengruppe Hokkaido, Mem, Fac, sei, kyushu univ. ser, D.Geol.14(3): 159-271.
- Thanikaimoni, G., Caratini, C., Venkatachala, B.S., Ramanujam, C.G.K and Kar, R.K. 1984. Selected Tertiary angiosperm pollens from India and their relationship with African Tertiary pollens. *Institute Francis de Podicherry, Trav aux de la section scientifique et Technique XIX* : 1-93
- Thiergart, F. 1938. Die pollen flora der Niederlausitzer Braunkohle, Jb Preuss, geol. 58: 282-351.
- Theirgart, F. 1940. Mikropalantologic als pollen analysis is Dienst der Braunkohlen forschung. Brenn, geol. 13: 1-82.
- Thiergard, F.X Frantoz, U 1962. Some spores and pollen grains from the Tertiary brown coal of Neyveli. *Palaeobotanist*, 11 : 43-45.
- Thomoson, P.W. & Pflug, H. 1953. Pollen and sporen des mittete-Uropaischen tertials, palaeontographics, 94: 1-138.
- Tripathi, S.K.M 1989. Algal and fungal remains from Jowai Sonapur Road section (Paleocene-Eocene) Meghalaya. *The Palaeobotanist* 37(1):63-76.
- Tripathi, S.K.M. & Singh H.P. 1985. Palynology of the Jaintia Group (Paleocene -Eocene) exposed along Jowri-Sonapur Road, Menghalaya India Part-I systematic palynology *Geophytology* 15(2):164-187.
- Trivedi G.K. 1985. Palynology of the Kopili Formation (Upper Eocene) exposed along Jowai Badarpur Road, Meghalaya J.Indian bot.sco.64:66-72.
- Valmiki, D., Kothe, J.M.Rao, W.M.Naidu, B.D. 1990. Exploration strategy of Cambay Basin (During VIII five year plan, Unpubl. ONGC).

- Van Der Hammen, . 1954. El desarrollo de la flora colombiana en los periodos Geológicos. I. Maestrichtiano Hasta Terciario mas Inferior, Boln. Beol.Bogota, 11(1):49-106.
- Van der Hammen, T. 1956. A palynological systematic nomenclature. Bull.Geol. 4(2-3):63-101.
- Van der Hammen T. 1956. Description of some genera and species of fossil pollen and spores. Bull.Geol.4(2-3): 111-117.
- Van der Hammen, T. & Wijmstra, T.H. 1964. A palynological study on the Tertiary and Upper Cretaceous of British Guiana. Leidse Geol.Med., 30:183-241.
- Van der Hammen, T. and Gracia de Mutis, C. 1965. The Paleocene pollen flora of Colombia. Leidse geol.Med.,Vol.35,pp.105-116.
- Van Hoeken-Klinkenberg, P.M.J. 1966. Maestrichtian, Palaeocene and Eocene pollen and spores from Nigeria. Leidse geol.vol.33, pp.37-48.
- Varma, C.P. and Rawat, M.S. 1963. A note on some diporate grains recovered from Tertiary horizons of India and their potential marker value Grana Palynol.Vol.4, pp.130-139.
- Varma Y.N.R., Ramanujam, C.G.K. & Patil R.S. 1986. Palynoflora of Tertiary sediments of Tonakkal area. Kerala J.Polynol.22(1):39-53.
- Varma, Y.N.R. 1987. Palynomorpha of ecological significance from Tonakkal Tertiary clays of Kerala State Indian J.Bot.10(2):163-168.
- Venkatachala, B.S. 1969. Palynology of the Mesozoic sediments of Kutch- 4 spores and pollen from the Bhuj exposures near Bhuj, Gujarat state Paleobotanist 17(2): 208-219.
- Venkatachala B.S. 1972. Observation on some palynological contributions to Indian stratigraphy, Palaeobotanist, 19(3), 284-296.
- Venkatachala, B.S. 1973. Palynological evidence of the age of Caddalore sandstone. Geophytology 3(2):145-140.
- Venkatachala B.S. 1974. Palynological zonation of the Mesozoic and Tertiary subsurface sediments in the Cauvery Basin:pp. 476-495 in Surange K.R. et al (Editors)-Aspects and Appraisal of Indian Palaeobotany. Birbal Sahni Instt. of Palaeobotany, Lucknow.
- Venkatachala B.S., Caratinie. Tissot C. and Kar D.K.1988. Paleocene-Eocene marker pollen from India and Tropical Africa. The Paleobotonist 37(1):1-25.
- Venkatachala, B.S. & Kar R.K. 1968. Fossil pollen comparable to pollen of Barringtonia from the Laki sediments of Kutch pollen spores 10(2), 335-339.

- Venktachala, B.S. & Kar, R.K. 1969a. Palynology of the Tertiary sediments of Kutch-1. Spores and pollen from bore hole no.14. Paleobotanist, 17(2):157-178.
- Venkatachala, B.S. & Rawat, M.S. 1971. Palynology of Tertiary sediments in Cauvery basin-1, Palaeocene-Early Eocene palynoflora from the subsurface, Seminar on Palaopalynology and Indian Stratigraphy, Calcutta, 1971.
- Venkatachala, B.S. & Rawat, M.S. 1972. Palynology of the Tertiary sediments in the Cauvery Basin. I. Palaeocene-Eocene Palynoflora from the subsurface. Proc.Sem.Palaopalynol.Indian stratigr.Calcutta:292-374.
- Venkatachala B.S. & Rawat M.S. 1973a. Palynology of the Tertiary sediments in the Cauvery Basin-2, Oligocene-Miocene palynoflora from the subsurface. Poloeobotanist 20(2): 238-263.
- Venkatachala, B.S. & Rawat M.S. 1973b. Palynological zonation of the Tertiary subcrop sequence in the Cauvery Basin 73-78. Polynology of the Cenophytic. Proc. 3rd int. palynol. Conf.Novosibirsk 1971. Acad.Sci.USSR Moscow.
- Venkatachala, B.S. & Sharma K.D. 1974. Palynology of the Cretaceous sediments from the subsurface of Pondicherry area, Cauvery Basin. New Bot, 1 (3&4):170-200.
- Venkatachala, B.S. & Sharma, K.D. 1984. Palynological zonation in subsurface sediments in Well.no.1, Godavari-Krishna Basin India, pp 445-465 in Badre et al. proc, 10th Colloqy. Indian micropal, strat. Pune 1982 Maharastra association for the cultivation of Sci. Pune.
- Wall, D.1967. Fossil microplankton in deep sea cores from the Caribbean sea; Palaeotology V.10; 95-123.
- Wilson G.I. 1967a. Some new species of lower Tertiary dinoflagellates from Mc Murdo sound, Antarctica: New Zealand Jour.Botany,V.5:57-83.
- Wilson, L.R. & Webster, R.M. 1946. Plantmicrofossils from a Fort union coal of Montana, Am.J.Bot, 33:271-278.
- Williams, G.L. & Downie C. 1966a. The genus *hystrichokolpoma* Bull.B.C.Mus. (Not Hist) geol.Suppl.London 3:176-181.
- Williams G.L. & Downie C 1966b. The wetzeliella from the London clay, in Davey et al 1966 Bull.Br.Mus (Nat Hist) Geol.suppl .London:182-198.

- Williams G.L. & Downie, C. 1965c. Further dinoflagellate cysts from the London clay in Davey et al, 1966 a Bull Br.Mus. (Nat Hist) Geol. Suppl . London 215-235.
- Wodehouse, P. 1933. Tertiary pollen. The oil shales of the Eocene green river formation Bull.Torrey bot. club. 60: 479-524.
- Zubov, I.P. Naugolny, I.K.Zapivalov, N.P. and Chandra P.K. 1966. Problem of correlation and distribution of hydrocarbon bearing horizons in the Eocene of Cambay Basin Bull.ONGC. Vol.3, No.2.
- Zamaloa, M.C. and Romero, E.J. 1990. Some spores and pollen from the Cullen Foramtion (Upper Eocene to Middle Oligocene) Tierra Del Fuego Argentina, AASP. Vol.(14): 123-132.