

List of Figures

Figure No.		Page No.
1.1	Location map of the study area. L-Lunavada, S-Santrampur, G-Godhra, V-Vadodara, M.R.- Mahi River, P.R.- Panam River.	3
1.2	Shuttle Radar Topography Mission (SRTM) derived shaded relief map of the area lying towards the southeast of the Lunavada town. Study area is located within the prominent quartzitic fold which has occupied the Southern part of the image. Source: USGS Earth-explorer.	5
2.1	Generalised geological map of Aravalli Mountain belt, (modified after Singh et al., 2010).	9
3.1	Geological map of the study area, modified after Mamtani (1998)	34
3.2	Satellite imagery (False Colour Composite) of the part of the area lying southeast to the Lunavada town (Toposheet no.46E/12). Rectangle with dashed lines indicates the study area.	36
3.3	Field photograph showing:	37
	a) Unoriented/star shaped amphibole needles within calc-silicate rock, Loc. Jotangiya;	
	b) Maculose structure over calc-silicate surface, Loc. Ora;	
	c) Compositional bandings, Loc. Amliyat;	
	d) Differential weathering of calc-silicates, Loc. Panam river section;	
	e) Development of S_1 and S_2 foliations within calc-silicate, Loc. Talwadi;	
	f) Two sets of joints within calc-silicates, Loc. Junakheda;	
	g) Structure like convolute bedding, Loc. Balaji No Muvada;	
	h) Pegmatite vein discordant to calc-silicate rocks, Loc. Ora;	
	i) Boudinage of quartzite within calc-silicate rock, Loc. Junakheda;	

- j) Folded quartz vein within calc-silicate, Loc. Chari;
- 3.4 Field photograph showing: 39
- a) Jointed quartzite, Loc. Panchmuva;
 - b) Massive, buff coloured quartzites, Loc. Tadi;
 - c) Plumose markings in quartzite, Loc. Hoselav;
 - d) Reddish brown surfaces of quartzite, Loc. Chandsar.
- 3.5 Field photograph showing: 40
- a) Well section having intercalations of metapelite and quartzite, Loc. Boriya;
 - b) Mahi river section showing intercalated sequence of chlorite schist and quartzite;
 - c) Outcrop of biotite schist, Loc. Junakheda ;
 - d) Garnet mica schist, Loc near Labhi village.
- 3.6 Field photograph showing: 42
- a) Panoramic view of granite country, Loc. Erandana Muvada;
 - b) Fine grained grey coloured granite having xenolith of schist, Loc. Guneli;
 - c) Granite- quartzite contact, Loc. Kothamba;
 - d) Coarse grained pink granite having xenoliths of schist, Loc. Palikhanda;
 - e) Pegmatite within granite with segregated tourmaline crystals, Loc. Hosselav;
 - f) Granite - Metapelite contact, Loc. Jhaidi.
- 4.1 Photomicrographs (in crossed polars) of calc-silicate samples showing: 50
- a) Hornfelsic texture developed by unoriented actinolites,
 - b) Rarely observed preferred orientation of actinolites,
 - c) Kink seen in biotite porphyroblast ,

- d) Star shaped actinolites,
- e) Actinolite porphyroblast with quartz inclusions having concordant relationship with matrix suggesting post tectonic origin of it,
- f) Replacement of ankeritic mass by salite,
- g) Actinolite having inclusions of salite suggesting its formation from latter,
- h) Titanite grains in association with ilmenite and quartz,
- i) Biotite grain nucleating along the margins of actinolite,
- j) Epidote grains in association with actinolites,
- k) Scapolite associated with other minerals like quartz, calcite and biotite,
- l) Formation of myrmekitic intergrowth.

4.2 Back Scattered Electron images of major calc-silicate minerals: 52

- a) Star shaped actinolites,
- b) Poikiloblastic actinolites surrounded by titanite, microcline etc.,
- c) Replacement of diopside by actinolite,
- d) Wedge shaped titanite grain,
- e) Biotite development along edges of actinolites,
- f) Epidote grains associated with actinolite porphyroblasts.

4.3 Photomicrographs (in crossed polars) showing: 56

- a) Serrated quartz grains boundaries within quartzite lying far from intrusion,
- b) Development of granoblastic polygonal texture can be seen within quartzite lying closer to the intrusion,
- c) S_1 foliation defined by biotite and muscovite to which bigger porphyroblast of biotite are obliquely aligned forming S_2 foliation plane,
- d) Inclusion trails within garnet porphyroblast defining internal schistosity (S_i) having oblique relation with external schistosity (S_e) indicating syntectonic growth of this porphyroblast,
- e) Fine grained grey granite with abundant plagioclase feldspar biotite and quartz,
- f) Coarse grained pink granite with abundant microcline and quartz.

5.1	Ca-Amphibole, classification scheme after Leake et al., (1997).	60
5.2	Ca-Mg-Fe triangular plot showing clinopyroxene composition.	62
5.3	Mica, classification scheme after Deer et al., (1992).	66
5.4	An-Ab-Or diagram showing plagioclase feldspar having composition range from (An ₂₂ -An ₆₁) .	69
5.5	Nomenclature and classification of Chlorite (after Hey,1954).	71
6.1	Whole-rock concentrations of major-element vs. silica in weight percent oxides for calc-silicates samples of Lunavada. Concentrations of all major elements excluding Al ₂ O ₃ , K ₂ O, Na ₂ O and TiO ₂ exhibit near-linear inverse correlations with SiO ₂ . *All iron is reported as FeO.	79
6.2	Chondrites normalized rare earth element (REE) patterns for calc-silicates of the study area (after Sun and McDonough, 1989).	80
6.3	The bulk-rock compositions of calc-silicates are plotted on a CaO-Al ₂ O ₃ -FeO+MgO (wt. %) ternary diagram. General bulk compositions of unaltered arkoses, pelites, greywackes and marls are represented by shaded fields along with schematic regions for metasomatic types (after Barton et al.,1991).	81
6.4	Whole-rock compositions of selected trace-elements along with Na ₂ O and K ₂ O exhibit a positive correlation with Al ₂ O ₃ in these calc-silicates (with the exception of Sc and Ni) suggesting these components were bound in clays.	83
6.5	Whole-rock compositions of calc-silicates fall within the continuum between sandstone and shale (grey) on a ternary Al-Zr-Ti diagram (after	84

Garcia et al., 1994). The field outlined by the dashed line represents the typical curved trend exhibited in whole-rock compositions of rocks derived from calc-alkaline plutonic suites.

6.6	A-CN-K diagram (after Nesbitt and Young, 1982) using molar proportions of Al ₂ O ₃ , CaO, Na ₂ O and K ₂ O. As per the CIA scale, sample plots of calc-silicates indicate low to moderate and weathering of source rocks.	85
6.7	Th/Sc vs Zr/Sc ratio diagram showing the composition of the sources for the primary sediments of calc-silicate rocks and the extent of recycling or concentration of Zr in sediments, associated with the abundance of heavy minerals, particularly zircon (after McLennan et al.,1990).	86
6.8	Th/Co vs. La/Sc diagram (after Cullers, 2002) where sample data plotted show affinity towards the source rocks of homogeneous composition of felsic nature.	87
6.9	Sc-Th-Zr/10 diagram of Bhatia and Crook (1986), A: oceanic island arc; B: continental island arc; C: active continental margin D: passive/rifted margins; calc-silicate samples fall in the 'C' region of this diagram indicating deposition of primary sediments in active continental margin setting.	88
7.1	Isobaric T-X(CO ₂) diagram for sample Ora-03 at 1 kbar. Shading represents degrees of variance, such that lighter shades are low variance and darker shades are higher variance	95
7.2	Isobaric T-X(CO ₂) diagram for sample Jtng-09 at 1 kbar	96
7.3	Isobaric T-X(CO ₂) diagram for sample Tlwd-14 at 1 kbar	98
7.4	Isobaric T-X(CO ₂) diagram for sample Bor-P10 at 1 kbar	99

7.5	The sequence of CMS-HC chemographic diagrams for metamorphosed protolith. a) Appearance of talc mineral by low grade metamorphism of protolith; b) Development of tremolite with progression of metamorphism; c) Appearance of diopside with further progression of metamorphism. All chemographic calculations were conducted in Perple_X v.6.8.1 (Connolly,2005;Connolly,2009) using thermodynamic and end-member compositional data from the Holland and Powell (2003) dataset .	100
7.6	The temperature- pressure diagram, (after Winter, 2010).The calc-silicate rocks fall within the ‘Hornblende-hornfels facies’ of contact metamorphism which is shown by the symbol ‘*’ in the respective field.	101
7.7	Cartoon diagrams showing the different stages of evolution of the Lunavada basin, NW India. a) Stage 1, (~1.83 Ga) : Rifting led to the opening of Aravalli basin, b) Stage 2, (1.77 Ga) : Passive margin stage with deposition of Jhamarkotra carbonates and phosphorites in a shallow water along with deposition of carbonaceous shales in comparatively deeper water, c) Stage 3, (1.64-1.59 Ga) : Eastward subduction of oceanic crust leading to the development of accretionary prism, d) Stage 4, (~1.59 Ga): Crustal extension giving rise to back arc basin (Lunavada basin), (modified after Bhowmik and Dasgupta, 2012 ; Absar and Sreenivas,2015).	103
8.1	Calc-silicate sample locations within the Champaner Group, SAMB, Gujarat (modified after Gupta et al.,1992).	106
8.2	Geological map of the study area within the Champaner Group, (after Joshi, 2019). Orange coloured stars represent Calc-silicate sample locations.	108
8.3	Field photograph showing: a) Unoriented amphibole needles within calc-silicate rock, Loc. Intwada;	109

	b) Elephant skin weathering, Loc.Chalvad;	
	c) Foliations,Loc. Vav;	
	d) Calc-silicates having greenish colour, Loc. Goldungari.	
8.4	Photomicrographs (in crossed polars) of calc-silicate samples showing:	111
	a) Formation of tremolite by replacement of dolomite;	
	b) Diopside nucleating along the borders of tremolite;	
	c) Scapolite associated with calcite and poikiloblastic garnet having quartz and calcite inclusions.	
8.5	Back Scattered Electron images of major and important calc-silicate minerals and reaction textures:	111
	a) Tremolites replaced by diopside,	
	b) Poikiloblastic garnet.	
8.6	Calcic-amphibole, classification scheme after Leake et al.,(1997). (Orange solid circles indicate samples from Champaner region and black solid circles represent samples from Lunavada region).	120
8.7	Ca-Mg-Fe triangular plot showing Clinopyroxene composition.	120
8.8	Nomenclature and classification of Chlorite (after Hey,1954).	121
8.9	Whole-rock concentrations of major-element versus silica in weight percent oxides for calc-silicates samples. (Black solid circles represent calc-silicate rocks from Champaner region while blue solid circles denote calc-silicates from Lunavada region.) *All iron is reported as FeO.	124
8.10	Chondrites normalized rare earth element (REE) patterns for calc-silicates of the study area within Champaner Group (after Sun and McDonough,1989).	125

8.11	The bulk-rock compositions of calc-silicates are plotted on a CaO-Al ₂ O ₃ -FeO+MgO (wt. %) ternary diagram (after Barton et al.,1991).	125
8.12	Whole-rock compositions of selected trace-elements along with Na ₂ O and K ₂ O exhibit a positive correlation with Al ₂ O ₃ in these calc-silicates suggesting these components were bound in clays.	126
8.13	Whole-rock compositions of calc-silicates fall within the continuum between sandstone and shale (grey) on a ternary Al-Zr-Ti diagram (after Garcia et al.,1994).	127
8.14	CaO-MgO-SiO ₂ ternary diagram classification system for marbles, (Storey and Vos,1981).	127
8.15	Th/Sc vs Zr/Sc ratio diagram showing the composition of the sources for the primary sediments of calc-silicate rocks and the extent of recycling or concentration of Zr in sediments, associated with the abundance of heavy minerals, particularly zircon (after McLennan et al., 1990).	128
8.16	Th/Co vs. La/Sc diagram (after Cullers, 2002) where samples can be seen to be showing affinity for a source rock of homogeneous composition.	129
8.17	Sc-Th-Zr/10 diagram of Bhatia and Crook (1986). A: oceanic island arc; B: continental island arc; C: active continental margin D: passive/rifted margins.	129
8.18	Isobaric T-X(CO ₂) diagram for sample Gndh-06 at 1 kbar.	131