

## **LIST OF TABLES**

**Table 1:** Products containing *H. indicus* as an ingredient

**Table 2:** Murashige and Skoog's (MS, 1962) basal stock composition

**Table 3:** Preparation of PGR stock solutions

**Table 4:** Reaction mixture and PCR conditions for cDNA synthesis

**Table 5:** Reaction mixture and PCR conditions for  $\beta$ -actin amplification

**Table 6:** Reaction mixture and PCR conditions for gene amplification

**Table 7:** Effect of cytokinins (BA/Kn), auxins (IAA/NAA) and combinations of cytokinins (BA and Kn) on shoot regeneration from nodal explant of *H. indicus* (8 weeks)

**Table 8:** Effect of different combinations of cytokinins with auxins on shoot regeneration from nodal explant of *H. indicus* (8 weeks)

**Table 9:** Effect of cytokinins on shoot regeneration from *in vitro* nodes of *H. indicus* (8 weeks)

**Table 10:** Effect of different strengths of MS medium fortified with IBA on root induction from *H. indicus* shoots (4 weeks)

**Table 11:** Effect of different strengths of MS medium fortified with NAA on root induction from *H. indicus* shoots (4 weeks)

**Table 12:** Fingerprint of different samples for hexane extracts

**Table 13:** Fingerprint of different samples for ethyl acetate extracts

**Table 14:** Fingerprint of different samples for methanol extracts

**Table 15:** Lupeol content in different samples of *H. indicus*

**Table 16:** Lupeol content in shoot cultures of *H. indicus* after elicitation using yeast extract

**Table 17:** Lupeol content in shoot cultures of *H. indicus* after elicitation using salicylic acid

**Table 18:** Primers used for gene expression analysis