

References

- Abdulazeez A.M., et al,(2010). Effect of *Peristrophe bicalyculata* on Blood Pressure, Kidney and LiverFunctions of Two Kidney One Clip (2K1C) Hypertensive Rats *British Journal of Pharmacology and Toxicology* **1(2)**: 101-107.
- Aboutabl EA (1996). *Phytochemical profile and bioactivity validation of traditional use of Aerva lanata and Aerva javanica*.*Phytomedicine*;3:p 131.
- Adinortey MB, Galyon IK, Asamoah NO. (2013).*Trema orientalis* Linn. Blume: A potential for prospecting for drugs for various uses. *Pharmacognosy Reviews* ;7(13):67-72.
- Afaq SH, Tajuddin and Afidi R (1991). BisehriBooti (*Aerva lanata* Juss.): Some lesser known uses and pharmacognosy. *Ethnobotany*;3:pp 37-40
- Aiyer VN, Narayanan V, Seshadri TR and Vydeeswaran S (1973).*Chemical components of some Indian medicinal plants*. *Indian J.Chem.*;11:pp 89-90.
- Alagammal M, Tresina Soris P and Mohan V R (2011) Chemical Investigations of *Polygala Chinensis* L. by GC-MS *Science Research Reporter* **1(2)**:49-52.
- Anonymous (1999)*Pharmacognosy of Indigenous Drugs*, V-II, CCRAS, Govt.of India, New Delhi. p.608
- Anonymous,(1990) *Phytochemical investigations of certain medicinal plants used in ayurveda*, publisher CCRAS, Govt.of India, New Delhi.p.299.
- Anonyms (1966) *The Wealth of India, Raw Materials*, Vol. VII, N-Pe, CSIR, New Delhi.

Annonyms (1956), *Glossary of Indian medicinal plants*, publisher CCRAS, Govt.of India, New Delhi. pp. 199.

Anonymous (2005). *Database On Medicinal Plants Used In Ayurveda. Vol-III*, CCRAS, New Delhi. p.77.

Anonymous (1956). *The Wealth of India: Raw Materials. Vol. IV*. New Delhi: Publication and Information Directorate, Council of Scientific and Industrial Research; p. 68.

Anonymous (1989). *TheAyurvedic Pharmacopoeia of India*. 1st Edn.NISCAIR ;1(1):p.120-121.

Anonymous (1990) *phytochemical investigations of certain medicinal plants used in ayurveda*, CCRAS, New Delhi. pp.190.

Anonymous (1990) *The Ayurvedic Pharmacopoeia of India*. Part-I **Vol - I-IV** 1st Edition. NISCAIR, New Delhi. pp. 127.

Anonymous (1990). *Phytochemical investigations of certain medicinal plants used in ayurveda*, CCRAS, New Delhi.p.169.

Anonymous (1990). The Ayurvedic Pharmacopoeia of India. Part-I **Vol- I**. 1st Edn. NISCAIR, New Delhi; p.56.

Anonymous (1990). The Ayurvedic Pharmacopoeia of India. Part-I **Vol- I**. 1st Edn. NISCAIR, New Delhi. pp. 17-18.

Anonymous (1996). *Indian Medicinal Plants, Vol. III*. Hyderabad (India): Orient Longman Ltd; p. 123.

Anonymous (1999) *The Ayurvedic Pharmacopoeia of India*. Part-I **Vol-II**,1st Edition.. NISCAIR, New Delhi. p. 46.

Anonymous (1999) *WHO Monographs on selected Medicinal Plants Materials Vol-I* WHO, Geneva: pp.183.

Anonymous (1999). *Pharmacognosy of Indigenous Drugs, Vol-I*, CCRAS, New Delhi; p.446.

Anonymous (1999). *Pharmacognosy of Indigenous Drugs. Vol-I*, CCRAS, New Delhi; pp.129-133.

Anonymous (1999). *The Wealth of India (Raw Materials)*. **Vol. IX**. CSIR, New Delhi.

Anonymous (2001), *Ayurvedic Pharmacopeia of India*, Part I, **Vol-III**, pp.76-75.

Anonymous (2001). The Ayurvedic Pharmacopoeia of India. Part-I **Vol- III**. 1st Edn. NISCAIR, New Delhi;pp. 183-184.

Anonymous (2004). *The Ayurvedic Pharmacopoeia of India. Part-I Volumes IV*. 1st Edn. NISCAIR, New Delhi.p. 138.

Anonymous (2004).Reviews on Indian Medicinal Plants,Indian Council of Medicinal Research, New Delhi,**Vol.IV** (Ba-By) ; pp.188-198.

Anonymous (2005) *Database On Medicinal Plants Used In Ayurveda Vol-3* CCRAS, New Delhi. pp.562.

Anonymous (2007). *Reviews on Indian Medicinal Plants Vol - 5 (Ca-Ce)* Medicinal Plants Unit, Indian Council Of Medical Research, New Delhi; p.941.

Anonymous(2004) *WHO Monographs on selected Medicinal Plants Materials Vol.II* WHO, Geneva p.276.

Anonymous. *Wealth of India- Vol-I*, CSIR, New Delhi.

Anonymous. *Wealth of India- Vol-III*, CSIR, New Delhi.

Anonymous.(1948) *Wealth of India Vol-I*, A-B, Raw material CSIR, New Delhi pp.3.

Arun Bhatt, Ayurvedic Herbal Industry: Quest for Global Acceptance
www.clinivet.com

Atta-ur-Rahman, Bhatti M. K, Akhtar F, Choudhary M. I (1992). Alkaloid of *Fumaria indica*. *Phytochemistry*; **31**; p2869.

Atta-ur-Rahman, Bhatti M. K, Choudhary M. I (1992). Chemical constituent o *Fumaria indica*.*Fitoterapia*; **13**: 129.

Atta-ur-Rahman, Bhatti M.K, Choudhary M I, Ahmad S.K (1995). Alkaloidal constituent of *Fumaria indica*. *Phytochemistry*; **40**: p. 593.

Audani Rashmi,*et al.*, (2009). Pharmacognostical and Phytochemical Investigation on Aerial Parts of *Peristrophe bicalyculata*. *PHCOG J*. **1(1)**:1-7.

Ayush <http://indianmedicine.nic.in>.

Balakrishnan N *et.al*,(2009)The Evaluation of Nitric Oxide Scavenging Activity of Acalypha Indica Linn Root *Asian J. Research Chem.* **2(2)**.

Balraj P. and Nagarajan S. (1981) Flavonol glycosides from three members of the Lythraceae.*J Nat Prod*; **44**:660-661.

Bensky D and Gamble A. (1990). *Chinese Herbal Medicine: Materia Medica*. Eastland Press, Seattle, WA.

Bhakuni D.S., Dhar M.L., Dhar M.M., Dhawan B.N., Gupta B. and Srimal R.C. (1971) Screening of Indian plants for biological activity. Part III. *Indian J Exp Biol*; **9** : 91-102.

Bhandari C.R. (1977). "Van aushadi chandrodaya"(Hindi), Chaukhamba Sanskrit sansthan, Varanasi.

Bhatnagar L.S., Singh V.K. and Pandey G. (1973) Medico-botanical studies on the flora of Ghatigaon forests, Gwalior. Madhya Pradesh. *J Res Ind Med*; **8**:67-100.

Broker, R and Bhat. J.V. (1953). Riboflavin and thiamine contents of gums. *Current Science* **22** :343.

Burkill H.M. (1985). The useful plants of West Tropical Africa. 2nd Edn. **Vol. I.** Families A-D. Royal Botanic Gardens Kew.

Chandra S and Sastry MS (1990). Chemical constituents of *Aerva lanata*. *Fitoterapia*; **61**: p 188.

Chang H.M and But P.P.H (1986). *Pharmacology and Applications of Chinese Materia Medica*. **Vol. II.** World Scientific, Singapore.

Chang Minyi. (1992). *Anticancer Medicinal Herbs*. Human Science and Technology Publishing House, Changsha.

Chang, (2002), Bioactive polysaccharides from traditional Chinese medicine herbs as anticancer adjuvants, *Jour. Alter. Complement. Ther.*, **8**: 561.

Charaka Samhita, Shree Gulab Kumverba Ayurvedic Society, Jamnagar,(1949), Sutra Sthana Adh. 4-15.

Chauhan SMS, Singh M, Narayan L. (2002). Isolation of 3 β -hydroxyolean-12-ene, friedelin and 7-methoxy-4-5-dihydroxyisoflavone from dry and fresh leaves of *Shorea robusta*. *Indian J Chem Sec B*; **41(5)**: 1097–1099.

Chauhan.J.S., sultan. M. and Srivastava, S. K.(1980) Constituents from *Salmalia malabaricum*. *Can J Chem* ;**58** :328-330.

Chen C.C, Hsin W.W, Ko F.N, Huang Y.L, Ou J.C, Teng C.M (1996). Antiplatelet arylnaphthalide lignans from *Justicia procumbens*. *J Nat Prod*; **59**: 1149-1150.

Chetty K.M., Chetty M.L., Sudhakar A. and Ramesh C. (1998) Ethno-medico-botany of some aquatic angiospermae in Chittoor district of Andhra Pradesh, India. *Fitoterapia*; **69**: 7-12.

Chiang T-C, Chang H-M, Mak TCW(1983) New oleanene-type triterpenes from *Abrus precatorius* and X-ray crystal structure of abrusgenic acid-methanol 1 :1 solvate. *Planta Med* ; **49**: 165-169.

Chopra R.N., Nayar S.L. and Chopra I.C. (1956) “ *Glossay of Indian Medicinal Plant*” CSIR New Delhi; p. 74.

Chopra, H. (1959) Medicinal Herbs of Chhattisgarh, India Having Less Known Traditional Uses. <http://www.botanical.com>.

Chowdhury Abu Asad and Islam M. S.(2004). Antimicrobial Activity of *Trema orientalis*. *Dhaka University Journal of Pharmaceutical Sciences*; **3** :1-2.

Chowdhury D, Sayeed A, Islam A, Bhuiyan MSA and Khan GRMAM (2002). Antimicrobial activity and cytotoxicity of *Aervalanata*. *Fitoterapia*; **73**: pp 92-94.

Daniel M (2006). *Medicinal Plants- Chemistry and Properties*. Oxford & IBH Publishing Co. Pvt. Ltd., New Delhi.

Das S.R. (1995)Some beneficial weeds of cultivation from Nadia district, West Bengal. *Bull Med Ethnobot Res*; **16**:74-81.

Datta S.C.and Mukerji B. (1950) *Pharmacognosy of Indian root and rhizome drugs* bulletin no. 1, Ministry of health, Publications, Delhi.pp.121.

De Eds, F. (1968). In: Comprehensive Biochemistry (M. Florkin and E. H. Stotz ed.) Elsevier publishing Co., Amsterdam.

De Smet PAGM (2002)Herbal Remedies N Engl J Med 347: 2046-56.

Desai V. G. (1975) "Aushadi Sangraha "(Marathi), Shri Gaganan Book Depot Prakashan, Mumbai.

Dey, A.C. *Indian medicinal plants used in ayurvedic preparations*, Bishen Singh Mahendra Pal Singh; DehraDun, 1980; pp. 202-203.

Dhar M.L., Dhar M.M., Dhawan B.N., Mehrotra, B.N., Srimal R.C. and Tandon J.S. (1973).Screening of Indian plants for biological activity. Part IV. *Indian J Exp Biol*; 11: 43-54.

Dhar. D.N. and Munial. R.C.(1976) Chemical examinatin of the seeds of the *Bombax malabaricum*. *Planta Medica* ;29:148-150.

Duke J.A. (1997).*The Green Pharmacy*. Rodale Press, PA, USA.

Duret S and Paris R.R. (1977). The flavonoids of several species of Bauhinia. *Plant. Med. Phytother* ; 11: 213-15.

Durga K. Ramya *et.al*, (2010) Isolation of potential antibacterial and antioxidant compounds from *Acalypha indica* and *Ocimum basilicum* , *African Journal of Plant Science*.4(5);163-166.

Dwivedi, S.(2002). Ethnomedicinal Uses of Some Plant Species by Ethnic and Rural Peoples of Indore District of Madhya Pradesh, India.<http://www.pharmainfo.net>.

Dymock W., Warden C.J.H & David Hooper (1976)" *Pharmacographia Indica*" M/s Periodical Experts,

Fang, Y.Z, et al., (2002) *Nutrition*, 872.

Frank JL. *Herbalism*.(1996). Duncan Baird Publishers, London.

Frankel, E. N. and Meyer A.S (2000). The problems of using one dimensional methods to evaluate multifunctional food and biological antioxidants, *J. Sci. Food Ageic*; **80**: 231.

Gamble JS.(1921)*Flora of The Presidency of Madras*, Vol. II. London: Published Under the

Geissman, T.A. (1962). Chemistry of Flavonoid Compounds, Pergamon Press, Oxford Syst. Bot. **3**: 353-368.

Ghazanfar SA.(1994). *Handbook of Arabian Medicinal Plant*.CRC Press, Florida, USA; p. 342.

Ghosal S, Dutta SK.(1971) Alkaloids of Abrus precatorius,*Phytochemistry*; **10**:195-198.

Giwa O.E. et al, (2010). Screening of Antimicrobial Ethanolic Extract of *Peristrophe bicalyculata* .*Ethnobotanical Leaflets* **14**: 766-73.

Gopal. H and Gupta. R.K.(1972). Chemical constituents of *Salmalia malabarica* Schott. and Endt flowers. *J Pharmaceut Sci* ;**61**: 807-808.

Gordon, M.C. and David J.N., (2001). Natural Product Drug Discovery in the next millennium, *Pharm Biol* ;**39**: 8-17.

Gupta A.K. , Tandon Neeraj ,Sharma Madhu (2008) *Quality standards of indian medicinal plants Vol-8*, Medicinal plants unit, indian council of medical research new delhi. pp.12.

Gupta A.K. and Chauhan J.S. (1984). Constituents from the stem of *Bauhinia variegata*. *Natl Acad Sci Lett* ; **7**: 174-76.

Gupta, A.K, Vidyapati T.J. and . Chauhan J.S (1980). Chemical examination of the stem of *Bauhinia variegata*. *Planta Med.* ;**38**: 174-76.

Gupta, A.K. Vidyapati T.J. and Chauhan J.S. (1979). 5, 7-dihydroxyflavonone-4 -O -L rhamnopyranosyl -D - glucopyranoside from the stem of *Bauhinia variegata*. *Ind. J. Chem* ;**18B**: 85-86.

GuptaA.K.,TandonNeeraj and SharmaMadhu (2008).*Quality Standards Of Indian Medicinal PlantsVol-* **3** Medicinal Plants Unit, Indian Council Of Medical Research, New Delhi.

Harbone JB. (1999).Recent advances in chemical ecology. *Nat Prod Rep*; **16(4)**: 509-523.

Harborne J.B. (1967 & 1984). *Phytochemical methods*,(1st and 2nd Edns.). Chapman and Hall, London.

Harborne J.B. and Smith D.M. (1978). Correlations between anthocyanin chemistry and pollination ecology of the Polemoniaceae. *Biochem. Syst. Ecol.* **6**: 127.

Harsoliya MS, Pathan JK, Khan N, Bhatt D, Patel UM. (2011). Effect of ethanolic extracts of *Bergenia ligulata*, *Nigella sativa* and combination on calcium oxalate urolithiasis in rats. *Int Drug Formulation Res*, **2(2)**:268-280.

Hase K., Kadota S., Basnet P (1997). *Planta Medica*; **63**: 216—219.

Hideaki O, Kayoko Y, Kazuo Y and Cantoria M.C. (1991).Isolation of 10-O-Acyl Iridoid Glycosides from the Philippine Medicinal plant, *Oldenlandia corymbosa L.* *Chem Pharm Bull.*; **39(8)**:2049-2052.

Hossain, T., Guha, R. C. and Mukerji B., (1943), *Science and Culture* ; **9**:167.

Hota RK, Bapuji M. (1993). Triterpenoids from the resin of *Shorea robusta*. *Phytochem*; **32(2)**: 466-468.

Hullatti K. K. and Bhattacharjee Prasenjit (2011). Pharmacognostical Evaluation of Different Parts of *Coleus amboinicus* lour., Lamiaceae. *Pharmacognosy Journal* ; **3(24)**:pp.39-45.

Husain W. and Siddiqui M.B. (1987). Ethno-botanical approach of North-western U.P. *Acta Bot Indica*; **15**: 94-97.

Idrahim, R.K. and G.H.M. Towers (1940). The identification by paper chromatography of plant phenolic acids. *Arch.Biochem. Biophys.* **87**:125.

Jain R. and Gupta, R.K. (1986a) Chemical investigation of *Alysicarpus bupleurifolius* DC. *J Indian Chem Soc* **63**:531-532.

Jain, R. and Gupta, R.K. (1985) Amino acid and fatty acid composition of *Alysicarpus longifolius* seeds. *Indian J Pharmaceut Sci* **47**:171-172.

Jain, R. and Gupta, R.K. (1986b) Flavonoid glucosides from *Alysicarpus longifolius*. *Fitoterapia* **57**:183-184.

Jain, R. and Gupta. R.K. (1981) Chemical investigation of *Alysicarpus longifolius* leaves. *J Indian Chem Soc* **58** :819-820.

Jain. R. and Gupta, R.K. (1984) Structure of phegopolin. *Indian J Chem* **23B** :999-1000.

Johansen, D.A. (1940). *Plant Microtechnique*, (1st Edn). McGraw Hill Book Company. New York.

Jong Ting Ting: Hwang Chien Chih (1995). Two rare isoflavones from *Celosia argentia*. *Planta Medica*; **61(6)**: 584-585.

Joshi AR, Joshi K (2000). Indigenous knowledge and uses of medicinal plants by local communities of the Kali Gandaki Watershed Area, Nepal. *J Ethnopharmacol* ;73: 175-183.

Joshi K.C, Prakash, L. and Singh, P. (1973). Quinones and other constituent from *Tabebuia rosea*, *Phytochemistry* 12: 942.

Joshi P. (1991). Herbal druds used in guinea worm diseases by the tribals of southern Rajasthan (India), *Int J Pharmacog*; 29:33-38.

Jyothi B., Sudarsanam G., Bulusu Sitaram and . Prasada Babu G (2010). Pharmacognosy of a local market sample of parpataka Polycarpaea corymbosa (L.) Lam. *International Journal of Plant Sciences*,5 (2) : 613-619.

jyothi B., sudarsanam G., Bulususitaram and prasadababu G. (2010) Pharmacognosy of a South Indian market sample of parpataka *Rungia repens*(L.) Nees *International Journal of Plant Sciences*, 5 (2) : pp 404-411.

Kako M, Miura T., Nishiyama Y., . Ichimaru M, . Moriyasu M, and Kato A.,(1996.) “Hypoglycemic effects of the rhizomes of *Polygala senega* in normal and diabetic mice and its main component, the triterpenoid glycoside senegin-II,” *Planta Medica*, 62(5): 440–443.

Kapoor. I. D., Singh. A ., Kapoor. SI and Srivasta, S.N.(1969). Survey of Indian plants for saponins. alkaloids and flavonoids. I. *Lloydia* ;32: 297-304.

Karawayya *et al.*, (1980) phytochemical studies of abrus precatorius alkaloids, *Herba hung* ; 19: 21-25.

Katkar KV , Suthar AC , Chauhan VS_ (2010). The chemistry, pharmacologic, and therapeutic applications of *Polyalthia longifolia*.*Pharmacognosy Review*; 4(7) : 62-68.

Katselis G. S, . Estrada A, . Gorecki D. K. J, and Barl, B.(2007).Adjuvant activities of saponins from the root of *Polygala senega L.*,” *Canadian Journal of Physiology and Pharmacology*, **85(11)**: 1184–1194.

Kaur S, Dayal R, Varshney VK, Bartley JP (2001). GC-MS analysis of essential oils of heartwood and resin of *Shorea robusta*. *Planta Med*; **67(9)**: 883-886.

Khaleqe A, Aminuddin M, Mulk SAU .(1996) Investigations on *Abrus precatorius*. I.Constituents of dry root. *Pak CSIR Bull Monogr* **3**:203-207.

Khare C.P. (2007) Indian Medicinal Plants, An Illustrated Dictionary, Springer Science + Business Media, LLC,USA, pp.393.

Kindscher K.,(1992) *Medicinal Wild Plants of the Prairie*, University of Kansas Press, Lawrence, Kan, USA.

Kirtikar K.R, Basu BD (1985). *Indian Medicinal Plants*, Vol. III.Lalit Mohan Basu, editor. Allahabad .p.138.

Kirtikar KR and Basu BD. (1995). *Indian Medicinal Plants*. International Book Distributors, Dehradun; p.562.

Kirtikar KR, Basu BD (1994). *Indian Medicinal Plants*. Vol. III. 2ndedn. International Book Distributors , Dehradun ; pp.1908-09.

Kuo S-C *et al.*,(1995) potent antiplatelet, anti-infiammatory and antiallergic isoflavanquinones from the roots of *Abrus precatorius*, *Planta Med* ; **67**: 307-312.

KuoS-C, *et al.*, (1999)Abruquinone derivatives useful in the treatmeni of inflammation and allergic reactions. *Patent- US* , **5** : 856,352.

Lacaille-Dubois M.-A. and Mitaine-Offer A.-C.,(2005) “Triterpenesaponins from Polygalaceae,” *Phytochemistry Reviews*, **4(2,3)** : 139–149.

Laping, J. (1984). Pluralism in the ayurvedic system of medicine. *Anc.Sci.life* ;4(2): 83.

M Aminur Rahman *et.al*, (2010) Analgesic And Antiinflammatory Activity Of Methanolic Extract Of *Acalypha Indica* Linn., *Pak. J. Pharm. Sci.*, **23(3)**;256-258.

Mabry, T.J., Markham and M.B./Thomas (1970). *The Systematic Identification of Flavonoids*. Springer – Verlage, Berlin.

Majumdar, Kaviraj A. (1989). Ayurveda and modern medicine. *Anc.Sci.life*; **8(3&4)**: 117.

Malati G and Pillai APG. (2005). Microscopic profile of powdered drugs used in Indian systems of medicine; **Vol-1**: Bark drugs. Institute of Ayurvedic Medicinal Plant Sciences, Jamnagar;p.50.

Malati G and Pillai APG. (2005). *Microscopic profile of powdered drugs used in Indian systems of medicine*; **Vol-1**: Bark drugs. Institute of Ayurvedic Medicinal Plant Sciences, Jamnagar; pp.88-89.

Malati G and Pillai APG. (2005). Microscopic profile of powdered drugs used in Indian systems of medicine; **Vol-1**: Bark drugs. Institute of Ayurvedic Medicinal Plant Sciences, Jamnagar.pp.20-21.

Malati G and Pillai APG. (2005). Microscopic profile of powdered drugs used in Indian systems of medicine; **Vol-1**: Bark drugs. Institute of Ayurvedic Medicinal Plant Sciences, Jamnagar.

Manandhar, N.P. (2002). *Plants and People of Nepal* Timber Press. Oregon.

Mangalan S.(1988).*Chemosystematics of some selected taxa of the Caryophyllales*. Ph.D. Thesis, M.S. University of Baroda.

Markham, K.R. (1982). *Techniques of Flavonoid Identification*, AP, London.

Methanolic Extract Of *Acalypha Indica* Linn., *Pak. J. Pharm. Sci.*, **23(3)**;256-258.

Meyers, S. F. H., Jawetz E. And Goldfine (1972). *Review of Medical Pharmacology*, 3rd edn. Large medical publications, California.

Miler, A.L (1996). Antioxidant flavonoids: structure, function and clinical usage. *Altern. Med. Rev.* **1**: 103.

Mishra LN, Ahmed A.(1997). Triterpenoids from *Shorea robusta* resin. *Phytochem*; **45(3)**: 575-578.

Mooss, N.S. (1980).*Ganas of Vahata*, Kottayam.p.63.

Mukherjee, J. and Roy. B. (1971), Chemical examination of *Salmania malabarica* Schott A: Hull. syn. *Bombax malabaricum* DC. *J Indian Chem Soc* ;**48** : 769-770.

Muthaliar Murugesa (1988). *Materia Medica (Vegetable section)*, **Vol- I**, 4th

Nadkarni K M (1954) The indian materia medica, **Vol-I** pp.760.

Nadkarni KM (1954) *Indian Materia Medica*. **Vol. I**. Popular Prakashan Mumbai ; p.580

Nadkarni KM, Nadkarni AK (2002). *Indian MateriaMedica*.**Vol. I**. Popular Prakashan, Mumbai ; p. 1081.

Nahrstedt A. *et.al*, (September 2006)'Flavonoids from *Acalypha indica*' *Fitoterapia* **77(6)**;484-486.

Nevin KG and Vijayammal PL (2003). *Effect of Aerva lanata* on solid tumour induced by DLA cells in mice. *Fitoterapia*; **74**: pp 578-582.

Niranjan. G.S. and Gupta. P.C. (1973). Anthocyanins from the flowers of *Bombax malabaricum*. *Planta Med*; **24**: 196-199.

Nishanthini, A. Mohan, V.R (2013). Antioxidant activity of *polycarpa corymbosa* l.

Nishihama Y, Masuda K, Yamaki M, Takagi S and Sakina K (1981). Iridoid glycosides isolated from *Oldenlandia corymbosa* L. *Planta med*; **43**:28-33.

Noiarsa P, Ruchirawat S, Otsuka H and Kanchapoom. T (2008). Chemical constituents from *Oldenlandia corymbosa* L. of Thai origin. *Nat Med* (Tokyo); **62(2)**:249-250.

Nordin H.J and Ahmad. R (2006). Phytochemical Studies and Pharmacological activities of Plants in Genus *Hedyotis/Oldenlandia*. Studies in Natural Products Chemistry, **Vol-33**, Copyright Elsevier.

Ogunwande IA *et al*, (2010)Essential oil constituents and biological activities of *Peristrophe bicalyculata* and *Borreria verticillata*. *Nat Prod Commun*. Nov; **5(11)**:1815-8.

Ou Ming. (1990). *An Illustrated Guide to Antineoplastic Chinese Herbal Medicine*. The Commercial Press, Hong Kong. p. 313.

Paech, K. and Tracey M. V. (1965). *Modern methods of plant analysis II*. Springer-Verlag, Berlin.

Pandey CN, Raval BR, Mali S and Salvi H (2005). *Medicinal plants of Gujarat*. Gujarat Ecological Education and Research Foundation, Gandhinagar.

Pandey M.B, Singh A.K, Singh J.P, Singh V.P, Pandey V.B (2008). Fuyuziphine, a new alkaloid from *Fumaria indica*. *Nat Prod Res*; **22(6)**:pp. 533–536.

Pandey VB, Das Gupta B, Bhattacharya SK, Lal R, Das PK (1971). Chemistry and pharmacology of the major alkaloid of *Fumaria indica*. *Curr Sci*; **40**: pp 455-457.

Pandey VB, Das Gupta B, Ray AB (1979) .Minor Alkaloid of *Fumaria indica* seeds.*Phytochemistry*; **18**: p 695.

Pandey VB, Das Gupta B, Ray AB. (1974) Letters to the editor, *Curr Sci.*; **43**: p 749.

Pandey VB, Tripathi VK (1992). Stem alkaloid of *Fumaria indica*. *Phytochemistry*; **31**: p 2189.

Pandey VB, Tripathi YC, Pathak NKR, Biswas M (1988). A Seco-phthalideisoquinoline alkaloid from *Fumaria indica*. *Phytochemistry*; **27**: pp 1918-1919.

Patra A, Dey AK, Kundu AB. (1992). Shoreaphenol, a polyphenol from *Shorea robusta*. *Phytochem*; **37(7)**: 2561-2562.

Pengelly, A. (2006). *The constituents of medicinal plants*, Allen & unwil, Australia, P. 126.

Pervykh LN, Zapesochnaya GG and Karasatov BS (1992). A study of the herb *Aervalanata. IV. Flavonoid glycosides*. *Khim Prir Soedin*; **5**: pp 581-583.

Pharmaceutical Botany by Heber W. Youngken, .(1951). 7 th edn, The Blackiston Company , Toronto. p.559.

Pradhan P., Joseph L., Gupta V., Chulet R., Arya H., Verma R. and Bajpai A. (2009) *Saraca asoca* (Ashoka): A Review. *Journal of Chemical and Pharmaceutical Research* ; **1 (1)**:62-71.

Prakash EO, Rao JT. (1999). A new flavone glycoside from the seeds of *Shorea robusta*. *Fitoter*; **70(6)**: 539-541.

Prakash, A. and Khosa, R.L. (1976). Chemical studies on *Bauhinia variegata*. *Curr. Sci* ; **45**:705.

Prakash, A. and Khosa, R.L. (1978). Neutral constituents of *Bauhinia variegata*. *J. Res. Indian Med. Yoga. Homeopath*; **13**: 96-97.

Raj J. and Singh K.P. (2000) *CCRH Quarterly Bulletin* (Drug Standardization Special - I)

Raja R.Vinoth *et al* (2009) Studies on wound healing property of coastal medicinal plants,

Rao,P.G *et.al*,(1982) *J.Proc.Indian Acad. Sci.(Ser) Plant Sci.*,**91**;495.

Rastogi RP, Mehrotra BN.(1979) Compendium of Indian Medicinal Plants. Lucknow: CDRI and NewDelhi: Publication and Information Directorate; p. 201.

Rastogi R.P, Mehrotra B.N (1970-79). *Compendium of Indian Medicinal Plants. Vol-II*.Lucknow: Central Drug Research Institute; p. 357.

Rath Bikash (2005). Globalisation, global trend in herbal market and the impact thereof on medicinal plants in Orissa www.vasundharaorissa.org reprint, Peiodical Experts, New Delhi.

Retnam K R & Martin P. *Ethnomedicinal Plants* . Agrobios , Jodhpur.

Sadashivan S, Latha PG and Sasikumar. (2006). Hepatoprotective studies on *Hedyotis corymbosa*. *J Ethnopharm*; **106**(2): 245.

Salisbury, E.J. (1932). The interpretation of soil climate and the use of stomatal frequency as an interesting index of water relation to the plant. *Beih. Bot. Zentralb.* **49**: 408.

Sankar SS, Joseph KJ, Nair AGR (1964). Occurrence of isosalipurposide in the flowers of *Rungia repens*, *Ind J Chem*; **8**: pp 338-39.

Saoji A.G., Saoji A. N. and Deshmukh V.K. (1972). Presence of lawsone in *Ammannia baccifera* Linn and *Woodsonia fruticosa* Salisb., *Curr Sc*; **41**:192.

Saraswathy A.,*et al.*, (2006). Pharmacognostic studies on *Peristrophe paniculata*. *Journal of Medicinal and Aromatic Plant Sciences*. **28**:209-215.

Satish S, Bhakuni DS (1972). Constituents of Indian and other plants. *Phytochem*; **11**: pp 2888-2890.

Savithramma N, Sulochana C, Rao K.N (2007). Ethnobotanical survey of plants used to treat asthma in Andhra Pradesh, India. *J Ethnopharmacol* ; **113**: 54-61.

Saxena HO, Brahman M. (1995) *The Flora of Orissa, Vol. III*, Bhubaneswar (India): RRL.

Saxena VK, Sharma DN.(1999) A new isoflavone from the roots of *Abrus precatorius*. *Fitoterapia*; **70**: 328-329.

Sayeed Ahmad*, Madhukar Garg, Maksood Ali, Mhaveer Singh, Md Tanvir Athar and Shahid Husain Ansari,(2009). A phyto-pharmacological overview on *Adhatoda zeylanica* Medic. syn. *A. vasica*(Linn.) Nees *Natural Product Radiance*, **Vol. 8(5)**, , pp.549-554.

Selvam R, Kalaiselvi P, Govindraj A, Balamurugan V and Sathishkumar AS (2001). *Effect of Aerva lanata* leaf extract and Vediuppuchurnam on the urinary risk factors of calcium oxalate urolithiasis during experimental hyperoxaluria. *Pharmacol Res*; **43**: pp 89-93.

Seshadri TR, Vydeeswaran S (1972). Chrysoeriol glycosides and other flavonoids of *Rungia repens* flowers. *Phytochemistry* ; **2**: pp 803-6.

Singh N. *et.al.*, (2011). Phytochemical Screening and Blood Glucose level lowering effect of *Bergenia ligulata* root extracts, *Journal of Natural Remedies*; **11(1)**:19 – 23.

Singh R.H and Singh K.P (1990). Perspective in plant drug reseaech. *Anc.Sci.Life* ;**9(3)**: 154.

Singh RS.,et. al,(2000).Aliphatics from *Peristrophe bicalyculata*.Fitoterapia.**71(1)**: 80-1.

Singh,R. Jain,S.C. Jain, R (2009). Antioxidant activity of some medicinally important arid zone plants. *Asian Journal of Experimental Sciences* **23 (1)** :215-221.

Sivarajan V V and Indira B (1994). *Ayurvedic Drugs and their Plant Sources*.Oxford and IBH Publishing Co. Pvt. Ltd, Delhi.

Somanadhan B, Varughese G, Palpu P, Sreedharan R, Gudikshen L, Smith UV and Nyman U (1999). An ethnopharmacological survey for potential angiotensin converting enzyme inhibitors from Indian medicinal plants. *J. Ethnopharmacol*; **65**: pp 103-112.

Soni S. , Anandjiwala Sheetal , Patel G. , and Rajani M. (2008). Validation of Different Methods of Preparation of *Adhatoda vasica* Leaf Juice by Quantification of Total Alkaloids and Vasicine *Indian J Pharm Sci.* **70(1)**: 36–42.

Soni S. , Anandjiwala Sheetal, Patel G. , and Rajani M. (2008).Validation of Different Methods of Preparation of *Adhatoda vasica* Leaf Juice by Quantification of Total Alkaloids and Vasicine *Indian J Pharm Sci* ;**70(1)**: 36–42.

Srinivasan, S. et al. (1971). Effects of some flavonoids an the surface change characteristics of the vascular system and their antithromabogenic characteristics. *European Conf. Microcircuaation* **6**: 394.

Srivastava C.K., Wali J.B., ChakravartyB.K.,Dhawan R.B. and Rastogi R.P. (1978).Screening of Indian plants for biological activity,*Indian J ExpBiol*; **16**:330-349.

Sutarjadi, Santosa M.H, Bendryman and Dyatmiko W. (1991).Immunomodulatory activity of *Hedyotis corymbosa*, grown in Indonesia. *Planta Medica*; **57**: 136.

Swain S. R., Sinha B. N. and Murthy P. N. (2008) Toxicological studies of the hydroalcoholic extract of *rungia repens* leaves.*Pharmacology on line*; Vol. **3**: pp 58-64.

Tagaki S, Yamaki M, Masuda K, Nishihama Y and K Sakina. (1981). Six iridoids isolated and identified. *Yakugaku Zasshi*; **101**(7): 659.

Tandon,J.S.,Agrawal,K.P. and Dharm,M.M. (1996) Chemistry of the triterpenoids from *Randia dumetorum* Indian J. Chem.**4** (11):483-487.

Tchamo D. Noungoué, Cartier G., Dijoux-Franca M.-G., Tsamo E. and Mariotte A.-M.(2001). Xanthones and Other Constituents of *Trema orientalis*. *Pharmaceutical Biology*; **39**(3):202-205.

Thakkar S.M., Deshmukh V.K., Saoji A.N. and DuragkarN.J. (1986). Chemical examination of *Ammannia baccifera* (Lythraceae),*J Indian ChemSoc*; **63**:619-620.

Theodore Cooke (1967).*The Flora of the Presidency of Bombay* ,Vol.-II.BSI, Calcutta.

Thokchom P. Singh, Okram M. Singh, and Huidrom B.(2011) Singh *Adhatoda vasica* Nees:

Trease GE, Evans WC.(2002) *Pharmacognosy*, 15th edn. W.B. Saunders, Elsevier Science

Tripathi HC, Suresh C and Jawahar L (1985).Cardiovascular effects of *Aerva lanata*. *Indian J. Pharmacol*; **17**: p 42.

Trump, B. S. (1961). A method for staining epoxy section for light microscopy. *J. Ultra Structure. Research* **5**: 343-348.

Udupa, K.N, (1983). Ayurveda for the Service of common name. *Anc. Sci. Life.* **2**(3): 117.

Udupihille M and Jiffry MTM (1986). Diuretic effect of *Aerva lanata* with water, normal saline and coriander as controls. *Ind J Phys Pharmacol*; **30**: pp 91-97.

Usmanghani K, Saeed A, Alam M.T (1997). *Indusyunic Medicine*. Karachii: University of Karachii Press;pp. 240-241.

Vaidyaratnam et al., (1994) *Indian Medicinal Plants* a compendium of 500 species Vol.1,Orient Longman Ltd., Madras. p.10.

Vetrichelvan T and Jagadeeshan M (2002).Antidiabetic activity of the alcohol extract of *Aerva lanata* in rats.*J. Ethnopharmacol*; **80**: pp103-107.

Vetrichelvan T, Jagadeeshan M, Paniappan MS, Murali NP and Sasikumar K (2002).Diuretic and anti-inflammatory activities of Aervalanata in rats. *Indian J. PharmaceutSci*; **62**: pp 300-302.

Vileges, J.H., DeMarchi E. and Lancas E.M.,(1997).Phytochemical studies of some medicinal plants.*Anal*; **8**: 266-270.

Wallis TE. *Textbook of Pharmacognosy*. (1960) 4th edition. J. and A. Churchill, London.Pg.413.

Wallis,T.E. (1953). *Practical Pharmacognosy*, J. and A. Churchill Ltd., Gloucester Place, W.I.

Wang L. and Chen H.P. (2005) Study on Chemical Constituents in the Volatile Oil of *Coleus amboinicus* Lour. by Gas Chromatography-Mass Spectrometry,*Journal of Chinese Mass Spectrometry* ; **26** (1): 62.

Warrier PK, Nambiar VPK and Ramankutty C (1994). *Indian Medicinal Plants* a compendium of 500 species. Vol-II, Orient Longman Ltd. Madras; p.289.

Warrier PK, Nambiar VPK and Ramankutty C (1994). *Indian Medicinal Plants* a compendium of 500 species Vol-V, Orient Longman Ltd. Madras,p.124.

Warrier PK, Nambiar VPK and Ramankutty C (1994).*Indian Medicinal Plants- A Compedium of 500 Species*. Vol.- I, Orient Longman Pvt. Ltd.;pp 67-69.

Wassel GM and Amnar NM (1987). *Phytochemical study of Aerva lanata*. *Fitoterapia*; **58**: p 367.

Watt G (1972) “*A Dictionay of The economic product of Indian*”, Vol. I & VI, second

Whole plant in alloxan induced diabetic rats, *World Journal Of Pharmacy And Pharmaceutical Sciences* **2(1)** : 343-351.

Svanidze NV (1986). Investigation of the herb *Aerva lanata*.I. O-Acylglycosides of flavonoids. *Khim Farm Zh*; **20**; pp855-858.

Zapesochnaya GG, Kurkin V, Okhanov V and Miroshnikov A (1992). Canthin-6-one and beta-carboline alkaloids from *Aerva lanata*. *PlantaMed* ; **58**: pp 192-196.

Zapesochnaya GG, Pervykh LN, Kurkin V, Okhanov V and Miroshnikov A (1991).A study of the herb Aervalanata.III. Alkaloids. *ChemNat Comp.*; **27**: pp 336-340.

Zapesochnaya GG, Pervykh LN, Kurkin V, Okhanov V and Miroshnikov A (1992).The structure of alkaloids of *Aerva lanata*. *Chem Nat Comp*; **27**: pp 725-728.

Zhang Ailian, Qi Huayi , Ye Qi , Zhang Guolin (2006). Chemical Study on *Rostellularia procumbens* *Chinese Journal Of Applied & Environmental Biology* ;**12**: 02.

Zhao Y.Y, Cui C.B, Cai B, Han B. and Sun Q.S. (2005). A new phenanthraquinone from the stems of *Bauhinia variegata* Linn. *J.Asian Nat.Prod.Res*; **7(6)**: 835-38.

Zore G.B. *et al.*, (2008) Chemoprofile and bioactivities of *Taverniera cuneifolia* (Roth) Arn.: a wild relative and possible substitute of *Glycyrrhiza glabra* L. *Phytomedicine : International j. of phytotherapy & phytopharmacology* , 1-10.