

ACKNOWLEDGEMENT

The work for this thesis was carried out in The Pharmacology Laboratory, Research Division, Alembic Chemical Works Co. Ltd., Baroda, under the guidance of Dr. A.H.Amin, Ph.D.(Maryland), Ph.D.(Edin.), Director of Research, Alembic Chemical Works Co. Ltd., Baroda.

I take this opportunity to acknowledge my indebtedness to Dr. A.H. Amin for his valuable guidance and help, which I received from him in the course of the work incorporated in this thesis.

I also express my sincere thanks to Dr.O.D. Gulati, M.D.(Pharmacology); M.S.(Colorado), Senior Professor and Head, Department of Pharmacology, Medical College, Baroda and to Dr. H.M. Parikh, M.D.(Pharmacology) for their valuable comments and suggestions.

My thanks are due to Management, Alembic Chemical Works Co., Ltd., Baroda for providing all the facilities required in this investigation.

I gratefully acknowledge the generous gift of: McN-A-343 by McNeil Laboratories, U.S.A.,

paromomycin by Parke Davis & Co. (India), viomycin by Ciba Research Centre (Bombay).

I am also grateful to Mr. C.N.Kachhia for the excellent typing of the manuscript, Mr. R.R. Patel for photographic reproduction of tracings and to my colleagues for their co-operation and suggestions.

- S.S.SAMARTH.