



Index

Introduction		
	Introduction	01
I	<i>Streptomyces</i> , Antibiotics and its regulation	01
II	History and definition of Polyketide	08
III	Polyketides as secondary metabolites	13
IV	Polyketide synthase (PKS)	17
V	Aromatic Polyketide synthases	21
V.1	Fungal Aromatic PKS	21
V.2	Type III Polyketide Synthase	23
V.3	Aromatic Polyketide synthase Type II	25
V.3.1	Phosphopantetheinyl Transferase	26
V.3.2	Acyl Carrier Protein	27
V.3.3	β -Ketoacyl Synthase	27
V.3.4	Cyclase and aromatase	27
V.3.5	Tailoring enzymes	28
VI	Cloning and sequencing of the Genes	29
VII	Role of tailoring enzymes in the biosynthesis and their role in synthesizing new molecules	34
VIII	About peptide antibiotics	38
Materials and Methods		
MI	Materials: Chemicals and Biochemicals	39
MII	Culture Media for <i>Streptomyces</i>:	41

MII.1	Solid media	41
MII.2	Liquid Media	44
MIII	Media used for culturing <i>E.coli</i>	45
MIII.1	Solid Media	45
MIII. 2	Liquid media	45
MIV	Reagents for plasmid preparation	46
MV	Buffers	47
MVI	Solutions for agarose gel electrophoresis	48
MVII	Miscellaneous reagents	50
MVIII	Solutions for Southern hybridization transfer	51
MIX	For radioactive southern hybridization	52
MX : Methods		
MX. 1	Maintenance of <i>streptomyces</i> strains	53
MX. 2	Maintenance of <i>E. coli</i> strains	53
MX. 3	Maintenance of bacteriophages	53
MX. 4	Growth Condition for <i>Streptomyces</i>	53
MX. 5	Growth condition for <i>E coli</i>	53
MX. 6	Titration of lambda phage	53
MXI. 7	Plasmid isolation (Alkali lysis method)	54
MX. 8	Total DNA isolation of <i>Streptomyces</i>	54
MX. 9	Preparation of competent cells	55
MX. 10	Electroporation <i>E. coli</i>	55
MX. 11	Transformation of <i>E. coli</i> (using calcium chloride method)	56
MX. 12	Transformation (<i>Streptomyces</i>)	56

MX. 12.1	Preparation of protoplasts	56
MX. 12.2	Preparation of protoplasts	57
MX. 12.3	Transformation of protoplasts	57
MX. 12.4	Spot-transformation	57
MX. 12.5	Preparation of single stranded DNA for protoplast transformation by denaturation by alkali	58
MX. 13	Transduction – λ lysate preparation	58
MX. 13.1	Broth Method	58
MXI. 13.2	Agar Method	58
MX. 13.3	Transduction	59
MX. 14	Conjugation	59
MX. 14.1	Between <i>E.coli</i> and <i>E.coli</i>	59
MX. 14.2	Between <i>E.coli</i> and <i>Streptomyces</i>	60
MX. 14.3	For control	60
MX. 15	Extraction of antibiotic	60
MX. 16	Thin layer chromatography	61
MX. 17(a)	Agarose gel electrophoresis	61
MX. 17(b)	Staining the gel	62
MX. 17(C)	Elution of DNA from agarose gel	62
MX. 18	Bioactivity tests	62

MX. 18.1	Bioautograms	62
MX. 18.2	Agar well assay	63
MX. 18.3.	Test organism	63
MX. 19	HPLC analysis of mycelia extract	63
MX. 20	Southern blotting (non radioactive)	64
MX. 21	Southern blotting using radiolabelled probe and colony hybridisation	64
MX. 22	Sucrose density gradient of <i>Sau3AI</i> partially digested genomic DNA	66
Chapter 1: Genomic DNA Library Construction and deducing the overlap		
	Introduction	68
1	Results and discussion	69
1.1	Construction of genomic DNA library and colony hybridization	69
1.1.1	Size fractionation of the genomic DNA	69
1.2	Preliminary characterization of library	74
1.3	Colony hybridization of the genomic DNA library	76
1.4	Analysis of the digestion pattern of the cosmid clones.	77
1.5	Southern hybridization of the different cosmid clones	82
1.5.1	Southern hybridization of <i>EcoRI</i> digested cosmid clones	82
1.5.2	Southern hybridization of <i>BglII</i> digested cosmid clones	83
1.5.3	Southern hybridization of <i>PstI</i> digested cosmid clones	84
1.6	Conclusion	85

Chapter 2 : Construction of Restriction Map		
	Introduction	87
2.1	Results and discussion	91
2.1.1	Restriction Map of 5.0 kb <i>EcoRI</i> fragment	91
2.2	Restriction digestion analysis of 11 kb <i>EcoRI</i> fragment	94
2.2.1	<i>Eco RI</i> , <i>Bgl II</i> single and <i>Eco RI</i> – <i>Bgl II</i> double digestion of the 11.0 kb (+)	94
2.2.2	<i>Eco RI</i> , <i>PstI</i> single and <i>Eco RI</i> – <i>PstI</i> double digestion of the 11.0 kb(+) <i>EcoRI</i> fragment	96
2.2.3	<i>EcoRI</i> , <i>BamHI</i> single and <i>EcoRI</i> – <i>BamHI</i> double digestion of the 11.0 <i>Eco RI</i> fragment	98
2.2.4:	<i>PstI</i> , <i>BglII</i> single and <i>PstI</i> – <i>BglII</i> double digestion of the 11.0 (+) <i>Eco RI</i> fragment	100
2.2.5	<i>BamHI</i> digestion of the 11.0 Δ <i>PstI</i> (in both +/- orientation)	102
2.2.6	Single and double digestion of the 11.0 kb (-) orientation clone with respect to four enzymes <i>BamHI</i> , <i>BglII</i> , <i>PstI</i> and <i>EcoRI</i> .	104
2.3	<i>BamHI</i> , <i>PstI</i> and <i>BamHI-PstI</i> double digestion of 4.0 kb <i>EcoRI</i> fragment	107
2.4	Digestion analysis of 8.0 kb Clone	109
2.4.1	Single and Double digestion of the 8.0 kb <i>EcoRI</i> fragment w.r.t three enzymes <i>PstI</i> , <i>BglII</i> and <i>BamHI</i> .	110
2.4.2:	Single and Double digestion of the 8.0 kb <i>EcoRI</i> fragment (in both (+) /(-) orienatation) with respect to three enzymes <i>EcoRI</i> , <i>BglII</i> and <i>BamHI</i> .	115
2.4.3	Single and Double digestion of the 8.0 kb <i>EcoRI</i> fragment with restriction enzymes <i>EcoRI</i> and <i>PstI</i>	119

2.4.4	Single and Double digestion of the 8.0 kb(+) <i>EcoRI</i> fragment with enzymes <i>BglII</i> and <i>PstI</i> .	122
2.4.5	8.0 kb <i>EcoRI</i> , digestion with enzymes <i>EcoRI</i> and <i>BamHI</i>	124
2.4.6	<i>BglII</i> , <i>BamHI</i> and <i>PstI</i> digestion of the 4.5kb <i>PstI</i> fragment	126
2.4.7	<i>BglII</i> , <i>BamHI</i> and <i>PstI</i> digestion of the 4.5kb <i>PstI</i> fragment	128
2.4.8	Digestion of 8.0 kb Δ <i>PstI</i> with <i>PstI</i> , <i>BglII</i> and <i>BamHI</i>	131
2.4.9	<i>PstI</i> , <i>BglII</i> and <i>BamHI</i> digestion of 8.0 kb Δ <i>PstI</i> in (-) orientation and 8.0 kb Δ <i>BglII</i>	132
2.5	Digestion analysis of 17.0 kb <i>EcoRI</i> fragment	134
2.5.1	<i>EcoRI</i> , <i>BglII</i> , <i>PstI</i> single and <i>EcoRI</i> – <i>BglII</i> and <i>EcoRI</i> – <i>PstI</i> double digestion of the 17.0 kb <i>EcoRI</i> fragment derived from 1.1 cosmid clone.	134
2.6	Conclusion	136
Chapter 3: Sequence analysis		
3.1	Introduction	139
3.2.1	Generating subclones for sequencing 8.0 kb <i>EcoRI</i> DNA	139
3.2.2	Analysis of Sequence of the 8.0 kb of PKS DNA	141
3.2.2.1	Sequencing result of 8.0 kb <i>EcoRI</i> fragment.	141
3.2.3	Sequencing result of 5.0 kb <i>EcoRI</i> fragment.	148
3.3	Conclusion	153
Chapter 4: Functional analysis of tailoring enzymes		
	Introduction	154
4.1	General strategy used for construction of a gene deletion mutant	156

4.2	General strategy used for the disruption of the gene function	157
4.3	Results and discussion	158
4.3.1	Deletion of TDP-glucose-4,6-dehydratase gene	158
4.3.1.1	Construction of the vector, 2Δ <i>Bam</i> HI for mutagenesis	158
4.3.1.2	Construction of TDP-glucose-4,6-dehydratase (DH) mutant using homologous recombination strategy	161
4.3.1.3	<i>trans</i> complementation of the DH ⁻ mutan	163
4.3.1.4	TLC and HPLC analysis of the culture extract of the deletion mutant	163
4.3.2	Insertion Mutagenesis of the TDP-glucose-4,6-dehydratase gene	166
4.3.2.1	Construction of the vector for insertion mutagenesis	166
4.3.2.2	Construction of insertion mutant using homologous recombination strategy	168
4.3.2.3	Phenotypic characteristic of the insertion mutant	169
4.3.2.4	Genetic confirmation of the mutant	169
4.3.2.5	Bioautogram, TLC and HPLC analysis of the insertion mutant	170
4.3.3	Insertion Mutagenesis of O-Methyltransferase gene	172
4.3.3.1	Construction of the (OMT) disruption vector	173
4.3.3.2	Construction of insertion mutant using homologous recombination strategy	174
4.3.3.3	Phenotypic characteristics of the mutant	174
4.3.3.4	Extraction of bio-active component, and TLC analysis of the insertion mutant	175
4.4	Conclusion	176
	Summary and Perspective	178
	Bibliography	185