

THE UNIVERSITY OF CHICAGO PRESS

- Abdul-Wahab, A. S. and E. L. Rice. (1967). Plant inhibition by Johnson-grass and its possible significance in old field succession. Bull. Torrey Bot. Club. 94 : 486-497.
- Ahn, Byung Zun, Ulrike, Degen, Charoon Lienjayetz etc., Peter Pachaly and Felix Zymalkowski. (1978). Constituents of Cassia siamea. Arch. Pharm. (Wein Heim) 311 (7) : 569 - 578.
- Akkasaeng, R., Gutteridge, R. C. and M. Wanapat. (1989). Evaluation of trees and shrubs for forage and fuelwood in Northeast Thailand. Int. Tree Crops Journal. 5 (4) : 209-220.
- Alam, S. M. and A. R. Azmi. (1989). Influence of Prosopis glandulosa water extract on the seedling growth of wheat cultivars. Pak. J. Sci. Ind. Res. 32 : 708.
- Alam, M. M., Siddiqui, S. A. and A. M. Khan. (1977). Mechanism of control of plant-parasitic nematodes as a result of the application of organic amendments to the soil : 3 : Role of phenols and amino acids in host roots. Ind. J. Nematol. 7 (1) : 27 - 31.
- Al-Naib, F. A. and E. L. Rice. (1971). Allelopathic effects of Plantanus occidentalis. Bull. Torrey Bot. Club. 98 : 75-82.
- Alsaadawi, I. S., Arif, M. B. and A. J. AlRubeaa. (1985). Allelopathic effects of Citrus aurantium L. II. Isolation, characterization, and biological activities of phytotoxins. J. Chem. Ecol. 11 : 1527-34.

- Altieri, M.A. (1989). Weed Ecology. In Agroecology : The Scientific Basis of Alternative Agriculture. Boulder, Colorado : Westview. pp 173-85.
- Altieri, M. A. and J. D. Doll. (1978). The potential of allelopathy as a tool for weed management in crop fields. Pans. 24 : 495-502.
- Anaya, A. L., Gliessman, S. R., CruzOrtega, R., Rosado-May, F. and V. Nava Rodriguez. (1988). Effects of allelopathic weeds used as cover crops on the floristic potential of soils. In Proc. of the Sixth Int. Sci. Conf. of the Int. Fed. Org. Agric. Movements. Patricia Allen and Debra Van Dusen (eds.). Vol. Two : 607-613.
- Anaya, A. L., Ramos, L., Cruz O., R. and J. Hernandez. (1987). Studies of allelopathy in Mexico. In Allelochemicals : Role in agriculture and forestry, G. R. Waller (ed.). American Chemical Society. Symposium Series, 330 : 89-101.
- Anaya, A. L., Roy-Ocotla, G., Ortiz, L. M. and Y. L. Ramos. (1982). Potencial allelopatico de las principales plantas de un cafetal. En Estudios ecologicos en el agroecosistema cafetalero, E. Jimenez Avila and A. Gomez - Pompa (eds.) Simposiodel INIREB.
- Aneja, R., Khanna, R. N. and T.R., Seshadri. (1963). 6-methoxyfuranoflavone, a New Component of the seeds of P. glabra. J. Chem. Soc., 23 : 163.

- Aoki, T. and T. Suga. (1978). Triterpenoid saponins from the flower buds of Fatsia japonica. Phytochemistry. 17 : 771-773.
- Arntzen, C.J., Falkenthal, S.V. and S. Bobick. (1974). Inhibition of photophosphorylation by kaempferol. Plant Physiol. 53 : 304-306.
- Asplund, R. O. (1968). "Monoterpenes : Relationship between structure and inhibition of germination". Phytochemistry. 7 : 1995-1997.
- Athar, M. and A. Mahmood. (1980). A qualitative study of the nodulating ability of legumes of Pakistan. List 2. Trop. Agric. 57 (4) : 319 - 324.
- Azami, A. R. and S. M. Alam. (1989). Effect on some wild plants reduces and wheat straw on germination and growth of wheat cultivars. Cer. Res. Commun. 17 : 59-62.
- Balasubramanian, M. and D. Purshothaman. (1972). Indole acetic acid in the eriophyid mite galls on P. glabra Vent. caused by Eriophyes cherianii Massee. (Eriophyidae : Acarina). Labdev J. Sci. Technol. Part B. Life Sci. 10(3/4) : 172-173.
- Balke, N.E. (1977). Inhibition of ion absorption in Avena sativa L. roots by diethylstilbestrol and other phenolic compounds. Ph.D. Thesis, Purdue Univ., W. Lafayette, Indiana. Diss. Abstr. No. 7813025.
- Balke, N.E., Davis, M.P. and C.C. Lee. (1987). Conjugation of Allelochemicals by Plants. In. Allelochemicals : Role in

- Agriculture and Forestry, G.R. Waller (ed.). American Chemical Society. Washington, D.C., pp 214-227.
- Balke, N.E. and T.K. Hodges. (1977). Inhibition of ion absorption in oat roots : Comparison of diethylstilbestrol and oligomycin. Plant. Sci. Lett. 10 : 319-325.
- Bansal, G.L. (1988). Allelopathic effects of aqueous extracts of stem and leaves of three tree species on the germination of some crops and weeds. In Proc. Trends in Tree Science. Solan (India). Indian Society of Tree Scientists. pp. 118-23.
- Bansal, G. L. (1989). Allelopathic potential of linseed on Buttercup (Ranunculus arvensis L.) In Plant Science Research in India. Trivedi, M. L., Gill, B. S. and S. S. Saini (eds.). Today and Tomorrow's Printers and Publishers, New Delhi-5. pp 801-805.
- Bapna, Saroj., Renapurkar, D. M. and S. R. Chavan. (1988). Cyclopical activity of three indigenous plant extracts. Insect. Sci. Appl. 9 (2) : 211-212.
- Barnes, J.P., Putnam, A.R. and A. Burke. (1986). Allelopathic activity of rye (Secale cereale L.). In The Science of Allelopathy. Putnam, A.R. and C.S. Tang (eds.). John Wiley and Sons, New York. pp. 271-286.
- Barz, W. and J. Koster. (1981). In "The Biochemistry of Plants, A Comprehensive Treatise". Stumpf, P.K. and Conn, E.E., (eds.), Academic Press. New York, Vol. 7, Chapter 3.

- Baveja, S. K., Rao K. V. R. and J. Arora. (1989). Examination of natural gums and mucilages as sustaining materials in tablet dosage forms. Part II. Ind. J. Pharm. Sci. 51 (4) : 115-118.
- Behmer, D.E. and T.M. McCalla. (1963). The inhibition of seedling growth by crop residues in soil inoculated with Penicillium urticae Bainer. Plant Soil. 18 : 119-206.
- Bell, E. A. (1981). The physiological role(s) of secondary (natural) products. In The Biochemistry of Plants. E. E. Conn (ed.) Academic Press, New York, Vol. 7 : 1-19,
- Benoit, R.E. and R.L. Starkey. (1968a). Enzyme inactivation as a factor in the inhibition of decomposition of some organic matter by tannins.
- Benoit, R.E. and R.L. Starkey. (1968b). Inhibition of decomposition of cellulose and some other carbohydrates by tannin. Soil. Sci. 105 : 291-296.
- Bergmark, Christine, L., William A. Jackson, Richard J. Volk and Udo Blum. (1992). Differential inhibition by ferulic acid of nitrate and ammonium uptake in Zea mays L. Plant Physiol. (Bethesda) 98 (2) : 639-645.
- Bhatia, R. K. and D. D. Chawan. (1976). Occurrence of phenolic substances in seed coat of Cassia species and their effect on early seedling growth. Geobios (Jodhpur) 3 (6) : 214-216.
- Bisla, S., Nandal, D. P.S. and S. S. Narwal. (1992). Influence of aqueous leaf extracts of Eucalyptus and poplar on the germination and seedling growth of winter crops. In Proc. Forest Biology in the Service of Mankind. pp. 95-97.

- Biswas, K. M. and H. Mallik. (1986). Cassiadinine, a chromone alkaloid and dextro-6-hydroxymellein, a dihydroisocoumarin from Cassia siamea. Phytochemistry (OXF.). 25 (7) : 1727-1730.
- Blair, G. J., Masud Panjaitan, Ivory, D. A., Palmer, B. and M. Sudjadi. (1988). An evaluation of tree legumes on an acid ultisol in south Sumatra, Indonesia. J. Agric. Sci. 111 (3) : 435-442.
- Blum, U., Dalton, B. R. and J.O. Rawlings. (1984). Effects of ferulic acid and some of its microbial metabolic products on radicle growth of cucumber. J. Chem. Ecol. 10 (8) : 1169-1191.
- Blum, U. and E.L. Rice. (1969). Inhibition of symbiotic nitrogen - fixation by gallic and tannic acid and possible roles in old - field succession. Bull. Torrey Bot. Club. 96 : 531-544.
- Bogdan, G.P. (1977). Mutual effect of couch grass and cultivated plants in phytocenoses. In "Interactions of plants and Microorganisms in Phytocenoses". A.M. Grodzinsky (ed.), Naukova Dumka, Kiev. pp. 36-43.
- Bogdan, G.P. and A.M. Grodzinsky. (1974). Role of sulfhydryl groups in protective reactions of plants during allelopathic damage. Ukr. Bot. Zh. 30 : 771-772.
- Bonner, J. (1965). The Isoprenoids. In Plant biochemistry. James Bonner and J. E. Varner (eds.) Academic Press, New York. pp. 665-692.

- Bonner, J. and A. W. Galston. (1944). Toxic substances from the culture media of the guayule which may inhibit growth. Botan. Gaz. 106 : 185-198.
- Borner, H. (1959). The apple replant problem. I. The excretion of phlorizin from apple root residues. Boyce. Thompson. Inst. Contrib. 20 : 39-56.
- Borner, H. (1960). Liberation of organic substances from higher plants and their role in the soil sickness problem. Bot. Rev. 26 : 393 - 424.
- Boucher, D. H., Espinoza, J., Romero, S. and S. R. Gliessman. (1983). Out-of-season planting of grain legumes as green manures for a tropical raised field agroecosystem. Biol. Agri. and Hort. 1 : 127-133.
- Bradow, J. M. (1985). Germination regulation by Amaranthus palmeri and Ambrosia artemisiifolia. Am. Chem. Soc. Symp. Ser. 268 : 285-99.
- Bradow, J. M. and W. J. Connick, Jr. (1988). Volatile methyl ketone seed-germination inhibitors from Amaranthus palmeri, S. Wats. residues. J. Chem. Ecol. 14 (7) : 1617-1632.
- Brain, P. W. (1957). Effects of antibiotics on plants. Ann. Rev. Plant Physiol. 8 : 413-426.
- Breman, J. W. and D. L. Wright. (1984). Using winter legume mulches as a nitrogen source for no tillage corn and grain sorghum production. Proc. South. Region No Till. Conf., Agric. Exp. Stn., Auburn Univ., Auburn, AL. pp. 6-17.

- Caceres, Armando, Brenda. R. Lopez, Melba A. Giron and Heidi Logemann. (1991). Screening of antimycotic activity of plants used in Guaetmala for the treatment of dermatophytoses. Rev. Mex. Micol. 7 (0) : 21-38.
- Cameron, H.J. and G.R. Julian. (1980). Inhibition of protein synthesis in lettuce (Lactuca sativa L.) by allelopathic compounds. J. Chem. Ecol. 6 : 989-995.
- Campbell, N. E. R., and H. Lees. (1967). The nitrogen cycle. In Soil Biochemistry, McLaren, D. A. and G. H. Peterson (eds.), Marcel Dekker, Inc., New York. pp. 194-215
- Capasso, R., Cristinzio, G., Evidente, A. and F. Scognamiglio. (1992). Isolation, spectroscopy and selective phytotoxic effects of polyphenols from vegetable waste waters. Phytochemistry 31 (12) : 4125 - 4128.
- Casal, J. F., Reigosa, M. J. and A. Carbelleira. (1985). Allelopathic potential of Acacia dealbata Link. Revue d'Ecologie et de Biologie du sol. 22 : 1-12.
- Chan, Gian and P. C. Shukla. (1973). Use of unconventional feeds in broiler rations. Ind. J. Anim. Sci. 43 (1) : 1013-1017.
- Chandra, U., Saxena, S.K. and M. Akram. (1981). Sphaerotheca fuliginea, causal organism of powdery mildew of Indian beans. Acta Bot. Indica. 9 (1) : 69-73.
- Chandramohan, D., Purushothaman, D. and R. Kothadaraman. (1973). Soil Phenolics and Plant growth inhibition. Plant and Soil, 39 : 303-308.

- Chatterjee, A. and S. R. Bhattacharjee. (1964). New Dianthraquinones from Cassia siamea Lam. Part I. Structure of Cassianin and Siameanin. J. Ind. Chem. Soc. 41 : 415.
- Chatterjee, A. and S. R. Bhattacharjee. (1965). Bull. Nat. Inst. Sci. India. 141.
- Chavan, P. B. and U. V. Kulkarni. (1974). Additions to the fungi from Maharashtra, India. Maharashtra Vidnyan Mandir Patrika. 9 (112) : 132-139.
- Chavan, P.B. and S.K. Patil. (1972). Studies in some rust fungi from India. Sydowia Ann. Mycol. 26 (1-6) : 277-281.
- Chopra, R. N., Chopra, I. C., Handa, K. L. and L. D. Kapur. (1982). Chopra's Indigenous Drugs of India. Second Edn. Academic Publishers, New Delhi.
- Chou, Chang-Hung and Mei-Hui, Hou. (1981). Allelopathic researches of subtropical vegetations in Taiwan : 1. Evaluation of allelopathic potential of bamboo (Phyllostachys edulis) vegetation. Proc. Natl. Sci. Council. Repub. China. Part B. Basic Sci. 5 (3) : 284-292.
- Chou, Chang-Hung and Lih-Ling, Leu. (1992). Allelopathic substances and interactions of Delonix regia (Boj.) Raj. J. Chem. Ecol. 18 (12) : 2285-2303.
- Chou, C. H. and C. H. Muller. (1972). Allelopathic mechanisms of Arctostaphylos glandulosa var. zacaensis. Amer. Midl. Natur. 88 : 324-347.

- Chou, C. H. and Y. L. Kuo. (1986). Allelopathic exclusion of understorey by Leucaena leucocephala (Lam) de Witt. J. Chem. Ecol. 12 : 1431-48.
- Colvin, B. M., Harrison, L. R., Sangster, L. T. and S. G. Harvey. (1986). Cassia occidentalis toxicosis in growing pigs. J. Amer. Vet. Med. Assoc. 189 (4) : 423-426.
- Cooper-Driver, G., Corner-Zamodits, J.J. and T. Swain. (1972). Z. Naturforsch. 27b : 943-946.
- Corcoran, M.R., Geissman, T.A. and B.D. Phinney. (1972). Tannins as gibberellin antagonists. Plant Physiol. 49 : 323-330.
- Cornman, I. (1946). Alteration of mitosis by coumarin and parasorbic acid. Am. J. Bot., 33 : 217.
- Cosio, E.G. and J.W. McClure. (1984). Plant Physiol. 74 : 877-881.
- Cutler, A.J. and E.E. Conn. (1982). Recent Adv. Phytochem. 16 : 249-271.
- Daiya, K. S., Sharma, H. K., Chawan, D. D. and D. N. Sen. (1980). Effect of salt solutions of different osmotic potential on seed germination and seedling growth in some Cassia spp. Folia Geobot. Phytotaxon. 15 (2) : 149-154.
- Dalal, M. R., Dahiya, D. S., Sarmah, M. K. and S. S. Narwal. (1992). Supression effects of Arid Zone Trees on Plant stand and Growth of Crops. In Proc. Forest Biology in the Service of Mankind. pp. 132-135.

- Danks, M.L., Fletcher, J.S. and E.L. Rice. (1975). Effects of phenolic inhibitors on growth and metabolism of glucose-UL-¹⁴C in Paul's Scarlet Rose cell-suspension cultures. Am. J. Bot. 62 : 311-317.
- Danso, A. A. and P. Morgan. (1993). Alley cropping maize (Zea mays-var-Jeka) with Cassia (Cassia siamea) in the Gambia : Crop production and Soil fertility. Agroforestry Syst. 21 (2) : 133-146.
- Daulatabad, C. D., Hosamani, K. M. and A. M. Mirajkar. (1988). Cassia siamea seed oil : A Minor source of vernolic and cyclopropenoid fatty acids : J. Am. Oil Chem. Soc., 65 (6) : 952-953.
- DeBell, D. S. (1971). Phytotoxic effects of cherrybark oak. For. Sci., 17 : 180-185.
- del Moral, R. and C. H. Muller. (1969). Fog drip : a mechanism of toxin transport from Eucalyptus globulus. Bull. Torrey Bot. Club. 96 : 467-475.
- del Moral, R. and C. H. Muller. (1970). The allelopathic effects of Eucalyptus camaldulensis. Amer. Midl. Natur. 83 : 254-282.
- Deshmukh, S. D. and M. N. Borle. (1975). Studies on the insecticidal properties of indigenous plant products. Ind. J. Entomol. 37 (1) : 11-18.


- Dutta, N. L., Ghosh, A. C., Nair, P. M. and K. Venkataraman. (1964). The structure of Cassiamin. A new plant pigment. Tetrahedron Lett. 40 : 3023-3030.
- Dzyubenko, N. N. and N. I. Petrenko. (1971). On biochemical interactions of cultivated plants and weeds. In Physiological biochemical basis of plant interactions in phytocenoses, A. M. Grodzinsky, (ed.) Vol. 2. Kiev : Naukova Dumka.
- Einhellig, F.A. and L. Kuan. (1971). Effects of scopoletin and chlorogenic acid on stomatal aperture in tobacco and sunflower. Bull. Torrey Bot. Club. 98 : 155-162.
- Einhellig, F.A. and G.R. Leather. (1988). Potentials for exploiting allelopathy to enhance crop production. J. Chem. Ecol. 14 : 1829-1844.
- Einhellig, F. A. and J. A. Rasmussen. (1979). Effects of three phenolic acids on chlorophyll content and growth of soybean and grain sorghum seedlings. J. Chem. Ecol. 5 (5) : 815-824.
- Einhellig, F.A., Rice, E.L., Risser, P.G. and S.H. Wender. (1970). Effects of scopoletin on growth CO_2 exchange rates and concentration of scopoletin, scopolin and chlorogenic acids in tobacco, sunflower and pigweed. Bull. Torrey Bot. Club. 97 : 22-33.
- Elakovich, S. D. and K. L. Stevens. (1985). Volatile constituents of Lippia nodiflora. J. Nat. Prod. 48 : 504-506.

- Elakovich, S. D. and B.O. Oguntimein. (1987). The essential oil of Lippia adoensis leaves and flowers. J. Nat. Prod. 50 : 503-506.
- Elliott, L. F. and J. W. Elaylock. (1975). Effects of wheat straw and alfalfa amendments on solubilization of manganese and iron in soil. Soil Sci., 120 : 205-211.
- Espinoza - Moreno, J. (1984). Allelopathic effects of Chenopodium album and Amaranthus retroflexus on rhizobial nodulation and growth of Phaseolus vulgaris. Master's thesis, University of California, Santa Cruz.
- Eyini, M., Jayakumar, H. and S. Pannirselvam. (1989). Allelopathic effect of bamboo leaf extract on the seedlings of groundnut. Trop. Ecol. 30 : 138-41.
- Fischer, N.H. (1986). The function of mono and sesquiterpenes as plant germination and growth regulators. In The Science of Allelopathy. Putnam, A.R. and C.S. Tang.(eds.). Willey Interscience, New York, pp. 203-218.
- Fischer, N. H., Olivier, E. J. and H.D. Fischer. (1979). The biogenesis and chemistry of sesquiterpene lactones. Prog. Chem. Org. Nat. Prod. 38 : 47-390.
- Fischer, N. H. and L. Quijano. (1985). Allelopathic agents from common weeds Amaranthus palmeri, Ambrosia artemisiifolia and related weeds. Am. Chem. Soc. Symp. Ser. 268 : 133-47.

- Floyd, G. L. and E. L. Rice. (1967). Inhibition of higher plants by three bacterial growth inhibitors. Bull. Torrey Bot. Club. 94 : 125-129.
- Fowden, L. (1963). Amino-acid analogues and the growth of seedlings. J. Exp. Bot. 14 : 387-398.
- Gaba, R.K. (1987). Roles of Allelopathy in Social Forestry. In Social Forestry for Rural Development. Khosla, P.K. and R.K. Kohli. (eds.) ISTS, pp. 228-234.
- Galvez, Guillermo. (1974). Amplitud de hospedantes, purificación y. microscopio electrónico del virus del. mosaico de la soya. (variety of hosts, purification and electron microscopy of the soybean mosaic virus) Rev. Inst. Colomb. Agropecu. 9 (2) : 159-182.
- Gant, R. E. and E. E. C., Clebsch. (1975). The allelopathic influences of Sassafras albidum in old field succession in Tennessee. Ecol. 56 : 604-615.
- Garg, G. P. (1979). A new component from leaves of Pongamia glabra. Planta Med. 37 (1) : 73-74.
- Garg, G. P., Sharma, N.N. and R. N. Khanna. (1978). Two new furano compounds, glabra I and glabra II from the stem bark of Pongamia glabra. Ind. J. Chem. Sect. B. Org. Chem. Incl. Med. Chem. 16 (8) : 658-661.
- Garg, G. P. and R. N. Khanna. (1983). Chemical examination of flowers in Pongamia glabra. Int. J. Crude Drug Res. 21 (1) : 43-47.

- Gaur, Y. D. (1989). Nodulation studies in species of genus Cassia. Zentralbl Bakteriol. 135 (3) : 201-204.
- Gbeassor, M., Kossou, Y., Amegbo, K., De Souza, C., Koumaglo, K. and A. Denke. (1989). Antimalarial effects of eight African medicinal plants. J. Ethnopharmacol. 25 (1) : 115-118.
- Geissman, T. A. (1962). The occurrence of flavonoid compound in nature. In Chemistry of Flavonoid Compounds. T. A. Geissman (ed.) Pergamon Press, London. pp. 1-5.
- Geissman, T.A. and B.O. Phinney. (1972). Tannins as gibberellin antagonists. Plant Physiol. 49 : 323-330.
- Gibbs, R. D. (1974). Chemotaxonomy of Flowering Plants. McGill Queens University Press, Montreal.
- Gillon, D., Adam, F., Hubert, B. and G. Kahlem. (1983). Seed production and seed consumption in a Sahelo-Sudanian woodland in Senegal : The ecological balance sheet. Terre vie. 38 (1) : 3-36.
- Glass, A.D.M. (1974). Influence of phenolic acids upon ion uptake. II. A structure-activity study of the inhibition of phosphate uptake by benzoic acid derivatives. In Mechanisms of regulation of plant growth. Bielecki et al., (eds.) Roy. Soc. N. Z. Bull. 12 : 159-164.
- Glass, A. D. M. and B. A. Bohm. (1971). The uptake of simple phenols by barley roots. Planta. 100 : 93-105.
- Goel, U. and T. S. Sareen. (1986). Allelopathic effect of trees on the understorey vegetation. Acta. Bot. Indica. 14 (2) : 162-166.

- Gray, R. and J. Bonner. (1948a). An inhibitor of plant growth from the leaves of Encelia farinosa. Am. J. Bot. 35 : 52-57.
- Gray, R. and J. Bonner (1948b). Structure determination and synthesis of a plant growth inhibitor 3-acetyl-6-methoxy benzaldehyde, found in the leaves of Encelia farinosa. J. Am. Chem. Soc. 70 : 1249-1253.
- Green, F.B. and M.R. Corcoran. (1975). Inhibitory action of five tannins on growth induced by several gibberellins. Plant Physiol. 56 : 801-806.
- Grodzinsky, A.M. and G.P. Bogdan. (1972). Histochemical study of pectins, lignin, suberin and melanins in plants treated with allelopathically active substances. Ukr. Bot. Zh. 29 : 137-143.
- Grodzinsky, A.M. and G.P. Bogdan. (1973). Role of ascorbic acid in formation of the brown mass in xylem of plants under the effect of allelopathic factors. Ukr. Bot. Zh. 30 : 28-35.
- Grummer, G. (1955). Die gegenseitige. Beeinflussung hoherer Pflanzen - Allelopathie, Gustav Fisher Verlag, Jena.
- Grummer, G. (1961). The role of toxic substances in the interrelationships between higher plants. In Mechanisms in biological competition. Cambridge Univ. Symp. Cambridge Press. pp. 219-228.
- Gu, Mao-Bim. (1983). Biology and Control of Catopsilia pomona. Acta Entomol. Sin. 26 (2) : 172-178.
- Gudauskas, R., David, T., Teem, H. and Gareth M-J. (1977). Anthracnose of Cassia occidentalis caused by Colletotrichum



- dematium f. truncata. Plant Dis. Rep. 61 (6) : 468-4702
- Guenzi, W.D. and T.M. McCalla. (1966). Phytotoxic substances extracted from soil. Soil Sci.Soc.Amer.Prox. 30
- Guenzi, W.D., McCalla, T.M. and F.A.Norstadt. (1967). Presence and persistence of phytotoxic substances in wheat, oat, corn and sorghum residues. Agron J. 59 : 163-165.
- Guha, S., Basu, S. and C.V.N. Rao. (1985). Structural features of the degraded polysaccharide from Pongamia glabra gum. Ind. J. Chem. 24 (2) : 171-174.
- Gupta, H.C., Verma, J.P., Bareth, S.S. and B.N. Mathur. (1988). Evaluation of some non-edible oils as grain protection in wheat and their subsequent effect on germination. Ind. J. Entomol. 50 (2) : 147-150.
- Halligan, J.P. (1975). Toxic terpenes from Artemisia californica Ecol. 56 : 999-1003.
- Harborne, J. B. (1964). Phenolic glycosides and their natural distribution. In Biochemistry of phenolic compounds. J. B. Harborne (ed.) Academic Press, New York. pp. 129-169.
- Harborne, J. B. (1967). Comparative Biochemistry of Flavonoids. Academic Press, London.
- Harborne, J.B. (1973). Phytochemical methods. Chapman and Hall, London.
- Harborne, J.B. (1980). In "Encyclopedia of Plant Physiology". Bell, E.A. and B.V. Charlwood (eds.). Springer-Verlag, New York. Vol. 8. pp. 329-402.

- Harborne, J. B. (1984). Phytochemical methods (2nd edn.) Chapman and Hall, London.
- Harborne, J. B. and J. J. Corner. (1961). Plant polyphenols. Four Hydroxycinnamic acid-sugar derivatives. Biochem. J. 81 : 242-250.
- Harborne, J. B. and N.W. Simmonds. (1964). The natural distribution of the phenolic aglycones. In Biochemistry of phenolic compounds. J. B. Harborne (ed.) Academic Press, New York. pp. 77-127,
- Harper, J.R. and N.E. Balke. (1980). Inhibition of potassium absorption in excised oat roots by phenolic acids. In "Plant Membrane Transport : Current Conceptual Issues". Spanswick, R.M., Lucas, W.J. and J. Dainty (eds.). Elsevier/North-Holland, New York. pp. 399-400.
- Hassan, Y. M., EL-Hindawy, S., Bassiony, S. and M. A., Abdalla. (1974). Cassia occidentalis L. as coffee substitute in Egypt. Egypt. J. Hortic. 1 (1) : 137-143.
- Hauser, S. (1993). Effect of Acioa barteri, Cassia siamea, Flemingia macrophylla and Gmelina arborea leaves on germination and early development of maize and cassava. Agric. Ecosyst. Environ. 45 (3-4) : 263-273.
- Henderson, M.E.K. (1963). Fungal metabolism of certain aromatic compounds related to lignin. Pure and Appl.Chem. 7:589-602.
- Henniquin, J-R. and C. Juste. (1967). Presence D'acides phenols libres dans le sol. Etude de leur influence sur la germination et la croissance des vegetaus. Ann. Agron. 18 : 545-569.

- Hogetsu, T., Shibaoka, H. and M. Shimokoriyama. (1974).
Involvement of cellulose in actions of gibberellin and
kinetin-coumarin interactions on stem elongation. Plant Cell
Physiol. 15 : 265-272.
- Horsley, S. B. (1977). Allelopathic interference among plants. II
Physiological modes of action. In Proceedings of the Fourth
North American Forest Biology Workshop. Wilcox, H.E. and
A.F. Hamer (eds.). College of Environmental Science and
Forestry, Syracuse, N.Y., U.S.A. pp. 93-136.
- Hungund, B. and C. H. Pathak. (1971). A survey of Gujarat forests
for sources of alkaloids, saponins and tannins. U.S.D.A.
Forest Service Research Paper. NE-201.
- Hurst, H. M. and N. A. Burges. (1967). Lignin and humic acids. In
Soil Biochemistry. McLaren, D.A. and G.H. Peterson (eds.)
Marcel Dekker, Inc., New York. pp. 260-286.
- Hutchinson, A., Roy, C. and G. H. N. Towers. (1958). Synthesis of
phlorin and other phenolic glucosides by plant tissues.
Nature. 181 : 841-842.
- Ibrahim, R. K. and G.H.N. Towers. (1960). The identification by
paper chromatography of plant phenolic acids. Arch. Biochem.
Biophys. 87 : 125-128.
- Inoue, M., Nishimura, H., Li, H-H. and J. Mizutani. (1992).
Allelochemicals from Polygonum sachalinense Fr. Schm.
(Polygonaceae). J. Chem. Ecol. 18 (10) : 1833-1840.

- Jadhav, B. B. and D. G. Gaynar. (1992). Allelopathic effects of Acacia auriculiformis A. Cunn. on germination of rice and cowpea. Ind. J. Plant Physiol. 35. (1) : 86-89.
- Jain, A.C., Lal, P. and T.R. Seshadri. (1969). Nuclear Prenylation of 2-Methyl-5,7-dihydroxychromone and Resacetophenone : Synthesis of Peucenin, Isopeucenin, Isobavachalkone and Isobavachin. Ind. J. Chem. 7 : 1072-1075.
- Jain, P. P., Suri, R.K., Deshmukh, S. K. and K. C., Mathur. (1987). Fatty oils from oilseeds of forest origin as antibacterial agents. Indian For. 113 (4) : 297-299.
- Jankay, P. and W. H. Muller. (1976). The relationships among umbelliferone, growth and peroxidase levels in cucumber roots. Amer. J. Bot. 63 : 126-132.
- Jeffrey, D. W., Timothy, C. M. and J. T. Romeo. (1987). Solution volume and seed number : Often overlooked Factors in Allelopathic Bioassays. J. Chem. Ecol. 13 (6) : 1481-1491.
- Jimenez - Osornio, J. J., K. Schultz, A. L., Anaya, J. Fernandez and O. Espejo. (1983). Allelopathic potential of corn pollen. J. Chem. Ecol. 9 : 1011 - 1025.
- Jimenez-Osornio, J. J. and S. R. Gliessman. (1987). Plant herbivore interaction in a wild mustard (Brassica campestris L.) and broccoli (Brassica oleraceae L. var. italica) intercrop agroecosystem. In Allelochemicals : Role in Agriculture and Forestry. Symposium series 330. G. R. Waller (ed.). American Chemical Society. Washington, D. C.

- Jobidon, R. and J. R. Thibault. (1981). Allelopathic effects of balsam poplar (Populus balsamifera) on green alder (Alnus crispa var. mollis) germination. Bull. Torrey Bot. Club. 108 (4) : 413-418.
- Jobidon, R. and J. R. Thibault. (1982). Allelopathic growth inhibition of nodulated and unnodulated Alnus crispa seedlings by Populus balsamifera. Am. J. Bot. 69 (8) : 1213-1223.
- Joshi, P.C. and O. Prakash. (1992). Allelopathic effects of litter extract of some tree species on germination and seedling growth of agricultural crops. Proc. First Nat. Symp. "Allelopathy in Agroecosystems". Tauro, P. and S.S. Narwal (ed.), Indian Society of Allelopathy, Har. Agri. Univ., Hisar, India. pp. 127-128.
- Kanungo, Das, P., Ganguly, A., Guha, A., Bhattacharyya, A. and N. A. Chaudhury. (1987). Glabone, a new furanoflavone from Pongamia glabra. Phytochemistry. 26 (12) : 3373-3374.
- Keck, R.W. and T.K. Hodges. (1973). Membrane permeability in plants : Changes induced by host specific pathotoxins. Phytopathology. 63 : 226-230.
- Khailov, K. M. (1974). Biochemical trophodynamics in marine coastal ecosystems (ed.). Naukova Dumka, Kiev. (In Russian).
- Khan, Gohar, Kapoor, V. P. and M. I. H. Farooqi. (1988). Structural investigation of Cassia siamea L. Seed galactomannan. Ind. J. Chem. Sect. B. Org. Chem. Incl. Med. Chem. 27 (9) : 821-824.

- Khanna, R.N. and T. R. Seshadri. (1963). Pongaglabrone, a new component of the seeds of Pongamia glabra : Its constitution and synthesis. Tetrahedron. 19 : 219-225.
- Khosla, S. N. and S. N. Sobti. (1981). Parthenin : A promising root inhibitor from Parthenium hysterophorus. Ind. J. For. 4 (1) : 56-60.
- Kirtikar and Basu. (1933). "Indian Medicial Plants" Vol. I. 2nd Edn., p. 830.
- Koepe, D.E. (1972). Some reactions of isolated corn mitochondria influenced by juglone. Physiol. Plant. 27 : 89-94.
- Kohli, R.H., Kaur, K., Chaudhary, P., Kumari, A. and D.B. Saxena. (1987). Negative aspects of Eucalyptus farming. In Agroforestry for Rural Needs. Khosla, P.K. and D. K. Khurana (eds.). Solan (India). Indian Society of Tree Scientists. pp. 225-241.
- Komai, K., Suguguka, Y. and S. Sato. (1981). Plant growth retardant of extracts obtained from water nutgrass (Cyperus serotinus Rottb.). Chem. Abst. 95 : 162961 C.
- Kononova, M. M. (1961). Soil organic matter. Pergamon Press, New York.
- Kopp, B., Loffehardt, W. and W. Kubelka. (1978). Z. Naturforsch. 33c : 646-650.
- Krylov, Y.V. (1970). Influence of potatoes on an apple tree and its photosynthesis. In "Physiological-Biochemical Basis of Plant Interactions in Phytocenoses". Grodzinsky, A.M. (ed.),

- Kuo, Y.L., Chou, C.H. and T.W. Hu. (1982). Allelopathic potential of Leucaena leucocephala. In Proc. Seminar on allelochemicals and pheromones. Taipei (Taiwan). pp. 107-119.
- Ladipo, J. L. (1988). Viruses associated with a mosaic disease of Crotalaria juncea in Nigeria : I. Cowpea mosaic virus. J. Phytopathol (Berl.). 121(1) : 8-18.
- Lakshmi, P., Srimannarayana, G. and N. V. Subba Rao. (1974). Pongaflavone, a new chromeno - chromone and an analogue of karanjin isolated from Pongamia pinnata (Linn.) Pierre (Syn. *P. glabra*).
- Leather, Gerald, R. (1983). Sunflowers (Helianthus annuus) are allelopathic to weeds. Weed Sci. 31 (1) : 37-42.
- Lee, D. W., Veronica, H. Y. and F.Y. Liew. (1975). Lectins in selected Malaysian legumes. Malays. J. Sci. 3 : 89-93.
- Lee, T.T. (1966). Effects hydroxybenzoic acids on oxidation of reduced nicotinamide adenine dinucleotide by enzymes from tobacco leaves. Physiol. Plant. 19 : 660-671.
- Lim, G. and H. L. Ng. (1977). Root nodules of some tropical legumes in Singapore. Plant Soil. 46 (2) : 317-327.
- Limaye, D. B. (1936). Rasayanam. 1 : 1.
- Lodhi, M.A.K. and E. L. Rice. (1971). Allelopathic effects of Celtis laevigata. Bull. Torrey Bot. Club. 98 : 83-89.

- Lohar, D. R., Chawan, D. D. and S.P. Garg. (1975). Phytochemical studies on Cassia species of Indian Aridzone. Curr. Sci. 44 (2) : 67.
- Lorber, P. and W. H. Muller. (1976). Volatile growth inhibitors produced by Salvia leucophylla. Effects on seedling root tip ultrastructure. Am. J. Bot. 63 : 196-200.
- Lorber, P. and W.H. Muller. (1980). Volatile growth inhibitors produced by Salvia leucophylla. Effects on metabolic activity in mitochondrial suspension. Comp. Physiol. Ecol. 5 : 68-75.
- Ly, C. Y., Lu, K.C., Trappe, J. M. and W. B. Balleu. (1970). Separation of phenolic compounds in alkali hydrolysates of a forest soil by thin layer chromatography. Can. J. Soil Sci. 50 : 458-460.
- Mabry, T. J., Markham, K. R. and M. B. Thomas. (1970). The Systematic Identification of Flavonoids. Springer-Verlag, New York.
- Makkar, H. P. S., Singh, B. and S. S. Negi. (1990). Tannin levels and their degree of polymerization and specific activity in some agroindustrial by-products. Biological wastes. 31 (2) : 137-144.

- Malik, S. B., Seshadri, T.R. and Pushpa Sharma. (1976). Minor components of the leaves of P. glabra. Ind. J. Chem. 14 (3) : 229-230.
- Malik, S.B., Sharma, P. and T. R. Seshadri. (1977a). Furano-flavonoids from the leaves of Pongamia glabra. Ind. J. Chem. Sect. B. Org. Chem. Incl. Med. Chem. 15 (6) : 536-538.
- Malik, S. B., Sharma, P. and T. R. Seshadri. (1977b). Conversion of Glabrachromene-I into corresponding flavanone, flavone, aurone and flavonol. Ind. J. Chem. Sect. B. Org. Chem. Incl. Med. Chem. 15 (6) : 539-540.
- Mandal, B., Ghosh, S., Majumdar, I. and C. R. Maity. (1985). Protease inhibitors and invitro protein digestibility of defatted seed cakes of akashmoni (Acacia auriculaeformis) and karanj (Pongamia glabra). J. Am. Oil, Chem. Soc. 62 (7) : 1124-1126.
- Mandal, B., Ghosh, S., Majumdar, I. and C. R. Maity. (1986). Isolation and evaluation of protein from processed karanj (Pongamia glabra) seed meal. Acta Aliment. 15 (1) : 69-77.
- Mandal, B., Ghosh, S., Majumdar, I. and C. R. Maity. (1988). A rapid color test for the detection of karanj (Pongamia glabra) oil in other vegetable oils. Fett. Wiss. Technol. 90 (6) : 239-240.
- Mandava, B. N. (1985). Chemistry and Biology of Allelopathic agents. In The Chemistry of Allelopathy : Biochemical Interactions Among Plants. A. Thompson (ed.). ACS Symp. Ser.

- No. 268, Washington, D.C. : American Chemical Society. pp. 33-54.
- Manissery, J. K., Narayana Gowda, B.S., Shankar, K. M. and T. J. Varghese. (1988). Growth of common carp (Cyprinus carpio var. communis) fed on feed incorporated with Cassia tora leaves. Ind. J. Anim. Sci. 58 (6) : 737-740.
- Manthe, B., Margot, S. and S. Heide. (1992). Effects of salicylic acid on growth and stomatal movements of Vicia faba L. Evidence for salicylic acid metabolization. J. Chem. Ecol. 18 (9) : 1525-1539.
- Markham, K. R. (1982). Techniques of Flavonoid Identification, Academic Press, London.
- Martins, E. V., Martins, M. V., Franklin R-C., Ricardo, A.S. and S. V. Paraboni. (1986). Intoxication by Cassia occidentalis (Leguminosae) in swine. Pesqui. Vet. Bras. 6 (2) : 35-38.
- Martin, P. and B. Rademacher. (1960). Studies on the mutual influences of weeds and crops. Brit. Ecol. Soc. Symp. 1 : 143-152.
- Massey, A. B. (1925). Antagonism of the walnuts (Juglans nigra L. and J. cinera L.) in certain plant associations. Phytopathol. 15 : 773-84.
- McCalla, T. M., Guenzi, W. D. and F. A. Norstadt. (1963). Microbial studies of phytotoxic substances in stubble-mulch system. Zeit. Allg. Mikrobiol. 3 : 202-210.

- McCahon, C. B., Kelsey, R. G., Sheridan, R. P. and F. Shafizadeh.
(1973). Physiological effects of compounds extracted from
sagebrush. Bull. Torrey Bot. Club. 100 : 23-28.
- McIlroy, R. J. (1951). The plant glycosides. Edward Arnold and
Co., London.
- Melkania, N. P. (1984). Influence of leaf leachates of certain
woody species on agricultural crops. Ind. J. Ecol. 11 : 82-
86.
- Miller, L.P. (1938). Formation of β -(2-chloroethyl)-d-glucoside
by Gladiolus corms from absorbed ethylene chlorohydrin.
Boyce Thompson Inst. Contrib. 9 : 425-429.
- Miller, L.P. (1939). Synthesis of β -(2-chloroethyl)-d-glucoside
by potato tubers treated with ethylene chlorohydrin. Boyce
Thompson Inst. Contrib. 10 : 139-141.
- Miller, L.P. (1940). Formation of β -O-chlorophenyl gentiobioside
in Gladiolus corms from absorbed O-chlorophenol. Boyce
Thompson Inst. Contrib. 11 : 271-279.
- Miller, L. P. (1943). High yields of β -2-trichloroethyl-d-
glucoside and -2-trichloro-ethylgentiobioside from tobacco
plants treated with chloral hydrate Boyce Thompson Inst.
Contrib. 13 : 185-200.
- Miller, L.P. (1973). In "Phytochemistry", L.P., Miller (ed.).
Reinhold, New York. Chap. 11.

- Mishra, R.R. and V.B. Srivastava. (1977). Comparison of mycoflora associated with certain crop and weed seeds. Acta. Mycol. 13 (1) : 145-150.
- Mittal, O. P. and T. R., Seshadri. (1956). Demethoxykanugin : A New Crystalline Compound from P. glabra. J. Chem. Soc. 426 : 2176.
- Möje, W. (1966). Organic soil toxins. In Diagnostic criteria for plants and soils. H. D., Chapman (ed.). Univ. Calif. Press, Berkeley. pp. 633-569.
- Molisch, H. (1937). Der Einfluss einer pflanze auf die andere - Allelopathie. Gustav Fischer-Verlag, Jena.
- Morita, V. H. (1965). The phenolic acids in organic soils. Can. J. Biochem. 43 : 1277 - 1280.
- Morita, H. (1975). Polyphenols in the lime water extractives of peat. Soil. Sci. 120 : 112-116.
- Morrison, R. I. (1963). Products of the alkaline nitrobenzene oxidation of soil organic matter. J. Soil. Sci. 14 : 201-216.
- Mothes, K. (1955). Physiology of Alkaloids. Ann. Rev. Plant. Physiol. 6 : 393 - 432.
- Mukerjee, S.K., Sarkar, S.C. and T.R. Seshadri. (1969a). Natural occurrence of Tetra-O-methylfisetin in Root and Stem barks of Pongamia glabra. Ind. J. Chem. 7 (12) : 1275.

- Mukerjee, S.K., Sarkar, S.C. and T.R. Seshadri. (1969b). The structure and synthesis of Pongachromene, a new component of Pongamia glabra. Tetrahedron. 25 : 1063-1069.
- Muller, C.H. (1966). The role of chemical inhibition (Allelopathy) in vegetational composition. Bull. Torrey Bot. Club. 93 : 332-351.
- Muller, C. H. (1969). Allelopathy as a factor in ecological process. Vegetatio. 18 : 348-357.
- Muller, W.H., Lorber, P. and B. Haley. (1968). Volatile growth inhibitors produced by Salvia leucophylla : effect on seedling growth and respiration. Bull. Torrey Bot. Club. 95 : 415-422.
- Muller, W.H. and C.H. Muller. (1964). Volatile growth inhibitors produced by Salvia species. Bull. Torrey Bot. Club. 91 : 327-330.
- Murti, P.B.R. and T.R., Seshadri. (1944). Proc. Ind. Acad. Sci. 20A : 279.
- Nadkarni. (1954). "Indian Materia Medica". 3rd ed., p. 1002.
- Naik Satam, P.G. and Bringi, N.V. (1972). Occurrence of (-) Isolonchocarpin and Demethoxykanugin in karanja (Pongamia glabra) seed oil. Ind. J. Chem. 11 (3) : 209-210.
- Nandal, D.P.S., Bisla, S.S. and S.S. Narwal. (1992). Allelopathic influence of Eucalyptus and Poplar leaf extracts in the germination and seedling growth of winter vegetables. In Proc. Forest Biology in the service of mankind. pp. 111-13.

- Narayanaswamy, S., Rangaswami, S. and T.R. Seshadri. (1954). Chemistry of Pongamol. Part II. J.Chem.Soc. 6 : 1871-1873.
- Niemeyer, H.M. (1988). Hydroxamic acids (4-hydroxy-1, 4-benzoxazin-3-ones), defence chemicals in the Gramineae. Phytochem. 27 (11) : 3349-3358.
- Nimbal, C.I., Patil, V.S. and Y.C. Panchal. (1990). Studies on allelopathic effect of honey mesquite. Journal of Maharashtra Agricultural Universities. 15 : 390-391.
- Nishimura, H., Kaku, K., Nakamura, T., Fukazawa, Y. and J. Mizutani. (1982). Allelopathic substances (+) - p-menthane-3, 8-diols isolated from Eucalyptus citriodora Hook. Agric. Biol. Chem. 46 : 319-320.
- Norman, A.G. (1955). The effect of polymyxin on plant roots. Arch. Biochem. Biophys. 58 : 461-477.
- Norman, A.G. (1960). Microbiol products affecting root development. 7th Int. Congr. Soil Sci. Madison, Wis. pp. 531-536.
- Norstadt, F. A. and T.M. McCalla. (1968). Microbially induced phytotoxicity in stubble mulched soil. Soil Sci. Soc. Amer. Proc. 32 : 241-245.
- Ogner, G. (1973). Permanganate oxidation of organic matter leached from forest soil after fertilization with urea. Norw. For. Res. Inst. Rep. 30 : 463-469.

- O'Hara, P.J. and K.R. Pierce. (1974). A toxic cardiomyopathy caused by Cassia occidentalis : I Morphological studies in poisoned rabbits. Vet. Pathol. 11 (2) : 97-109.
- Ohman, J.H. and T. Kommedahl. (1960). Relative toxicity of extracts from vegetative organs of quackgrass to alfalfa. Weeds 8 : 666-670.
- Oleszek, W. (1987). Allelopathic effects of volatiles from some cruciferae species on lettuce, barnyard grass and wheat growth. Plant Soil. 102 (2) : 271-274.
- Oleszek, W. and M. Jurzysta. (1987). The allelopathic potential of alfalfa root medicagenic acid glycosides and their fate in soil environments. Plant Soil. 98 (1) : 67-80.
- Overland, L. (1966). The role of allelopathic substances in the "Smother Crop" barley. Am. J. Bot. 53 : 423-432.
- Owens, L.D. (1969). Toxins in plant disease structure and mode of action. Science. 165 : 18-25.
- Parenti, R. L. and E. L. Rice. (1969). Inhibitional effects of Digitaria sanguinalis and possible role in old field succession. Bull. Torrey Bot. Club. 96 : 70-78.
- Park, K-H, Kim, S-J. and K-H. Hyun. (1993). Brassio-steroid substances in immature Cassia tora seeds. J. Korean Agric. Chem. Soc. 36 (2) : 99-104.
- Patel, K. S. (1977). Phyllosphere microflora of Pongamia glabra. Vent and their ability to degrade various aromatic compounds. Geobios (Jodhpur). 4 (5) : 194-195.

- Patil, V. B., Rama Rao, A.V. and K. Venkataraman. (1970). Cassiamin A, B and C, three 2, 2-Bianthraquinonyls in Cassia siamea. Ind. J. Chem. 8 : 109-112.
- Patrick, Z. A. (1955). The peach replant problem in Ontario. II. Toxic substances from microbial decomposition products of peach root residues. Can. J. Bot. 33 : 461-486.
- Patrick, Z. A. (1971). Phytotoxic substances associated with the decomposition in soil of plant residues. Soil Sci. 111 : 13-18.
- Paulino, V.T., Sanchez, M.J.F., Werner, J.C. and V. Consola. (1987). Allelopathic effects of Eucalyptus in forage growth. Revista De Agricultural (Piraciacoba) 62 : 17-35.
- Pavanram, S. K. and L.R. Row. (1956). Aust. J. Chem. 9 : 132.
- Perez, F.J. (1990). Allelopathic effect of hydroxamic acids from cereals on Avena sativa and A. fatua. Phytochemistry. 29 (3) : 773-776.
- Phillips, R. E., Blevins, R. L., Thomas, G. W., Frye, W. W. and S. H. Phillips. (1980). No tillage agriculture. Science. 208 : 1108-1113.
- Picman, A. K. (1986). Biological activities of sesquiterpene lactones. Biochem. Syst. Ecol. 14 : 225-81.
- Prasad, B., Prasad, R. and J. Prasad. (1986). Evaluation of nitrification retardation property of non-edible oils and

- their influence on yield and nitrogen uptake by wheat in calcareous soils J. Ind. Soc. Soil Sci. 34 (2) : 281-285.
- Pridham, J. B. (1960). The formation and possible function of phenolic glucosides. In Phenolics in health and disease. J. B. Pridham (ed.), Pergamon Press, New York. 9-15,
- Pridham, J. B. and M. J. Saltmarsh. (1960). The formation of phenolic glycosides by germination broadbean (Vicia faba) seeds. Biochem. J. 74-42.
- Putman, A.R. and W.B. Duke. (1978). Allelopathy in agroecosystems. Ann. Rev. Phyto. Pathol. 16 : 432-451.
- Ram, M. R. and K. V. Mallaiah. (1993). Pseudocoreospora pongamiae pinnatae, new species from India, Mycol. Res. 97 (1) : 127-128.
- Ramos, L., A. L. Anaya. and J. Nieto de Pascual. (1983). Evaluation of allelopathic potential of dominant herbaceous species in a coffee plantation. J Chem. Ecol. 9 (8) : 1079-1097.
- Randall H. W., Worsham, A.D. and U. Blum. (1989). Allelopathic potential of Legume debris and aqueous extracts. Weed Sci. 37 : 674-679.
- Ranga Rao, N. and N. S. Subba Rao. (1974). Factors responsible for non nodulating nature of some legumes. Proc. Ind. Natl. Sci. Acad. Part B. Biol. Sci. 40 (6) : 613-617.
- Rangaswami, S. (1946). Curr. Sci. 15A : 127.

- Rangaswami, S. and T. R. Seshadri. (1940). Pongamol, A new crystalline compound from Pongamia oil. Curr. Sci. 9 (4) : 179.
- Rao, M. R., Muraya, P. and P.A. Huxley. (1993). Observations of some tree root systems in agroforestry intercrop situations, and their graphical representation Exp. Agric. 29 (2) : 183-194.
- Rao, M. V. and P.S. Dubey. (1990). Biochemical aspects (Antioxidants) for development of tolerance in plants growing at different low levels of ambient air pollutants. Environ. Pollut. 64 (1) : 55-66.
- Rao, N.S. and P.C. Reddy. (1984). Studies on the inhibitory effects of Eucalyptus (hybrid) leaf extracts in the germination of certain food crops. Ind. For. 110 : 218-222.
- Rao, V. R., Subba Rao, N.S. and K.G. Mukerji. (1973). Inhibition of Rhizobium in vitro by non-nodulating legume roots and root extracts. Plant Soil. 39 (2) : 449-452.
- Rao, J.V.S., Rama Mohan Rao, K. and S.S. Murthy. (1979). Allelopathic effect of some weeds of vegetable crops on the germination and early seedling growth of Bajra (Pennisetum typhoideum). Trop. Ecol. 29 (1) : 5-8.
- Rathore, A. L., Patel, S. L. and G. P. Pali. (1993). Integrated nitrogen management with green manures in rice chickpea cropping system. J. Agron. Crop. Sci. 170 (3) : 158-162.
- Reddy, K.R. (1988). Folk medicine from Chittoor District, Andhra Pradesh, India, used in the treatment of jaundice. Int. J. Crude Drug Res. 26 (3) : 137-140.

- Ribereau-Gayon, P. (1972). Plant phenolics. Oliver and Boyd, Edinburgh. pp. 254.
- Rice, E.L. (1965). Inhibition of nitrogen-fixing and nitrifying bacteria by seed plants. II. Characterization and identification of inhibitors. Physiol. Plant. 18 : 225-268.
- Rice, E. L. (1974). Allelopathy. Academic press, New York.
- Rice, E.L. (1979). Allelopathy - An update. Bot. Rev. 45 : 15-109.
- Rice, E.L. (1984). Allelopathy. 2nd Edn. Academic Press, Inc.
- Rice, E. L. and S. K. Pancholy. (1974). Inhibition of nitrification by climax ecosystems. III. Inhibitors other than tannins. Am. J. Bot. 61 : 1095-1103.
- Rovira, A.D. (1969). Plant root exudates. Bot. Rev. 35 : 35-57.
- Roy, D. and R. N. Khanna. (1979). Structure and synthesis of pongol : A new component from immature seeds of Pongamia glabra. Ind. J. Chem. Sect. B. Org. Chem. Incl. Med. Chem. 18 (6) : 525-528.
- Roy, D., Sharma, N.N. and R. N. Khanna. (1977). Structure and synthesis of isopongaflavone, a new component of the seeds of Pongamia glabra. Ind. J. Chem. Sect. B. Org. Chem. Incl. Med. Chem. 15 (12) : 1138-1139.
- Roy, S., Dutta, A. K. and D.P. Chakraborty. (1982). Amasterol, an ecdysone precursor and growth inhibitor from Amaranthus viridis. Phytochem. 21 : 2417-20.

- Sadique, J., Chandra, T., Thenmozhi, V. and V. Elango. (1987). Biochemical modes of action of Cassia occidentalis and Cardiospermum halicacabum in inflammation. J. Ethnopharmacol. 19 (2) : 201-212.
- Sahrawat, K. L. and S. K. Mukerjee. (1977). Nitrification inhibitors : I. Studies with karanjin, a furano-flavonoid from karanja (Pongamia glabra) seeds. Plant Soil. 47 (1) : 27-36.
- Sahrawat, K. L., Parmar, B.S. and S. K. Mukerjee. (1974). Note on the nitrification-inhibitors in the seeds, bark and leaves of Pongamia glabra Vent. Ind. J. Agri. Sci. 44 (6) : 415-418.
- Saiki, H. and K. Yoneda. (1982). Possible dual roles of an allelopathic compound, cis-dehydromatricaria ester. J. Chem. Ecol. 8 (1) : 185-194.
- Sama, S.K., Krishnamurthy, L., Ramachandran, K. and K. Lal. (1976). Efficiency of an indigenous compound preparation (Liv-52) in acute viral hepatitis : A double blind study. Ind. J. Med. Res. 64 (5) : 738-742.
- Sashida, Y., Nakata, H., Shimomara, H. and Kagaya, M. (1983). Sesquiterpene lactones from Pyrethrum flowers. Phytochemistry. 22 : 1219-1222.
- Satam, P. G. Naik and N. V. Bringi. (1973). Karanjachromene, a new flavone from Pongamia glabra seed oil. Ind. J. Chem. 11 (11) : 1188-1189.

- Saxena, A. P. and K. M. Vyas. (1986). Antimicrobial activity of seeds of some ethnomedicinal plants. J. Econ. Taxon. Bot. 8 (2) : 291-300.
- Schwartz, S. M., Varner, J.E. and W. P. Martin. (1954). Separation of organic acids from several dormant and incubated Ohio soils. Soil. Sci. Soc. Am. Proc. 18 : 174-177.
- Schwimmer, S. (1958). Influence of polyphenols and potato components on potato phosphorylase. J. Biol. Chem. 232 : 715-721.
- Seaman, F. C. (1982). Sesquiterpene lactones as taxonomic characters in the Asteraceae. Bot. Rev. 48 : 121-595.
- Sharma, K.K. (1992). Wheat cultivation in association with Acacia nilotica (L.) Willd ex Del. Field-bund plantation in a case study. Agroforestry systems. 17 : 43-51.
- Sharma, K.M.S., Dhillon, M.S. and K.K. Dhingra. (1967). Presence of germination inhibitors in the leaf leachate of some farm grown trees. Ind. For. 113 : 816-820.
- Sharma, P., Seshadri, R. T. and S. K. Mukerjee. (1973). Some Synthetic and Natural Analogues of glabrachromene. Ind. J. Chem. 11 (10) : 985-986.
- Shivappa Shetty, K., Balasubramanya, R.H., Siddarame Gowda, T.K. and R.B. Patil. (1974). Studies on the disease of Cassia siamea L. caused by Fusarium. Misore J. Agric. Sci. 8 (3) : 384-390.

- Sidiyasa, K. (1986). Tree-flora on the ridges and upper slopes of dry climate area at Poboya Nature Reserve, Central Sulawesi, (Indonesia). Bul. Penelitian Hutan. 0 (485) : 31-38.
- Sigmund, W. (1924). Ueber die Einwirkung, von Stoffwechsel - Endprodukten auf die Pflanzen. Biochem. Z. 146 : 389-419.
- Singh, D. and R. K. Kohli. (1992). Impact of Eucalyptus tereticornis Sm. shelterbelts on crops. Agroforestry Syst. 20 (3) : 253-266.
- Singh, G.B. (1983). Role of agroforestry in improving the environment. Ind. Farm. 33 : 15-19.
- Singh, K. K. and J. K. Maheshwari. (1983). Traditional phytotherapy among the tribals of Varanasi district, Uttar Pradesh (India). J. Econ. Taxon. Bot. 4 (3) : 829-838.
- Sinha, A. (1959). Studies on Sterol from Seeds of Pongamia glabra. Ind. J. Appl. Chem. 22 (2) : 86-88.
- Sinha, B. K. and P. S. Basu. (1981). IAA and its metabolism in root nodules of Pongamia pinnata. Biochem. Physiol. Pflanz (BPP). 176 (3) : 218-227.
- Stahl, C., Vanderhoef, L. N., Siegel, N. and J. P. Helgeson. (1973). Fusarium tricinctum T-2 toxin inhibits auxin-promoted elongation in soybean hypocotyl. Pl. Physiol. 52 : 663-666.
- Stenlid, G. (1968). On the physiological effects of phloridzin, phloretin and some related substances upon higher plants. Physiol. Plant. 21 : 882-894.

- Stevens, K. L. and G.B. Merrill. (1985). Sesquiterpene lactones and allelochemicals from Centaurea species. Am. Chem. Soc. Symp. Ser. 268 : 83-98.
- Stevenson, F. J. (1967). Organic acids in Soil. In Soil Biochemistry. D. A. Mc. Laren and G. H. Peterson (eds.) Marcel Dekker, Inc. New York. pp. 119-146,
- Subramanyam, S. V. (1987). Assessment of utility of some pulp wood species of Kerala State (India) based on fiber quality. Ind. For. 113 (6) : 427-433.
- Subrahmanyam, K., Rao, J.M. and K.V.J. Rao. (1972/73). Isolation of Pongachalcone - I from the heart-wood of Pongamia glabra L. Merr. Curr. Sci. 42 (4) : 128-129.
- Sulekha, M., Sharma, P., Seshadri, T. and S.K. Mukerjee. (1972). Structure and Synthesis of Glabrachromene, A New Constituent of P. glabra. Ind. J. Chem. 10 (6) : 585-588.
- Sullia, Shanker Bhat. (1973a). Effect of root exudates and extracts on rhizosphere fungi. Plant Soil. 39 (1) : 197-200.
- Sullia, Shanker Bhat. (1973b). Soil moisture content and root exudates in relation to rhizosphere effect in leguminous weeds. Proc. Ind Acad. Sci. Sect. B. 77 (6) : 264-275.
- Sunil Puri and A. Khara. (1991). Allelopathic effects of Eucalyptus tereticornis on Phaseolus vulgaris seedlings. Int. Tree Crops. J. 6 : 287-293.
- Swain, T. (1977). Secondary compounds as protective agents. Ann. Rev. Plant Physiol. 28 : 479-501.

- Swaminathan, C., Vinaya Rai, R. S. and K. K. Suresh. (1989).
Allelopathic proclivities of Acacia nilotica (L). Willd. ex
Del. J. Trop For. Sci. 2 (1) : 56-60.
- Talapatra, S. K., Mallik, A. K. and B. Talapatra. (1980).
Pongaglabol, a new hydroxyfuranoflavone and aurantiamide
acetate, a dipeptide from the flowers of Pongamia glabra
Vent. Phytochem. (Oxf.). 19 (6) : 1199-1202.
- Talapatra, S. K., Mallik, A. K. and B. Talaptra. (1982).
Isopongaglabol and 6-methoxy isopongaglabol. 2 new
hydroxyfuranoflavones from Pongamia glabra. Phytochem
(Oxf.). 21 (3) : 761-766.
- Taparia, A.L., Talmale, T.W. and V.V. Sharma. (1978). Utilization
of Cassia tora seeds in growth rations of buffalo calves.
Ind. J. Anim. Sci. 48 (11) : 804-810.
- Thomazini - Casagrande, Lilian Isolde. (1980). Mycorrhiza in
coffee senna plants (Cassia occidentalis). Phyton Rev. Int.
Bot. Exp. 39 (1) : 113-120.
- Tian, G., Kang, B.T. and L. Brussaard. (1992). Effects of
chemical composition on nitrogen, calcium and magnesium
during incubation of leaves from selected agroforestry and
fallow plant species. Biogeochem (DORDR). 16 (2) : 103-119.
- Towers, G.H.N. (1964). Metabolism of phenolics in higher plants
and microorganisms. In Biochemistry of phenolic compounds.
J.B. Harborne (ed.) Academic Press, New York. pp. 249-294.
- Tripathi, R.S. (1977). Weed problems - an ecological perspective.
Trop. Ecol. 18 : 138-148.

- Tripathi, A. K., Gupta, K.R. and J. Singh. (1993). Anthraquinone galactosides from the roots of Cassia siamea. Fitoterapia. 64 (1) : 63-64.
- Tukey, H.B., Jr. (1969). Implications of allelopathy in agricultural plant science. Bot. Rev. 35 : 1-16.
- Tukey, H.B., Jr. (1970). The leaching of substances from plants. Ann. Rev. Plant Physiol. 21 : 305-324.
- Tukey, H.B., Jr. and R.A. Mecklenburg. (1964). Leaching of metabolites from foliage and subsequent reabsorption and redistribution of the leachate in plants. Am. J. Bot. 51 : 737-742.
- Tyson, B.J., Dement, W.A. and H.A. Mooney. (1974). Volatilization of terpenes from Salvia mellifera. Nature. 252 : 119-120.
- Van Sumere, C.F., Cottenie, J. DeGreef and J. Kint. (1972). Biochemical studies in relation to the possible germination regulatory role of naturally occurring coumarin and phenolics. In Advances in Phytochemistry. J.C. Runeckles and J.E. Watkins (eds.). Appleton - Century - Crofts, New York. pp. 165-221.
- Varshney, I.P. and Rajpal. (1978). Chemical studies of the flowers of Cassia siamea Lam., Peltophorum ferrugineum Benth and Caesalpinia pulcherrima S.W. Ind. J. Pharm. 40 (1):15-16.
- Verma, V.S. and F.R. Niazi. (1974). Negro coffee mosaic virus. Z. Pflanzen Pflanzenschutz. 81 (10) : 608-610.

- Wagner, Hildebert, Samia Mohammed El-Sayyad, Otto Seligmann and V. Mohanchari. (1978). Chemical constituents of Cassia siamea Lam. I. 2-Methyl-5-acetonyl-7-hydroxychromone (cassiachromone). Planta Med. 33 (3) : 258-261.
- Waller, G.R. (1989). Allelochemical action of some natural products. In Phytochemical Ecology : Allelochemicals, Mycotoxins and Insect Pheromones and Allomones. Academia Sinica Monograph Ser. 9, Chou, C.K. and G.R. Waller (eds.). Taipei, R.O.C. pp. 129-154.
- Wang, T.S.C., Cheng, S-Y. and H. Tung. (1967a). Extraction and analysis of soil organic acids. Soil Sci. 103 : 360-366.
- Wang, T.S.C., Cheng, S-Y. and H. Tung. (1967b). Dynamics of soil organic acids. Soil. Sci. 104 : 138-144.
- Wang, T.S.C., Yang, T-K. and T-T. Chuang. (1967). Soil phenolic acids as plant growth inhibitors. Soil Sci. 103 : 239-246.
- Wang, T.S.C., Yeh, K-L., Cheng, S-Y. and T-K. Yang. (1971). Behavior of soil phenolic acids. In US. Nat. Comm. Int. Biol. Program. Biochemical interactions among plants. Nat. Acad. Sci. Washington, D.C. pp. 113-120.
- Weaver, T.W. and D. Klarich. (1977). Allelopathic effects of volatile substances from Artemisia tridentata. Nutt. Am. Midl. Nat. 97 : 508-512.
- Westlake, D.W.S., Talbot, G., Blakley, E.R. and F.J. Simpson. (1959). Microbial decomposition of rutin. Can. J. Microbiol. 5 : 621-629.

- Whitehead, D.C. (1963). Some aspects of the influence of organic matter on soil fertility. Soil and Fert. 26 : 217-223.
- Whitehead, D.C. (1964). Identification of -hydroxybenzoic, vanillic, -coumaric and ferulic acids in soils. Nature. 202 : 417-418.
- Whittaker, R.H. (1970). The Biochemical Ecology of Higher Plants. In Chemical Ecology. Sondheimer, E. and J. B. Simeone (eds.). Academic Press. New York, London. pp. 43-70.
- Whittaker, R.H. and P. P. Feeny. (1971). Allelochemicals : Chemical Interactions between Species. Science. 171 : 757-770.
- Wilson, R.E. and E.L. Rice. (1968). Allelopathy as expressed by Helianthus annuus and its role in old-field succession Bull. Torrey Bot. Club. 95 : 432-448.
- Winter, A.G. (1961). New physiological and biological aspects in the interrelationships between higher plants. In Mechanisms in biological competition. Soc. Exp. Biol. Symp. Cambridge Univ. Press, Cambridge. pp. 229-244.
- Woods, F.W. (1960). Biological antagonisms due to phytotoxic root exudates. Bot. Rev. 26 : 546-569.
- Wright, J.M. (1951). Phytotoxic effects of some antibiotics. Ann. Bot. 15 : 493-499.
- Wu, Yingdong and S. Guo. (1988). Karyotype-analysis of Cassia tora L. a traditional chinese medicine. J. Xi'an Med. Univ. 9 (1) : 21-23.