

List of Tables

Table 1.1: Characteristics of different varieties of Fig	25
Table 2.1 : Some commonly used plant hormones and their molecular weights.	32
Table 2.2 : Standard Basal Media Formulations of MS, Gamborg B5 and WPM medium	33
Table 3.1: Morphogenesis response of different ex-plants, sprouting and percentage survival rate after (15 days) interval of <i>Citrus aurantifolia</i> . (Each experiment has 3 replicates)	50
Table 3.2: Effect of BAP and NAA on shoot formation from axillary buds after 8 weeks.....	51
Table 3.3 : Effect of BAP and GA on shoot multiplication after 8 weeks in MS Medium	52
Table 3.4 : Effect of BAP, NAA and GA on shoot multiplication after 8 weeks in MS Medium	53
Table 3.5 : Effect of different concentration of IBA on rooting with combination of BAP (0.1mg/l) of in-vitro regenerated <i>Citrus aurantifolia</i> roots in ½ MS medium.	54
Table 4.1: Effect of individual cytokinin on shoot bud initiation from axillary bud explants after 4 weeks in MS Medium	67
Table 4.2: Effect of individual cytokinin on shoot bud initiation from axillary bud explants after 4 weeks in WPM Medium	68
Table 4.3 : Morphogenesis response of sprouting and percentage survival rate of <i>Ficus carica</i> grown MS media supplemented with different concentration of BAP with 0.1mg/l NAA. (Values determined after four weeks from three individual experiments).....	69
Table 4.4 : Effect of TDZ and IBA on shoot formation from axillary bud explants after 4 weeks in MS Medium	71
Table 4.5 : Effect of BAP and GA on shoot multiplication from axillary bud explants after 8 weeks in MS Medium.	71
Table 4.6 : Effect of BAP, NAA and GA on shoot multiplication from axillary bud explants after 8 weeks in MS Medium.....	72
Table 4.7: Effect of Basal Salt Concentration on <i>in vitro</i> plants.....	73
Table 4.8 : Effect of Activated Charcoal in full strength MS medium.....	74
Table 4.9: Effect of Auxins (IAA, IBA, NAA) in full strength MS medium for root induction.	75
Table 4.10: Use of different substrates for Primary Hardening Experiments:	76
Table 5.1 : General information and physical parameter of soil samples which were collected from Lemon and Fig tree roots.	94
Table 5.2 Percentage moisture in collected soil samples (20 samples from lemon and fig cultivation area).....	95
Table 5.3 Field data of Lemon plantation	95

Table 5.4 : Field data of Fig plantation.....	96
Table 5.5 : Isolation of AM spores in 100 gm soil sample (number of spores) of Lemon orchards.	101
Table 5.6 : Number of spores per 100 gm soil from Fig orchards.....	101
Table 6.1 : Optimized media gelling obtained on combination of Agar 3gm+ Rice flour 50gm (Plate 6.7, A-D).....	118
Table 6.2 : Optimized media gelling obtained on combination of Agar 4gm+ Isubgol 25gm (Plate 6.6, A-D).	118
Table 6.3 : Optimized media gelling obtained on combination of Agar 4gm+ Natural gum 10gm(Plate 6.4, A-D).....	119
Table 6.4 : Optimized media gelling obtained on combination of Agar 4gm+ Sabudana 10gm (Plate 6.5, A-D).....	119
Table 6.5 : Optimized media gelling obtained on combination of Agar 4gm+ Suji 10gm.....	120
Table 6.6 : Optimized media gelling obtained on combination of Agar 3gm+ Guar gum 25gm.(Plate 6.1, A-D).....	120
Table 6.7 : Optimized media gelling obtained on combination of Agar 4gm+ Singoda flour 30gm (Plate 6.2, A-D).....	120
Table 6.8 : Summary of above experiments (best gelling combination) and its costing.	121
Table 6.9 : Observations	121