

REFERENCES

Chapter V

Abd El-Moaty, H. I. (2015). Primary Metabolites and Saponin Constituents of *Taverniera lappacea* (Forssk.) DC. *Current Science International*, 4(4), 677-683.

Abe, N., Ebina, T., & Ishida, N. (1982). Interferon induction by glycyrrhizin and glycyrrhetic acid in mice. *Microbiology and immunology*, 26(6), 535–539. <https://doi.org/10.1111/j.1348-0421.1982.tb00207.x>.

Abed, A. K. M. (2007). Study of Amino Acids and Fatty Acids in Date Palm Fruit Cultivars Am Aldehin and Braime of Three Male Date Palm Pollinators. *Journal of Basrah Researches (Sciences)*, 33(3B).

Aboobucker, S.I., Suza, W.P. (2019). Why do plants convert sitosterol to stigmasterol? *Front. Plant Sci.*, 10, 354.

Adenan, M., S. T. Muhammad, T., B. Chung, C., A. Aizuddin, M., & Zahari, Z. (2022). Diisooctyl Phthalate Isolated from *Rourea mimosoides*: A Partial PPAR γ Agonist Potently Blocks Adipocyte Differentiation. *Tropical Journal of Natural Product Research (TJNPR)*, 6(8), 1197–1202.

Adjé, F., Lozano, Y. F., Lozano, P., Adima, A., Chemat, F., & Gaydou, E. M. (2010). Optimization of anthocyanin, flavonol and phenolic acid extractions from *Delonix regia* tree flowers using ultrasound-assisted water extraction. *Industrial Crops and Products*, 32(3), 439-444.

Afendi, F. M., Okada, T., Yamazaki, M., Hirai-Morita, A., Nakamura, Y., & Nakamura, K. (2012). KNApSACk family databases: integrated metabolite-plant species databases for multifaceted plant research. *Plant Cell Physiol.* 53: e1. [10.1093/pcp/pcr165](https://doi.org/10.1093/pcp/pcr165)

Agati, G., Azzarello, E., Pollastri, S., & Tattini, M. (2012). Flavonoids as antioxidants in plants: location and functional significance. *Plant science*, 196, 67-76.

Ahmad, A., Alkarkhi, A. F. M., Hena, S., Siddique, B. M., & Dur, K. W. (2010). Optimization of Soxhlet Extraction of *Herba Leonuri* Using Factorial Design of Experiment. *International Journal of Chemistry*, 2(1), 198–205. <https://doi.org/10.5539/ijc.v2n1p198>

Ajanal, M., Gundkalle, M. B., & Nayak, S. U. (2012). Estimation of total alkaloid in *Chitrakadivati* by UV-Spectrophotometer. *Ancient science of life*, 31(4), 198.

- Ajanal, M., Gundkalle, M. B., & Nayak, S. U. (2012). Estimation of total alkaloid in Chitrakadivati by UV-Spectrophotometer. *Ancient science of life*, 31(4), 198.
- Alam, Q., Vijayanarayana, K., & Satyanarayana, D. (2012). Evaluation of estrogenic activity of alcoholic extract of fruits of *Solanum xanthocarpum* using uterine wet weight, uterine glycogen content and uterine histology as parameter of assessment. *Int J Pharm Biol Sci*, 1:99-108.
- Albert, S., Horbach, R., Deising, H. B., Siewert, B., & Csuk, R. (2011). Synthesis and antimicrobial activity of (E) stilbene derivatives. *Bioorganic & medicinal chemistry*, 19(17), 5155-5166.
- Alcazar, A., Ballesteros, O., Jurado, J. M., Pablos, F., Martin, M. J., Vilches, J. L., & Navalón, A. (2007). Differentiation of green, white, black, Oolong, and Pu-erh teas according to their free amino acids content. *Journal of agricultural and food chemistry*, 55(15), 5960-5965.
- Alexiades, M. N., & Laird, S. A. (2002). Laying the foundation: equitable biodiversity research relationships. *Biodiversity and traditional knowledge: equitable partnerships in practice*. London, UK: Earthscan, 3-15.
- Ali, SI. (1977). Flora of West Pakistan, University of Karachi, 324.
- Al-Rifai, A. A., Aqel, A., Awaad, A., & Alothman, Z. A. (2015). Analysis of Quercetin and Kaempferol in an Alcoholic Extract of *Convolvulus pilosellifolius* using HPLC. *Communications in Soil Science and Plant Analysis*, 46(11), 1411-1418.
- Altemimi, A., Lightfoot, D. A., Kinsel, M., & Watson, D. G. (2015). Employing response surface methodology for the optimization of ultrasound assisted extraction of lutein and β -carotene from spinach. *Molecules*, 20(4), 6611–6625. <https://doi.org/10.3390/molecules20046611>
- Amit, N., & Daniel, M. (2014). Development of quality standards of *taverniera cuneifolia* (Roth) arn. root - A substitute drug for liquorice. *International Journal of Pharmacognosy and Phytochemical Research*, 6, 255–259.
- Ang, L. F., Yam, M. F., Fung, Y. T. T., Kiang, P. K., & Darwin, Y. (2014). HPLC method for simultaneous quantitative detection of quercetin and curcuminoids in traditional chinese medicines. *Journal of pharmacopuncture*, 17(4), 36.

- Angiosperm Phylogeny Group, Chase, M. W., Christenhusz, M. J., Fay, M. F., Byng, J. W., Judd, W. S., & Stevens, P. F. (2016). An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical journal of the Linnean Society*, 181(1), 1-20.
- Anil Kumar, M., Sravanthi Pammi, S. S., Sukanya, M. S., & Giri, A. (2018). Enhanced production of pharmaceutically important isoflavones from hairy root rhizoclones of *Trifolium pratense* L. *In Vitro Cellular & Developmental Biology-Plant*, 54(1), 94-103.
- Aniszewski, T. (2007). *Alkaloids-Secrets of Life: Alkaloid Chemistry, Biological Significance, Applications and Ecological Role*. Elsevier.
- Ansari, A., Patil, D., & Patil, A. (2015). HPTLC chemo-profiling of raw materials and formulations of Safoof-E-Bar: A unani formulation for vitiligo. *European Journal of Pharmaceutical and Medical Research*, 2(3): 575-588.
- AOAC (2019). Guidelines for Dietary Supplements and Botanicals (AOAC Official Method of Analysis, Appendix K). Available at: http://www.eoma.aoc.org/app_k.pdf (Accessed November 1, 2021).
- APG, IV. (2016). An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* 181: 1–20.
- Arase, Y., Ikeda, K., Murashima, N., Chayama, K., Tsubota, A., Koida, I., ... Kumada, H. (1997). The long term efficacy of glycyrrhizin in chronic hepatitis C patients. *Cancer*, 79(8), 1494–1500. [https://doi.org/10.1002/\(SICI\)1097-0142\(19970415\)79:8<1494::AID-CNCR8>3.0.CO;2-B](https://doi.org/10.1002/(SICI)1097-0142(19970415)79:8<1494::AID-CNCR8>3.0.CO;2-B)
- Archana, V., Kushagra, D. S. D., & Hemant, J. (2011). 4. Comparative Standardization Parameters of *Ficus Carica* Linn. And *Ficus Religiosa* Linn. By Vajpayee Archana¹, Dwivedi Sumeet, Dubey Kushagra and Joshi Hemant. *Life Sciences Leaflets*, 12, 376-383.
- Arimura, G. I., Ozawa, R., Shimoda, T., Nishioka, T., Boland, W., & Takabayashi, J. (2000). Herbivory-induced volatiles elicit defence genes in lima bean leaves. *Nature*, 406(6795), 512-515.

- Arnqvist, L., Persson, M., Jonsson, L., Dutta, P.C., & Sitbon, F. (2007). Overexpression of *CYP710A1* and *CYP710A4* in transgenic *Arabidopsis* plants increases the level of stigmasterol at the expense of sitosterol. *Planta*, 227, 309–317.
- Ashraf, M. F. M. R., & Foolad, M. R. (2007). Roles of glycine betaine and proline in improving plant abiotic stress resistance. *Environmental and experimental botany*, 59(2), 206-216.
- Atanasov, A. G., Waltenberger, B., Pferschy-Wenzig, E. M., Linder, T., Wawrosch, C., Uhrin, P., ... & Stuppner, H. (2015). Discovery and resupply of pharmacologically active plant-derived natural products: A review. *Biotechnology advances*, 33(8), 1582-1614.
- Aucamp, J. P., Hara, Y., & Apostolides, Z. (2000). Simultaneous analysis of tea catechins, caffeine, gallic acid, theanine and ascorbic acid by micellar electrokinetic capillary chromatography. *Journal of Chromatography A*, 876(1-2), 235-242.
- Audru, J. (1987). Maisons-Alfort: IEMVPT. La vegetation. Republique de Djibouti.
- Ayres, D.C., Loike, J. D. L. (1991). Chemical, Biological and Clinical Properties. Lignans Chem. *Biol. Clin. Prop.*
- Aziz, M., & Karboune, S. (2018). Natural antimicrobial/antioxidant agents in meat and poultry products as well as fruits and vegetables: A review. *Critical reviews in food science and nutrition*, 58(3), 486-511.
- Babbar, N. (2015). An introduction to alkaloids and their applications in pharmaceutical chemistry. *The Pharma Innovation Journal*, 4(10), 74-75.
- Bailly, C., & Vergoten, G. (2020). Glycyrrhizin: An alternative drug for the treatment of COVID-19 infection and the associated respiratory syndrome. *Pharmacology & therapeutics*, 214, 107618.
- Bairiganjan, GC. (1985). Fabaceae in Orissa. *J. Econ. Taxon Bot.*, 7:249-276.
- Bansal, A., Chhabra, V., Rawal, R.K., & Sharma, S. (2013). Chemometrics: A new scenario in herbal drug standardization. *J. Pharm. Anal*, 4:223–233. doi: 10.1016/j.jpha.2013.12.001.
- Bansal, S., & DeStefano., A. (2007). Key elements of bioanalytical method validation for small molecules. *AAPS J*, 9: E109-14.
- Baragi, P. C., Patgiri, B. J., & Prajapati, P. K. (2008). Neutraceuticals in Ayurveda with special reference to Avaleha Kalpana. *Ancient Science of Life*, 28(2), 29.

- Barbehenn, R. V., & Constabel, C. P. (2011). Tannins in plant–herbivore interactions. *Phytochemistry*, 72(13), 1551-1565.
- Bardakci, H., Celep, E., Gözet, T., Kurt-Celep, I., Deniz, I., Şen-Utsukarci, B., & Akaydin, G. A. L. İ. P. (2019). A comparative investigation on phenolic composition, antioxidant and antimicrobial potentials of *Salvia heldreichiana* Boiss. ex Benth extracts. *South African Journal of Botany*, 125, 72-80.
- Basar et al. (2014). *Phytochem. Anal*, 25, 399–404. DOI 10.1002/pca.2507399
- Bassi, R., & Sharma, S. S. (1993). Changes in proline content accompanying the uptake of zinc and copper by *Lemna minor*. *Annals of botany*, 72(2), 151-154.
- Bassi, R., & Sharma, S. S. (1993). Proline accumulation in wheat seedlings exposed to zinc and copper. *Phytochemistry*, 33(6), 1339-1342.
- Beccaria, M., Inferred, V., Rigano, F., Gorynski, K., Purcaro, G., Pawliszyn, J., ... & Mondello, L. (2017). Highly informative multiclass profiling of lipids by ultra-high performance liquid chromatography–Low resolution (quadrupole) mass spectrometry by using electrospray ionization and atmospheric pressure chemical ionization interfaces. *Journal of Chromatography A*, 1509, 69-82.
- Benbelaid, F., Abdoune, MA., Khadir, A., & Bendahou M. (2013). Drying effect on yield and antimicrobial activity of essential oils. *Int J Med Arom Plants*, 3:93-101.
- Bennett, B. C., & Balick, M. J. (2008). Phytomedicine 101: plant taxonomy for preclinical and clinical medicinal plant researchers. *Journal of the Society for Integrative Oncology*, 6(4), 150.
- Bennett, B. C., & Balick, M. J. (2014). Does the name really matter? The importance of botanical nomenclature and plant taxonomy in biomedical research. *Journal of ethnopharmacology*, 152(3), 387-392.
- Bentham, G., & Hooker, JD. (1862–1883) *Genera Plantarum*, Vols. 1-3. L. Reeve & Co., London:1862–1883.
- Bernardo-Gil, M. G., Roque, R., Roseiro, L. B., Duarte, L. C., Gírio, F., & Esteves, P. (2011). Supercritical extraction of carob kibbles (*Ceratonia siliqua* L.). *Journal of Supercritical Fluids*, 59, 36–42. <https://doi.org/10.1016/j.supflu.2011.08.007>

- Beutler, J. A. (2019). Natural products as a foundation for drug discovery. *Current protocols in pharmacology*, 86(1), e67.
- Bhandari, MM. (1978). Flora of the Indian desert, Jodhpur.
- Bhardwaj, SK., Dwivedi, K., & Agarwal, DD. (2015) A Review: HPLC Method Development and Validation. *International Journal of Analytical and Bioanalytical Chemistry*, 5(4): 76-81. 4.
- Bhattacharya, A., Sood, P., & Citovsky, V. (2010). The roles of plant phenolics in defence and communication during *Agrobacterium* and *Rhizobium* infection. *Molecular plant pathology*, 11(5), 705-719.
- Bodini, S.F., Manfredini, S., Epp, M., Valentini, S., & Santori, F. (2009). Quorum sensing inhibition activity of garlic extract and *p*-coumaric acid. *Lett. Appl. Microbiol.*, 49:551–555.
- Boissier, E. (1888) Umbelliferae. In: Buser, R. (Ed.) *Flora Orientalis Supplementum*. H.Georg, Genève, Basel, pp. 251–272
- Borgesen, F. (1929). Notes on the vegetation of Dwarka on the west coast of India. *J. Ind. Bot. Soc*, 8, 1-18.
- Borman, P., & Elder, D. (2017). Q2 (R1) validation of analytical procedures. *ICH Quality guidelines*, 5, 127-166.
- Boudet, A. M. (2007). Evolution and current status of research in phenolic compounds. *Phytochemistry*, 68(22-24), 2722-2735.
- Bravi, E., Perretti, G., Montanari, L., Favati, F., & Fantozzi, P. (2007). Supercritical fluid extraction for quality control in beer industry. *Journal of Supercritical Fluids*, 42(3 SPEC. ISS.), 342–346. <https://doi.org/10.1016/j.supflu.2007.02.012>
- Brinckmann, J. A. (2020). The long road to sustainable licorice. *Windowsill to The World of Medicinal Plants*, 19.
- British Pharmacopoeia (2020). London: The Department of Health and Social Care, British Pharmacopoeia Commission Office.
- Bruneau, A., Doyle, J. J., Herendeen, P., Hughes, C., Kenicer, G., ... & van Wyk, B. E. (2013). Legume phylogeny and classification in the 21st century: Progress, prospects and lessons for other species-rich clades. *Taxon*, 62(2), 217-248.

- Bryant, J. Burrell, M.M.; Kruger, N. N. (1999). Plant Carbohydrate Biochemistry. In Garland Sci./Taylor & Francis: New York, NY, USA.
- Butler, M. S. (2004). The role of natural product chemistry in drug discovery. *Journal of natural products*, 67(12), 2141-2153.
- Cai, H., Steward, W. P., & Gescher, A. J. (2005). Determination of the putative cancer chemopreventive flavone tricetin in plasma and tissues of mice by HPLC with UV–visible detection. *Biomedical Chromatography*, 19(7), 518-522.
- Cal, K., Kupiec, K., & Sznitowska, M. (2006). Effect of physicochemical properties of cyclic terpenes on their ex vivo skin absorption and elimination kinetics. *Journal of Dermatological Science*, 41(2), 137-142.
- Calapai, G. (2008). European legislation on herbal medicines: a look into the future. *Drug Saf.* 31, 428–431. doi: 10.2165/00002018-200831050-00009
- Canonica, L., Russo, G., & Bombardelli, E. (1996). Triterpenes of *Glycyrrhiza glabra*. 2. Liquiritic acid. *Gazz Chim Ital.*, 96:833–842.
- Cao, Y., Tang, X. Y., & Song, F. W. (2002). Compare of and beta-Glycyrrhetic acid alpha-Glycyrrhetic acid in the treatment of Chronic Hepatitis B. *China Journal of Modern Medicine*, 12(7), 91-92.
- Capilla, A. S., Sánchez, I., Caignard, D. H., Renard, P., & Pujol, M. D. (2001). Antitumor agents. Synthesis and biological evaluation of new compounds related to podophyllotoxin, containing the 2,3-dihydro-1,4-benzodioxin system. *European Journal of Medicinal Chemistry*, 36(4), 389–393. [https://doi.org/10.1016/S0223-5234\(01\)01231-4](https://doi.org/10.1016/S0223-5234(01)01231-4)
- Chaiyasit, W., Elias, R. J., McClements, D. J., & Decker, E. A. (2007). Role of physical structures in bulk oils on lipid oxidation. *Critical reviews in food science and nutrition*, 47(3), 299-317.
- Chalal, M., Klinguer, A., Echairi, A., Meunier, P., Vervandier-Fasseur, D., & Adrian, M. (2014). Antimicrobial activity of resveratrol analogues. *Molecules*, 19(6), 7679-7688.
- Charles, R., Garg, S. N., & Kumar, S. (2000). New gingerdione from the rhizomes of *Zingiber officinale*. *Fitoterapia*, 71(6), 716-718.

- Charlton, J. L. (1998). Antiviral activity of lignans. *Journal of Natural Products*, 61(11), 1447–1451. <https://doi.org/10.1021/np980136z>
- Chaudhari, V. S., Borkar, R. M., Murty, U. S., & Banerjee, S. (2020). Analytical method development and validation of reverse-phase high-performance liquid chromatography (RP-HPLC) method for simultaneous quantifications of quercetin and piperine in dual-drug loaded nanostructured lipid carriers. *Journal of Pharmaceutical and Biomedical Analysis*, 186, 113325.
- Chauhan, BP., Uniyal, DK., & Singh. (2003). *Taverniera cuneifolia* (Roth) Arn. (Fabaceae): A New Record for Jammu & Kashmir, Northwest Himalaya. *Annals of Forestry*, 11(98):10.
- Chauhan, S., Sharma, A., Upadhyay, N. K., Singh, G., Lal, U. R., & Goyal, R. (2018). In-vitro osteoblast proliferation and in-vivo anti-osteoporotic activity of *Bombax ceiba* with quantification of Lupeol, gallic acid and β -sitosterol by HPTLC and HPLC. *BMC complementary and alternative medicine*, 18, 1-12.
- Chen, L., Chen, Q., Zhang, Z., & Wan, X. (2009). A novel colorimetric determination of free amino acids content in tea infusions with 2, 4-dinitrofluorobenzene. *Journal of Food Composition and Analysis*, 22(2), 137-141.
- Chen, Y., Xie, M. Y., & Gong, X. F. (2007). Microwave-assisted extraction used for the isolation of total triterpenoid saponins from *Ganoderma atrum*. *Journal of Food Engineering*, 81(1), 162–170. <https://doi.org/10.1016/j.jfoodeng.2006.10.018>
- Chesnin, L., & Yein, CH. (1950). Turbidimetric determination of available sulphates. *Soil Sci. Soc. Am. Proc.*, 15: 149-151.
- Chirinos, R., Betalleluz-Pallardel, I., Huamán, A., Arbizu, C., Pedreschi, R., & Campos, D. (2009). HPLC-DAD characterisation of phenolic compounds from Andean oca (*Oxalis tuberosa* Mol.) tubers and their contribution to the antioxidant capacity. *Food Chemistry*, 113(4), 1243–1251. <https://doi.org/10.1016/J.FOODCHEM.2008.08.015>
- Cho, H. J., Lim, S. S., Lee, Y. S., Kim, J. S., Lee, C. H., Kwon, D. Y., & Park, J. H. Y. (2010). Hexane/ethanol extract of *Glycyrrhiza uralensis* licorice exerts potent anti-inflammatory effects in murine macrophages and in mouse skin. *Food chemistry*, 121(4), 959-966.
- Chong, J., Poutaraud, A., & Hugueney, P. (2009). Metabolism and roles of stilbenes in plants. *Plant science*, 177(3), 143-155. (CrossRef)

- Cinatl, J., Morgenstern, B., Bauer, G., Chandra, P., Rabenau, H., & Doerr, H. W. (2003). Glycyrrhizin, an active component of liquorice roots, and replication of SARS-associated coronavirus. *Lancet*, 361(9374), 2045–2046. [https://doi.org/10.1016/S0140-6736\(03\)13615-X](https://doi.org/10.1016/S0140-6736(03)13615-X)
- Clifford, M. N. (1999). Chlorogenic acids and other cinnamates - Nature, occurrence and dietary burden. *Journal of the Science of Food and Agriculture*, 79(3), 362–372. [https://doi.org/10.1002/\(SICI\)1097-0010\(19990301\)79:3<362: AID-JSFA256>3.0.CO;2-D](https://doi.org/10.1002/(SICI)1097-0010(19990301)79:3<362: AID-JSFA256>3.0.CO;2-D)
- Clifford, M. N., & Scalbert, A. (2000). Ellagitannins–nature, occurrence and dietary burden. *Journal of the Science of Food and Agriculture*, 80(7), 1118-1125.
- Combs, C. A. (2016). Tannins: biochemistry, food sources and nutritional properties. Nova Publishers.
- Committee, F. of C.E., Others. (2018). Flora of China. Flora of China.
- Cook, N. C., & Samman, S. (1996). Flavonoids—chemistry, metabolism, cardioprotective effects, and dietary sources. *The Journal of nutritional biochemistry*, 7(2), 66-76.
- Cooke, T. (1901-1908). The Flora of the presidency of Bombay. London. Vols. 1, 2 & 3 (Repr. eds. 1958. Vols. 1 & 2.BSI, Calcutta).
- Cooke, T. (1908). The flora of the presidency of Bombay (Vol. 2). Bishen Singh Mahendra Pal Singh.
- Corson, T. W., & Crews, C. M. (2007). Molecular understanding and modern application of traditional medicines: triumphs and trials. *Cell*, 130(5), 769-774.
- Cosa, P., Vlietinck, A. J., Berghe, D. V., & Maes, L. (2006). Anti-infective potential of natural products: How to develop a stronger in vitro ‘proof-of-concept’. *J. Ethnopharmacol.*, 106, 290–302.
- Coskun, O. (2016). Separation techniques: chromatography. *Northern clinics of Istanbul*, 3(2), 156.
- Cotton, C. M. (1996). *Ethnobotany: principles and applications*. John Wiley & Sons.
- Cragg, G. M., & Newman, D. J. (2013). Natural products: a continuing source of novel drug leads. *Biochimica et Biophysica Acta (BBA)-General Subjects*, 1830(6), 3670-3695.

- Cronquist, A. (1968). The evolution and classification of flowering plants. *The evolution and classification of flowering plants*.
- Cushnie, T. T., Cushnie, B., & Lamb, A. J. (2014). Alkaloids: An overview of their antibacterial, antibiotic-enhancing and antivirulence activities. *International journal of antimicrobial agents*, 44(5), 377-386.
- Da Porto, C., & Decorti, D. (2009). Ultrasound-assisted extraction coupled with under vacuum distillation of flavour compounds from spearmint (carvone-rich) plants: comparison with conventional hydrodistillation. *Ultrasonics sonochemistry*, 16(6), 795-799.
- Daffodil, E. D., Uthayakumari, F. K., & Mohan, V. R. (2012). GC-MS determination of bioactive compounds of *Curculigo orchoides gaertn.* *Science Research Reporter*, 2(3), 198-201.
- Dagne, E., Yenesew, A., Capasso, F., Mascolo, N., & Pinto, A. (1990). Preliminary studies on antipyretic and analgesic properties of *Taverniera abyssinica*. *Ethiopian medical journal*, 28(4), 155-161.
- Dalzell, N. A. (1850). *Phaseolus*. *Hooker's J Bot Kew Gard Misc*, 2, 33-37.
- Dash, S., Das, C., & Sahoo, D. C. (2010). Phytochemical and anthelmintic screening of crude bark extract of *Adenanthera pavonina* Linn. *International Journal of Comprehensive Pharmacy*, 2, 1-4.
- De Bruyne, T., Pieters, L., Deelstra, H., & Vlietinck, A. (1999). Condensed vegetable tannins: biodiversity in structure and biological activities. *Biochemical Systematics and Ecology*, 27(4), 445-459.
- De Meulder, M., Waldron, M. P., Li, L., Peay, M. G., Tingler, M. J., Hidy, B. J., ... & Jenkins, R. G. (2016). Development and validation of HILIC–ESI/MS/MS methods for simultaneous quantitation of several antipsychotics in human plasma and blood. *Bioanalysis*, 8(8), 765-794.
- De, B., Debbarma, T., Sen, S., & Chakraborty, R. (2010). Tribal life in the environment and biodiversity of Tripura, India. *Current World Environment*, 5(1), 59.
- Denisova, S. B., Danilov, V. T., Yunusova, S. G., Davydova, V. A., Murinov, Y. I., & Zarudii, F. S. (2007). Isolation and biological activity of lipids from licorice (*Glycyrrhiza glabra*) roots. *Pharmaceutical Chemistry Journal*, 41, 489-491.

- Desai, S., Tatke, P., Mane, T., & Gabhe, S. (2020). Isolation, characterization and quantitative HPLC-DAD analysis of components of charantin from fruits of *Momordica charantia*, *Food Chemistry*, doi: <https://doi.org/10.1016/j.foodchem.2020.128717>
- Desire, DDP., Sandrine, MNY., Claude, BD., Mireille, K., Bibi-Farouck, AO., & Theophile, D. (2015). In vivo estrogenic-like activities of *Gouania longipetala* Hemsl. (Rhamnaceae) bark extracts in a post-menopause-like model of ovariectomized Wistar rats. *J Ethnopharmacol*, 168:22-8.
- Dev, S. (1988). Ayurveda and modern drug development. *Proceedings of the Indian National Science Academy A: Physical Sciences*, 54, 12-42.
- Dev, S. (1997). Ethnotherapeutics and modern drug development: the potential of Ayurveda. *Current science*, 73(11), 909-928.
- Devi, V. K., Jain, N., & Valli, K. S. (2010). Importance of novel drug delivery systems in herbal medicines. *Pharmacognosy reviews*, 4(7), 27.
- Dey, P., Kundu, A., Chakraborty, H. J., Kar, B., Choi, W. S., Lee, B. M., ... & Kim, H. S. (2019). Therapeutic value of steroidal alkaloids in cancer: Current trends and future perspectives. *International journal of cancer*, 145(7), 1731-1744.
- Dhiman, A., Sharma, K., Sharma, A., & Sindhu, P. (2016). A review on the status of quality control and standardization of herbal drugs in India. *Drug Development and Therapeutics*, 7(2), 107-107.
- Ding, Y., Yu, H., & Mou, S. (2002). Direct determination of free amino acids and sugars in green tea by anion-exchange chromatography with integrated pulsed amperometric detection. *Journal of Chromatography A*, 982(2), 237-244.
- Dixon, R. A., Dey, P. M., & Lamb, C. J. (1983). Phytoalexins: enzymology and molecular biology. *Advances in enzymology and related areas of molecular biology*, 55(1), 69.
- Djilan, A., Legseir, B., Soulimani, R., Dicko, A., & Younos, C. (2006). New extraction technique for alkaloids. *Journal of the Brazilian Chemical Society*, 17, 518-520.
- Dowell, A., Davidson, G., & Ghosh, D. (2015). Validation of quantitative HPLC method for Bacosides in Keen Mind. *J Evid Based Complementary Altern Med*, 2105:1-8.

- Du, G., Zhao, H., Song, Y., Zhang, Q., & Wang, Y. (2011). Rapid simultaneous determination of isoflavones in *Radix puerariae* using high-performance liquid chromatography–triple quadrupole mass spectrometry with novel shell-type column. *Journal of separation science*, 34(19), 2576-2585.
- Dubey, N. K., Kumar, R., & Tripathi, P. (2004). Global promotion of herbal medicine: India's opportunity. *Current science*, 86(1), 37-41.
- Dubrovina, A. S., & Kiselev, K. V. (2017). Regulation of stilbene biosynthesis in plants. *Planta*, 246(4), 597-623. (CrossRef)
- Duddeck, H., Yenesew, A., & Dagne, E. (1987). Isoflavonoids from *Taverniera abyssinica*. *Bulletin of the Chemical Society of Ethiopia*, 1(1).
- Durling, N. E., Catchpole, O. J., Grey, J. B., Webby, R. F., Mitchell, K. A., Foo, L. Y., & Perry, N. B. (2007). Extraction of phenolics and essential oil from dried sage (*Salvia officinalis*) using ethanol-water mixtures. *Food Chemistry*, 101(4), 1417–1424. <https://doi.org/10.1016/j.foodchem.2006.03.050>
- Eagleson, M. (1994). Concise Encyclopedia Chemistry (1st ed.). Edn. Berlin: Walter deGruyter;
- Ebere, E. C., Obinna, I. B., & Wirnkör, V. A. (2019). Applications of column, paper, thin layer and ion exchange chromatography in purifying samples: Mini review. *SF Journal of Pharmaceutical and Analytical Chemistry*.
- Ehlers, D., Nguyen, T., Quirin, K., and Gerard, D. (2001). Analysis of essential basil oils-CO₂ extracts and steam-distilled oils. *Deutsche Lebensmittel-Rundschau*, 97(7), 245–250.
- El Khawand, T., Courtois, A., Valls, J., Richard, T., & Krisa, S. (2018). A review of dietary stilbenes: Sources and bioavailability. *Phytochemistry Reviews*, 17(5), 1007-1029. (CrossRef)
- Elachouri, M. (2018). Ethnobotany/ethnopharmacology, and bioprospecting: Issues on knowledge and uses of medicinal plants by Moroccan people. In *natural products and drug discovery* (pp. 105-118). Elsevier.
- Eller, F. (2002). Supercritical carbon dioxide extraction of fat. *Bus. Brief. Food Technol.*, 1–4.

El-Moaty, H. I. A. (2016). Antimicrobial activity of *Taverniera lappacea* forssk. extract affected by its active constituents. *Indo American journal of pharmaceutical sciences*, 3(10), 1092-1095.

EMA Guideline on Quality of Herbal Medicinal Products/Traditional Herbal Medicinal Products. 2011. [(accessed on 28 February 2022)]. Available online: https://www.ema.europa.eu/en/documents/scientific-guideline/guideline-quality-herbal-medicinal-products-traditional-herbal-medicinal-products-revision-2_en.pdf

Ende, W. V. D., & Peshev, D. (2013). Sugars as antioxidants in plants. In *Crop improvement under adverse conditions* (pp. 285-307). Springer, New York, NY.

Ernst, B.; Hart, G.W.; Sinay, P. (2000). Carbohydrates in Chemistry and Biology. *John Wiley & Sons: Weinheim, Germany*.

Ettre, L. S. (1979). 75 years of chromatography: A glimpse behind the scenes. *Journal of High Resolution Chromatography*, 2(8), 500-506.

Falodun, A., Okunrobe, L.O., & Uzoamaka, N. (2006). Phytochemical screening and anti-inflammatory evaluation of methanolic and aqueous extracts of *Euphorbia heterophylla* Linn9Euphorbiaceae). *Afr. J. Biotechnol*, 5(6):529-531.

Farag, M. A., & Wessjohann, L. A. (2012). Volatiles profiling in medicinal Licorice roots using steam distillation and solid-phase microextraction (SPME) coupled to chemometrics. *Journal of Food Science*, 77(11), C1179-C1184.

Farhan, N., Fitzpatrick, S., Shim, Y. M., Paige, M., & Chow, D. S. L. (2016). Ultrapressure liquid chromatography–tandem mass spectrometry assay using atmospheric pressure photoionization (UPLC-APPI-MS/MS) for quantification of 4-methoxydiphenylmethane in pharmacokinetic evaluation. *Journal of Pharmaceutical and Biomedical Analysis*, 128, 46-52.

Farmakope Herbal Indonesia (2017). Edisi II. Jakarta: Kementrian Kesehatan Republik Indonesia.

Farooqui, NA., Dey, A., Singh, GN., & Easwari, TS. (2015). Analytical method development and validation for the quantitative estimation of nimbin in bulk and solid dosage form by RP-HPLC. *Int J Pharm Bio Sci*, 6:479-90.

FDA (2016). Botanical Drug Development, Guidance for Industry, U.S. Department of Health and Human Services Food and Drug Administration Center for Drug Evaluation and Research

- (CDER). Available at: <https://www.fda.gov/media/93113/download> (Accessed August 26, 2021).
- Feng, G., Runyi, Y. (2006). New approach on similarity analysis of chromatographic fingerprint of herbal medicine. *Journal of Chromatography A*, 1104:100-5.
- Fenner, M. A., Chakrabarty, S., Wang, B., Pagnotti, V. S., Hoang, K., Trimpin, S., & McEwen, C. N. (2017). An LC/MS method providing improved sensitivity: electrospray ionization inlet. *Analytical chemistry*, 89(9), 4798-4802.
- Fich, E. A., Segerson, N. A., & Rose, J. K. (2016). The plant polyester cutin: biosynthesis, structure, and biological roles. *Annual review of plant biology*, 207-233. (crossref)
- Filgueira, D., Moldes, D., Fuentealba, C., & García, D. E. (2017). Condensed tannins from pine bark: A novel wood surface modifier assisted by laccase. *Industrial Crops and Products*, 103, 185-194.
- Fischer, E. Ber. (1890). The monosaccharides. (Vol. 23). Stanek, J.; Cerny, M.; Kocouvek, J.; Pacak, J.; Academic Press: New York.
- Frattini, C., Bicchi, C., Baretini, C., & Nano, G. M. (1977). Volatile flavor components of licorice. *Journal of Agricultural and Food Chemistry*, 25(6), 1238-1241.
- Fukai, T., Cai, B., Maruno, K., Miyakawa, Y., Konishi, M., & Nomura, T. (1998). An isoprenylated flavanone from *Glycyrrhiza glabra* and rec-assay of licorice phenols. *Phytochemistry*, 49, 2005–2013.
- Fukui, H., GOTO, K., & TABATA, M. (1988). Two antimicrobial flavanones from the leaves of *Glycyrrhiza glabra*. *Chemical and pharmaceutical bulletin*, 36(10), 4174-4176.
- Gad, HA., El-Ahmady, SH., Abou-Shoer, MI., & Al-Azizi. (2013). Application of chemometrics in authentication of herbal medicines: A review. *Phytochem Anal*, 24:1-24.
- Gadepalli, S. G., Deme, P., Kuncha, M., & Sistla, R. (2014). Simultaneous determination of amlodipine, valsartan and hydrochlorothiazide by LC–ESI-MS/MS and its application to pharmacokinetics in rats. *Journal of pharmaceutical analysis*, 4(6), 399-406.
- Gaggeri, R., Rossi, D., Collina, S., Mannucci, B., Baierl, M., & Juza, M. (2011). Quick development of an analytical enantioselective high performance liquid chromatography

- separation and preparative scale-up for the flavonoid Naringenin. *Journal of Chromatography A*, 1218(32), 5414-5422.
- Gamble, JS. (1918). Flora of Presidency of Madras, The Authority of the Secretary of state for India in council, (1):327.
- Garg, V., Dhar, V. J., Sharma, A., & Dutt, R. (2012). Facts about standardization of herbal medicine: a review. *Zhong xi yi jie he xue bao= Journal of Chinese Integrative Medicine*, 10(10), 1077-1083.
- Geng, Y., Liu, J., Lv, R., Yuan, J., Lin, Y., & Wang, X. (2007). An efficient method for extraction, separation and purification of eugenol from *Eugenia caryophyllata* by supercritical fluid extraction and high-speed counter-current chromatography. *Separation and Purification Technology*, 57(2), 237–241. <https://doi.org/10.1016/j.seppur.2007.04.015>
- Genovese, T., Menegazzi, M., Mazzon, E., Crisafulli, C., Di Paola, R., Dal Bosco, M., ... Cuzzocrea, S. (2009). Glycyrrhizin reduces secondary inflammatory process after spinal cord compression injury in mice. *Shock*, 31(4), 367–375. <https://doi.org/10.1097/SHK.0b013e3181833b08M>.
- Gershenzon, J., & Dudareva, N. (2007). The function of terpene natural products in the natural world. *Nature Chemical Biology*, 3(7), 408–414. <https://doi.org/10.1038/nchembio.2007.5>
- Ghasemi, M. (2019). High-performance thin-layer chromatography (HPTLC) method for estimation of Glycyrrhizic acid (GA) in an herbal extract of *Taverneira cuneifolia*. *International journal of recent scientific research*, 10(12), 36313-36315.
- Ghosh, AB., Bajaj, JC., Hasan, R., & Singh, D. (1983). Soil and water Testing Methods- A Laboratory Manual. *Division of soil science and Agricultural chemistry*, IARI, New Delhi.
- Giovando, S., Koch, G., Romagnoli, M., Paul, D., Vinciguerra, V., Tamantini, S., ... & Mugnozza, G. S. (2019). Spectro-topochemical investigation of the location of polyphenolic extractives (tannins) in chestnut wood structure and ultrastructure. *Industrial Crops and Products*, 141, 111767.
- Gohil, A. N., & Daniel, M. (2014). Development of Quality Standards of *Taverniera cuneifolia* (Roth) Arn. Root—A Substitute Drug for Licorice. *International Journal of Pharmacognosy and Phytochemical Research*, 6(02), 255-259.

- Goodarzi, M., Russell, P.J., & Vander Heyden, Y. (2013). Similarity analyses of chromatographic herbal fingerprints: A review. *Anal. Chim. Acta*, 804:16–28. doi: 10.1016/j.aca.2013.09.017
- Govt. of India. (2000). Medicinal Plants Introduction, Indian System of Medicine and Homoeopathy (ISMH), Department of ISMH, Ministry of Health and Family Welfare, Govt. of India. Available at: <http://indianmedicine.nic.in/html/plants/mimain.htm>
- Gray, A. (1886). Synoptical Flora of North America: The Gamopetalae, being a Second Edition of Vol I Part II, and Vol II Part I, Collected (Vol. 31). Ivison, Blakeman, Taylor.
- Grosso, C., Cardoso, M. a T., Figueiredo, a C., & Moldão-martins, M. (2007). Supercritical fluid extraction, Hydrodistillation and Soxhlet extraction of the aerial part of winter savory . Comparative evaluation of the extraction method on the chemical composition. *Proceedings of European Congress of Chemical Engineering*, September 16–20.
- Güçlü-Üstündağ, Ö., & Mazza, G. (2007). Saponins: properties, applications and processing. *Critical reviews in food science and nutrition*, 47(3), 231-258.
- Gulumian, M., Yahaya, E. S., & Steenkamp, V. (2018). African herbal remedies with antioxidant activity: a potential resource base for wound treatment. *Evidence-Based Complementary and Alternative Medicine*, 2018.
- Gumpricht, E., Dahl, R., Devereaux, M. W., & Sokol, R. J. (2005). Licorice compounds glycyrrhizin and 18 β -glycyrrhetic acid are potent modulators of bile acid-induced cytotoxicity in rat hepatocytes. *Journal of Biological Chemistry*, 280(11), 10556–10563. <https://doi.org/10.1074/jbc.M411673200>
- Gupta, D., Verma, M., Anand, R., & Khajuria, R. (2013). Development of a validated UPLC–qTOF-MS/MS method for determination of bioactive constituent from *Glycyrrhiza glabra*. *Journal of Pharmaceutical Analysis*, 3 (3). doi: 10.1016/j.jpha.2013.01.001
- Hagerman, A. E., Riedl, K. M., Jones, G. A., Sovik, K. N., Ritchard, N. T., Hartzfeld, P. W., & Riechel, T. L. (1998). High molecular weight plant polyphenolics (tannins) as biological antioxidants. *Journal of agricultural and food chemistry*, 46(5), 1887-1892.
- Hajimehdipoor, H., Amin, G. R., Adib, N., Rastegar, H., & Shekarchi, M. (2012). Development of a validated HPLC method for the simultaneous determination of flavonoids in *Cuscuta*

- chinensis Lam. by ultra-violet detection. *DARU Journal of Pharmaceutical Sciences*, 20(1), 1-6.
- Han, Y., Hou, Z., Zhang, X., Yan, K., Liang, Z., & He, Q. (2022). Important changes in germination, seedling tolerance, and active components content due to drought stress on three licorice (*Glycyrrhiza*) species. *Industrial Crops and Products*, 175, 114240.
- Hanai, T., & Hanai, T. T. (1999). *HPLC: a practical guide* (Vol. 6). Royal Society of Chemistry.
- Hanower, P., & Brzozowska, J. (1975). The effect of osmotic stress on the composition of free amino acids in cotton leaves. *Phytochemistry*.
- Hanuš, L. O., Řezanka, T., & Dembitsky, V. M. (2003). A trinorsesterterpene glycoside from the North American fern *Woodwardia virginica* (L.) Smith. *Phytochemistry*, 63(8), 869-875.
- Harborne, A. J. (1998). *Phytochemical methods a guide to modern techniques of plant analysis*. springer science & business media.
- Hare, P. D., Cress, W. A., & Van Staden, J. (1998). Dissecting the roles of osmolyte accumulation during stress. *Plant, cell & environment*, 21(6), 535-553.
- Hartmann, M.-A. (1998) Plant sterols and the membrane environment. *Trends Plant Sci.* 3, 170– 175.
- Haslam, E. (1982). The metabolism of gallic acid and hexahydroxydiphenic acid in higher plants. In *Fortschritte der Chemie organischer Naturstoffe/Progress in the Chemistry of Organic Natural Products* (pp. 1-46). Springer, Vienna.
- Haslam, E. (1989). Plant Polyphenols – Vegetable Tannins Revisited – Chemistry and Pharmacology of Natural Products. In Cambridge University Press, Cambridge.
- Hassan, A. R., Amer, K. F., El-Toumy, S. A., Nielsen, J., & Christensen, S. B. (2019). A new flavonol glycoside and other flavonoids from the aerial parts of *Taverniera aegyptiaca*. *Natural product research*, 33(8), 1135-1139.
- Hassanean, H. A. (1998). Saikosaponins from *Taverniera aegyptiaca*. *Phytochemistry*, 48(4), 693-697.
- Hayashi, H., & Sudo, H. (2009). Economic importance of licorice. *Plant Biotechnology*, 26(1), 101-104.

- Hayashi, H., Fukui, H., & Tabata, M. (1988). Examination of triterpenoids produced by callus and cell suspension cultures of *Glycyrrhiza glabra*. *Plant Cell Rep.*, 7, 508–511.
- Hayat, S., Hayat, Q., Alyemeni, M. N., Wani, A. S., Pichtel, J., & Ahmad, A. (2012). Role of proline under changing environments: a review. *Plant signaling & behavior*, 7(11), 1456-1466.
- Heim, K. E., Tagliaferro, A. R., & Bobilya, D. J. (2002). Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *The Journal of nutritional biochemistry*, 13(10), 572-584.
- Henry, M. (2005). Saponins and phylogeny: example of the “gypsogenin group” saponins. *Phytochemistry Reviews*, 4(2), 89-94.
- Hethelyi, E., Tetenyi, P., Dabi, E., & Danos, B. (1987). The role of mass spectrometry in medicinal plant research. *Biomedical & environmental mass spectrometry*, 14(11), 627-632.
- Hillis, W. E., & Rozsa, A. N. (1985). High temperature and chemical effects on wood stability. *Wood Science and Technology*, 19(1), 57-66.
- Hirano, T., Wakasugi, A., Oohara, M., Oka, K., & Sashida, Y. (1991). Suppression of Mitogen-Induced Proliferation of Human Peripheral Blood Lymphocytes by Plant Lignans. *Planta Med*, 57(04), 331–334.
- Hoefer, G., Baltina, L., Michaelis, M., Kondratenko, R., Baltina, L., Tolstikov, G. A., Cinatl, J. (2005). Antiviral activity of glycyrrhizic acid derivatives against SARS-coronavirus. *Journal of Medicinal Chemistry*, 48(4), 1256–1259. <https://doi.org/10.1021/jm0493008>
- Hooker, J. D. (1973). JD 1872-1897. The Flora of British India, 1(7).
- Hostettmann, K. (1995). Saponins/K. Hostettmann and A. Marston. Cambridge Univ. Press.
- Hostettmann, K., & Marston, A. (1995). Saponins. In UK: Cambridge University Press.
- <https://powo.science.kew.org/taxon/urn:lsid:ipni.org:names:520187-1/general-information>
- <https://www.berkem.com/en/expertise/en/plantextraction> (accessed on 19-12-2022)
- <https://www.ncbi.nlm.nih.gov>
- <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7398001/>
- <https://www.slideshare.net/KaustavDey21/recent-advances-in-extraction> (Last accessed: 5-10-2022)

<https://www.ssi.shimadzu.com/service-support/faq/liquid-chromatography/knowledge-base/hplc-basics/index.html> (accessed on 17-12-2022)

<https://www.tridge.com/intelligences/licorice-root2/export> (accessed on 19-12-2022)

<https://www.volza.com/p/medicinal-plant/export/export-from-india/>. (accessed on 19-12-2022).

ICH, H. T. G. (2005). Validation of analytical procedures: text and methodology, Q2 (R1). Current Step 4 Version, Parent Guidelines on Methodology Dated November 6, 1996. Incorporated in November 2005. In *International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use, Geneva, Switzerland*.

Ikeda, K. (2007). Glycyrrhizin injection therapy prevents hepatocellular carcinogenesis in patients with interferon-resistant active chronic hepatitis C. *Hepatology Research : The Official Journal of the Japan Society of Hepatology*, 37 Suppl 2, S287—93. <https://doi.org/10.1111/j.1872-034x.2007.00199.x>

Imai, T., Nomura, M., & Fukushima, K. (2006). Evidence for involvement of the phenylpropanoid pathway in the biosynthesis of the norlignan agatharesinol. *Journal of Plant Physiology*, 163(5), 483–487. <https://doi.org/10.1016/j.jplph.2005.08.009>

Indian Pharmacopoeia. (2010). Ghaziabad: The Indian Pharmacopoeia Commission.

Iwasaki, T., Kondo, K., Kuroda, T., Moritani, Y., Yamagata, S., Sugiura, M., Kikkawa, H., Kaminuma, O., & Ikezawa, K. (1996). Novel selective PDE IV inhibitors as antiasthmatic agents. Synthesis and biological activities of a series of 1-aryl-2,3-bis(hydroxymethyl)naphthalene lignans. *Journal of Medicinal Chemistry*, 39(14), 2696–2704. <https://doi.org/10.1021/jm9509096>

Jacob, J. (2012). Microwave Assisted Reactions in Organic Chemistry: A Review of Recent Advances. *International Journal of Chemistry*, 4(6), 29–43. <https://doi.org/10.5539/ijc.v4n6p29>

Jadhav, C. A., Vikhe, D. N., & Jadhav, R. S. (2020). Global and domestic market of herbal medicines: a review. *Research Journal of Science and Technology*, 12(4), 327-330.

Jahan, B., Vahidy, A. A., & Ali, S. I. (1994). Chromosome numbers in some taxa of Fabaceae mostly native to Pakistan. *Annals of the Missouri Botanical Garden*, 792-799.

- Jamdhade, V. C., & Surwase, B. S. (2013). Memory enhancing activity of *Taverniera cuneifolia* (Roth) Arn: A substitute for commercial Liquorice. *Int J Pharm Bio Sci.* Apr, 4(2), 277-285.
- Jamdhade, V. C., Tembhurne, S. V., Kamble, S., & Surwase, B. S. (2015). Gastroprotective and Antioxidant Potential of Methanol Root Extract of *Taverniera cuneifolia* (Roth) Arn. in Albino Wister Rat. *Journal of Biologically Active Products from Nature*, 5(2), 150-162.
- Jamdhade, VC., Surwase, BS., Joshi, EB., Joshi, PN., & Jain, BK. (2013). Ecology and Marketing Pattern of Ethnomedicinal Plants of Kachchh District, Gujarat (India). *Life Sciences Leaflets*, 35-46
- Jan, S., Ahmad, J., Dar, M. M., Wani, A. A., Tahir, I., & Kamili, A. N. (2022). Development and validation of a reverse phase HPLC–DAD method for separation, detection & quantification of rutin and quercetin in buckwheat (*Fagopyrum* spp.). *Journal of Food Science and Technology*, 59(7), 2875-2883.
- Jayaprakasam, R., Saleshier, MF., & Ravi, TK. (2014). Development and validation of HPTLC and HPLC methods for the standardization of stigmasterol from the extract of *Acalypha indica* and its formulation. *World J Pharm Pharm Sci.*, 3:729-41.
- Jeandet, P., Delaunois, B., Conreux, A., Donnez, D., Nuzzo, V., Cordelier, S., ... & Courot, E. (2010). Biosynthesis, metabolism, molecular engineering, and biological functions of stilbene phytoalexins in plants. *Biofactors*, 36(5), 331-341.
- Jha, D. K., Shah, D. S., Talele, S. R., & Amin, P. D. (2020). Correlation of two validated methods for the quantification of naringenin in its solid dispersion: HPLC and UV spectrophotometric methods. *SN Applied Sciences*, 2(4), 1-11.
- Jones, M. E. (1953). Albrecht Kossel, a biographical sketch. *The Yale Journal of Biology and Medicine*, 26(1), 80.
- Joshi, E. B., Jain, B. K., Joshi, P. N., & Soni, H. B. (2013). Prevalence of traditional medications through native floral elements among tribal communities of Kachchh arid ecosystem, Gujarat, India. *International journal of environment*, 2(1), 184-201.
- Joshi, PN., Soni, HB., Sunderraj, SFW & Joshua, J. (2013). Status and distribution of threatened and endemic *Helichrysum cutchicum* (Asteraceae) in Arid Ecosystem of Kachchh, Gujarat, India. *Int. Journ. Life Sciences Leaflets*, 11 (11): 93-100.
- Journal, A. I., Polu, P., Nayanabhirama, U., & Khan, S. (2015). Herbal medicinal plants as an

anticancer agents, 4(1), 37–45.

Kamboj, A. (2012). Analytical Evaluation of Herbal Drugs. In: Vallisuta O, editor. Europe: In Tech, p. 1-39.

Kamboj, V. P. (2000). Herbal medicine. *Current science*, 78(1), 35-39.

Kameoka, H., & Nakai, K. (1987). Components of essential oil from the root of *Glycyrrhiza glabra*. *Nippon Nageikagaku Kaishi*, 61(9), 1119-1121.

Karthikeyan, S & Kumar, A. (1993). Flora of Yavatmal District, Maharashtra.

Kashiwada, Y., Nonaka, G. I., Nishioka, I., Chang, J. J., & Lee, K. H. (1992). Antitumor agents, 129. Tannins and related compounds as selective cytotoxic agents. *Journal of Natural Products*, 55(8), 1033–1043. <https://doi.org/10.1021/np50086a002>

Kaufman, T. S., & Rúveda, E. A. (2005). The quest for quinine: those who won the battles and those who won the war. *Angewandte Chemie International Edition*, 44(6), 854-885.

Kaufmann, B., & Christen, P. (2002). Recent Extraction Techniques for Natural Products: Microwave-assisted Extraction and Pressurised Solvent Extraction. *Phytochemical Analysis*, 13, 105–113.

Kawazoe, K., Yutani, A., Tamemoto, K., Yuasa, S., Shibata, H., Higuti, T., & Takaishi, Y. (2001). Phenylanthralene compounds from the subterranean part of *Vitex rotundifolia* and their antibacterial activity against methicillin-resistant *Staphylococcus aureus*. *Journal of Natural Products*, 64(5), 588–591. <https://doi.org/10.1021/np000307b>

Kemppainen, K., Siika-aho, M., Pattathil, S., Giovando, S., & Kruus, K. (2014). Spruce bark as an industrial source of condensed tannins and non-cellulosic sugars. *Industrial Crops and Products*, 52, 158-168.

Keymanesh, K., Hamed, J., Moradi, S., Mohammadipanah, F., & Sardari, S. (2009). Antibacterial, antifungal and toxicity of rare Iranian plants. *IJP-International Journal of Pharmacology*, 5(1), 81-85.

Khalighi-Sigaroodi, F., Ahvazi, M., Hadjiakhoondi, A., Taghizadeh, M., Yazdani, D., Khalighi-Sigaroodi, S., & Bidel, S. (2012). Cytotoxicity and antioxidant activity of 23 plant species of Leguminosae family. *Iranian journal of pharmaceutical research: IJPR*, 11(1), 295.

- Khalighi-Sigaroodi, F., Jeddi-Tehrani, M., Ahvazi, M., Shahnazi, S., Bayat, A. A., Mohajer, N., & Zarei, S. (2014). Cytotoxicity evaluation of *Taverniera spartea* on human cancer cell lines. *Journal of Medicinal Plants*, 13(50), 114-128.
- Khan, M. K., Abert-Vian, M., Fabiano-Tixier, A. S., Dangles, O., & Chemat, F. (2010). Ultrasound-assisted extraction of polyphenols (flavanone glycosides) from orange (*Citrus sinensis* L.) peel. *Food chemistry*, 119(2), 851-858.
- Khan, SW., Pokle, DS., & Pal, SC. (2012). Isolation and Hptlc Estimation of Phytochemicals from *Taverniera Cuneifolia* (Fabaceae). *Int. J. Nat. Prod. Res.*, 1(4): 75-78).
- Khanbabaee, K., & van Ree, T. (2001). Tannins: Classification and definition. *Natural Product Reports*, 18(6), 641–649. <https://doi.org/10.1039/b101061l>
- Khare, CP. (2007) *Indian Medicinal Plants: An Illustrated Dictionary*, published by Springer Science and Business media, LLC, 233 spring street, New York, USA, 647.
- Khonsa, K., Setyaningrum, D. L., Saputro, A. H., Amelia, T., Ibrahim, S., & Damayanti, S. (2022). Analysis of B-Sitosterol in Supplement Using High-Performance Liquid Chromatography: Development and Validation. *methods*, 15(3), 1997-2003.
- Khoo, H. E., & Ismail, A. (2008). Determination of daidzein and genistein contents in *Mangifera* fruit. *Malaysian Journal of Nutrition*, 14(2), 189-198.
- Kim, D., Park, J. B., Choi, W. K., Lee, S. J., Lim, I., & Bae, S. K. (2016). Simultaneous determination of β -sitosterol, campesterol, and stigmasterol in rat plasma by using LC–APCI-MS/MS: Application in a pharmacokinetic study of a titrated extract of the unsaponifiable fraction of *Zea mays* L. *Journal of separation science*, 39(21), 4060-4070.
- Kim, J. K., Oh, S. M., Kwon, H. S., Oh, Y. S., Lim, S. S., & Shin, H. K. (2006). Anti-inflammatory effect of roasted licorice extracts on lipopolysaccharide-induced inflammatory responses in murine macrophages. *Biochemical and Biophysical Research Communications*, 345(3), 1215-1223.
- Kimbaris, A. C., Siatis, N. G., Daferera, D. J., Tarantilis, P. A., Pappas, C. S., & Polissiou, M. G. (2006). Comparison of distillation and ultrasound-assisted extraction methods for the isolation of sensitive aroma compounds from garlic (*Allium sativum*). *Ultrasonics sonochemistry*, 13(1), 54-60.

- Kinoshita, T., Saitoh, T., & Shibata, S. (1978). A new 3-arylcoumarin from licorice root. *Chemical and Pharmaceutical Bulletin*, 26(1), 135-140.
- Kishore, N., Mishra, B. B., Tripathi, V., & Tiwari, V. K. (2009). Alkaloids as potential anti-tubercular agents. *Fitoterapia*, 80(3), 149-163.
- Kiuchi, F., Itano, Y., Uchiyama, N., Honda, G., Tsubouchi, A., Nakajima-Shimada, J., & Aoki, T. (2002). Monoterpene hydroperoxides with trypanocidal activity from *Chenopodium ambrosioides*. *Journal of Natural Products*, 65(4), 509-512. (crossref)
- Koch, K. (2004). Sucrose metabolism: Regulatory mechanisms and pivotal roles in sugar sensing and plant development. *Current Opinion in Plant Biology*, 7(3), 235–246. <https://doi.org/10.1016/j.pbi.2004.03.014>
- Koch, K. E. (1996). Carbohydrate-modulated gene expression in plants. *Annual review of plant biology*, 47(1), 509-540.
- Kolbe, L., Immeyer, J., Batzer, J., Wensorra, U., Mundt, C., Wolber, R., ... & Wenck, H. (2006). Anti-inflammatory efficacy of Licochalcone A: correlation of clinical potency and in vitro effects. *Archives of dermatological research*, 298(1), 23-30.
- Korenaga, M., Hidaka, I., Nishina, S., Sakai, A., Shinozaki, A., Gondo, T., & Hino, K. (2011). A glycyrrhizin-containing preparation reduces hepatic steatosis induced by hepatitis C virus protein and iron in mice. *Liver International*, 31(4), 552-560. <https://doi.org/10.1111/j.1478-3231.2011.02469.x>
- Kroymann, J. (2011). Natural diversity and adaptation in plant secondary metabolism. *Curr. Opin. Plant Biol.* 14, 246–251. doi: 10.1016/j.pbi.2011.03.021
- Kulkarni, S., Palghadmal, S., Sawant, A., Bhope, S., & Padmanabhan, S. (2021). RP-HPLC method development and validation for simultaneous estimation of glabridin, glycyrrhizinic acid and total flavonoids in *Glycyrrhiza glabra* extract: stability studies of constituents, *Glob. J. Pharmaceu. Sci*, 8(3).
- Kulkarni, TR., Mateenuddin, M., Sahasrabude, RA., & Pandit, VA. (2012). Estrogenic effect of alcoholic extract of lemon seeds (*Citrus limonum*) on immature albino rats. *International Journal of Research in Pharmaceutical and Biomedical Sciences*, 3:1231-5.

- Kumar N., Gupta S., Yadav T.C., Pruthi V., Varadwaj P.K., Goel N. Extrapolation of phenolic compounds as multi-target agents against cancer and inflammation. *J. Biomol. Struct. Dyn.* 2019;37(9):2355–2369.
- Kumar, N., & Pruthi, V. (2014). Potential applications of ferulic acid from natural sources. *Biotechnology Reports*, 4(1), 86–93. <https://doi.org/10.1016/j.btre.2014.09.002>
- Kumar, S., & Pandey, A. K. (2013). Phenolic content, reducing power and membrane protective activities of *Solanum xanthocarpum* root extracts.
- Kumar, S., Mishra, A., & Pandey, A. K. (2013). Antioxidant mediated protective effect of *Parthenium hysterophorus* against oxidative damage using in vitro models. *BMC complementary and alternative medicine*, 13(1), 1-9.
- Kumari, V. C., Patil, S. M., Ramu, R., Shirahatti, P. S., Kumar, N., Sowmya, B. P., ... & Patrick-Iwuanyanwu, K. C. (2022). Chromatographic techniques: types, principles, and applications.
- Kupiec, T. (2004). Quality-control analytical methods: High-performance liquid chromatography. *International journal of pharmaceutical compounding*, 8, 223-227.
- Kurek, J. (Ed.). (2019). Alkaloids: Their importance in Nature and Human life. BoD–Books on Demand.
- Kushwaha, SKS., Kushwaha, N., Maurya, N., & Rai, AK. (2010). Role of markers in standardization of herbal drugs: A review. *Arch Appl Sci Res.*, 2:225-9.
- Laghari, A. Q., Memon, S., Nelofar, A., & Laghari, A. H. (2011). Extraction, Identification and Antioxidative Properties of the Flavonoid-Rich Fractions from Leaves and Flowers of *Cassia angustifolia*; *American Journal of Analytical Chemistry*, 02(08), 871–878. <https://doi.org/10.4236/ajac.2011.28100>
- Lee, H. G., Kim, H., Suh, M. C., Kim, H. U., & Seo, P. J. (2018). The MYB96 transcription factor regulates triacylglycerol accumulation by activating DGAT1 and PDAT1 expression in *Arabidopsis* seeds. *Plant and Cell Physiology*, 59(7), 1432-1442. (crossref)
- Lee, K. R., Chen, G. Q., & Kim, H. U. (2015). Current progress towards the metabolic engineering of plant seed oil for hydroxy fatty acids production. *Plant cell reports*, 34(4), 603-615. (crossref)

- Lee, M. K., Kim, S. H., Park, J. H., Cho, J. H., Kim, D. H., Baek, J. H., ... & Sung, S. H. (2009). Determination and identification of nine constituents in Siho-Gyeoji-Tang by HPLC-DAD and HPLC-MS/MS. *Journal of Liquid Chromatography & Related Technologies*®, 32(14), 2122-2133.
- Lee, S. B., & Suh, M. C. (2015). Advances in the understanding of cuticular waxes in *Arabidopsis thaliana* and crop species. *Plant cell reports*, 34(4), 557-572. (crossref)
- Lewis, N. G., & Davin, L. B. (1999). Lignans: Biosynthesis and Function. *Comprehensive Natural Products Chemistry*, 639–712. <https://doi.org/10.1016/b978-0-08-091283-7.00027-8>
- Li, P., Yin, Z. Q., Li, S. L., Huang, X. J., Ye, W. C., & Zhang, Q. W. (2014). Simultaneous determination of eight flavonoids and pogostone in *Pogostemon cablin* by high performance liquid chromatography. *Journal of Liquid Chromatography & Related Technologies*, 37(12), 1771-1784.
- Li, X., Sun, R., & Liu, R. (2019). Natural products in licorice for the therapy of liver diseases: Progress and future opportunities. *Pharmacological Research*, 144, 210-226.
- Li. Y., Shen. Y., Yao. C.L., & Guo, D.A. (2020). Quality assessment of herbal medicines based on chemical fingerprints combined with chemometrics approach: A review. *J. Pharm. Biomed. Anal*, 185:113215. doi: 10.1016/j.jpba.2020.113215.
- Lin, A., Liu, Y., Huang, Y., Sun, J., Wu, Z., Zhang, X., & Ping, Q. (2008). Glycyrrhizin surface-modified chitosan nanoparticles for hepatocyte-targeted delivery. *International Journal of Pharmaceutics*, 359(1), 247–253. [https://doi.org/https://doi.org/10.1016/j.ijpharm.2008.03.039](https://doi.org/10.1016/j.ijpharm.2008.03.039)
- Lin, L., Lin, H., Zhang, M., Dong, X., Yin, X., Qu, C., & Ni, J. (2015). Types, principle, and characteristics of tandem high-resolution mass spectrometry and its applications. *RSC advances*, 5(130), 107623-107636.
- London, E. (2002). Insights into lipid raft structure and formation from experiments in model membranes. *Curr. Opin. Struct. Biol.*, 12, 480–486.
- Loos, G., Van Schepdael, A., & Cabooter, D. (2016). Quantitative mass spectrometry methods for pharmaceutical analysis. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 374(2079), 20150366.

- Lope Pihie, A. H., Zakaria, Z. A., & Othman, F. (2012). Antiproliferative and proapoptotic effects of labisia pumila ethanol extract and its active fraction in human melanoma HM3KO cells. *Evidence-Based Complementary and Alternative Medicine*, <https://doi.org/10.1155/2012/123470>
- Lu, H., & Liu, G. T. (1992). Antioxidant activity of dibenzocyclooctene lignans isolated from schisandraceae. *Planta Medica*, 58(4), 311–313. <https://doi.org/10.1055/s-2006-961473>
- Luo, X., Bai, R., Zhen, D., Yang, Z., Huang, D., Mao, H., ... & Fu, C. (2019). Response surface optimization of the enzyme-based ultrasound-assisted extraction of acorn tannins and their corrosion inhibition properties. *Industrial Crops and Products*, 129, 405-413.
- Luthria, D. L. (2008). Influence of experimental conditions on the extraction of phenolic compounds from parsley (*Petroselinum crispum*) flakes using a pressurized liquid extractor. *Food Chemistry*, 107(2), 745–752. <https://doi.org/10.1016/j.foodchem.2007.08.074>
- Mahdi-Pour, B., Jothy, S. L., Latha, L. Y., Chen, Y., & Sasidharan, S. (2012). Antioxidant activity of methanol extracts of different parts of Lantana camara. *Asian Pacific journal of tropical biomedicine*, 2(12), 960-965.
- Maji, AK., Maity, N., Panerji, P., & Banerjee, DA. (2014). validated RP-HPLC-UV method for quantitative determination of puerarin in Pueraria tuberosa DC tuber extract. *Pharm Methods*, 3:79-83.
- Mangalorkar, P. (2014). *Bioprospecting the potential of Taverniera cuneifolia (Roth) Ali* (Doctoral dissertation).
- Mangalorkar, P. S. (2013). Bioprospecting the potential of Taverniera cuneifolia Roth Ali.
- Mangalorkar, P. S., Rana, K. G., Parikh, R. D., & Nagar, P. S. (2013). The Effect of Different Chemical Treatments and Salt Stress on the Germination Potential of Tavernieracuneifolia (Roth) Ali seeds. *Journal of Chemical, Biological and Physical Sciences (JCBPS)*, 4(1), 369.
- Mangalorkar, P. Shailajan, S., & Nagar, P.S. (2014). HPTLC Analysis of the roots of *Taverniera cuneifolia* (Roth.) Ali. *J. of Pharma. And Scientific Innovation*, 3 (2): 143-147.
- Mangalorkar, P., Patel, B., & Nagar, P. (2013). Anatomy and pharmacognosy of Taverniera cuneifolia root (roth) arn., a possible substitute of glycyrrhiza glabra l. *Int. J. Pharm. Biol. Sci.*, 4, 221-226.

- Mangalorkar, P., Rana, K.G., Parikh, R.D., & Nagar, P.S. (2014). The effect of Different Chemical treatments and Salt Stress on the Germination potential of *Taverniera cuneifolia* (Roth) Ali seeds. *J. Chem. Bio. Phy. Sci.*, Sec. 4(1): 369-374.
- Manns, M. P., Wedemeyer, H., Singer, A., Khomutjanskaja, N., Dienes, H. P., Roskams, T., ... Moroz, L. (2012). Glycyrrhizin in patients who failed previous interferon alpha-based therapies: Biochemical and histological effects after 52 weeks. *Journal of Viral Hepatitis*, 19(8), 537–546. <https://doi.org/10.1111/j.1365-2893.2011.01579.x>.
- Manske, R. H. F. (1965). The Alkaloids. *Chemistry and Physiology*. New York, NY: Academic Press.
- Marjan, G. (2019). High-Performance Thin-Layer Chromatography (HPTLC) Method for Estimation of Glycyrrhizic Acid (Ga) In A Herbal Extract of *Taverniera Cuneifolia*. *Int J Recent Sci Res.*, 10(12), pp. 36313-36315. <http://dx.doi.org/10.24327/ijrsr.2019.1012.4899>
- Marston, A. (2007). Role of advances in chromatographic techniques in phytochemistry. *Phytochemistry*, 68:2785-97.
- Martin, V. J., Pitera, D. J., Withers, S. T., Newman, J. D., & Keasling, J. D. (2003). Engineering a mevalonate pathway in *Escherichia coli* for production of terpenoids. *Nature biotechnology*, 21(7), 796-802.
- Martinelli, E., Schulz, K., and Mansoori, G. A. (1991). Supercritical fluid extraction/retrograde condensation with applications in biotechnology. *Bruno, T.J. and Ely, J.F., Eds.*; CRC Press, Boca Raton, FL, 451–478.
- McNaught, A. D. (1997). *Compendium of chemical terminology* (Vol. 1669). Oxford: Blackwell Science.
- Medina, I., Gallardo, J.M., Gonzalez, M.J., Lois, S., & Hedges, N. (2007). Effect of molecular structure of phenolic families as hydroxycinnamic acids and catechins on their antioxidant effectiveness in minced fish muscle. *J. Agric. Food Chem.*, 55:3889–3895.
- Meena, A.K., Singh, A., Sharma, K., Kumari, S., & Rao, M.M. (2010). Physicochemical and preliminary phytochemical studies on the rhizomes of *Glycyrrhiza glabra* Linn. *Int J Pharmacy Pharm Sci.* 2010;2(2):48–50.
- Metre, V., & Ghorpade, J. (2013). An overview of the research on texture based plant leaf classification. *arXiv preprint arXiv:1306.4345*.

- Modh & pandya (2019). *Journal of Pharmacognosy and Phytochemistry*, 8(3): 834-836.
- Moghimipour, E., & Handali, S. (2015). Saponin: properties, methods of evaluation and applications. *Annual Research & Review in Biology*, 207-220.
- Montoro, P., Maldini, M., Russo, M., Postorino, S., Piacente, S., & Pizza, C. (2011). Metabolic profiling of roots of liquorice (*Glycyrrhiza glabra*) from different geographical areas by ESI/MS/MS and determination of major metabolites by LC-ESI/MS and LC-ESI/MS/MS. *Journal of pharmaceutical and biomedical analysis*, 54(3), 535–544. <https://doi.org/10.1016/j.jpba.2010.10.004>
- Moon, H. I., & Lee, J. H. (2012). Neuroprotective effects of triterpene glycosides from *Glycine max* against glutamate induced toxicity in primary cultured rat cortical cells. *Int. J. Mol. Sci.*, 13, 9642- 9648.
- Moresco, HH., Pereira, M., Bretanha, LC., Micke, GA., Pizzolatti, MG., & Brighente, IMC. (2014). Myricitin as the main constituent of two species of *Myrica*. *J App Pharm Sci.*, 4:1-7.
- Mori H., Kawabata K., Yoshimi N., Tanaka T., Murakami T., Okada T., & Murai H. (1999). Chemo preventive effects of ferulic acid on oral and rice germ on large bowel carcinogenesis. *Anticancer Res.*, 19:3775–3778.
- Morikawa, T., Mizutani, M., Aoki, N., Watanabe, B., Saga, H., Saito, S., Oikawa, A., Suzuki, H., Sakurai, N., & Shibata, D. (2006). Cytochrome P450 *CYP710A* encodes the sterol C-22 desaturase in *Arabidopsis* and tomato. *Plant Cell*, 18, 1008–1022.
- Morlock, G. E., Meyer, S., Zimmermann, B. F., & Roussel, J. M. (2014). High-performance thin-layer chromatography analysis of steviol glycosides in *Stevia* formulations and sugar-free food products, and benchmarking with (ultra) high-performance liquid chromatography. *Journal of Chromatography A*, 1350, 102-111.
- Moutsatsu, P. (2007). The spectrum of phytoestrogens in nature: Our knowledge is expanding. *Hormones*, 6:173-93.
- Mukherjee, P. K. (2002). *Quality control of herbal drugs: an approach to evaluation of botanicals*. Business horizons.
- Mukherjee, P.W. (2002). *Quality Control of Herbal Drugs: An Approach to Evaluation of Botanicals*. New Delhi, India: Business Horizons Publishers.

- Munns, R. (2005). Genes and salt tolerance: bringing them together. *New phytologist*, 167(3), 645-663.
- N. P. Committee, (2020). Pharmacopoeia of the People's Republic of China. Part 1. China Medical Science and Technology Press.
- Naczki, M., & Shahidi, F. (2006). Phenolics in cereals, fruits and vegetables: Occurrence, extraction and analysis. *Journal of Pharmaceutical and Biomedical Analysis*, 41(5), 1523–1542. <https://doi.org/10.1016/j.jpba.2006.04.002>
- Nagar, P. (2008) Medicinal Plants of Saurashtra, Scientific Publishers, Jodhpur.
- Naidu, B. P., Paleg, L. G., Aspinall, D., Jennings, A. C., & Jones, G. P. (1991). Amino acid and glycine betaine accumulation in cold-stressed wheat seedlings. *Phytochemistry*, 30(2), 407-409.
- Nandhini & Ilango (2020). *Pharmacogn J.*, 12(4): 850-856. 10.5530/pj.2020.12.122
- Nasir, E., Ali, S. I., & Nasir, Y. J. (1974). Flora of west Pakistan. University of Karachi.
- Nelson, D.R. (2006). Plant cytochrome P450s from moss to poplar. *Phytochem. Rev.*, 5, 193–204.
- Newman, D. J., & Cragg, G. M. (2012). Natural products as sources of new drugs over the 30 years from 1981 to 2010. *Journal of natural products*, 75(3), 311-335.
- Newman, D. J., & Cragg, G. M. (2016). Natural products as sources of new drugs from 1981 to 2014. *Journal of natural products*, 79(3), 629-661.
- Ngo, L. T., Okogun, J. I., & Folk, W. R. (2013). 21st century natural product research and drug development and traditional medicines. *Natural product reports*, 30(4), 584-592.
- Noamesi, B. K., Bogale, M., & Dagne, E. (1990). Intestinal smooth muscle spasmolytic actions of the aqueous extract of the roots of *Taverniera abyssinica*. *Journal of ethnopharmacology*, 30(1), 107-113.
- Oakenfull, D. (1981). Saponins in food—a review. *Food chemistry*, 7(1), 19-40.
- Oakenfull, D., Sidhu, G. S., & Cheeke, P. R. (1989). Saponins.
- Ohlrogge, J., & Browse, J. (1995). Lipid biosynthesis. *The Plant Cell*, 7(7), 957. (crossref)

- Okuda, T., Yoshida, T., Ashida, M., & Yazaki, K. (1983). Tannis of Casuarina and Stachyurus species. Part 1. Structures of pendunculagin, casuarictin, strictinin, casuarinin, casuariin, and stachyurin. *Journal of the Chemical Society, Perkin Transactions 1*, 1765-1772.
- Olszewska, M. (2008). Separation of quercetin, sexangularetin, kaempferol and isorhamnetin for simultaneous HPLC determination of flavonoid aglycones in inflorescences, leaves and fruits of three Sorbus species. *Journal of pharmaceutical and biomedical analysis*, 48(3), 629-635.
- Orhan, E. I., Tosun, F., Tamer, U., Duran, A., Alan, B., & Kök, F. A. (2011). Quantification of genistein and daidzein in two endemic Genista species and their antioxidant activity. *Journal of the serbian chemical society*, 76(1), 35-42.
- Ortega, S. N., Santos-Neto, A. J., & Lancas, F. M. (2017). Development and optimization of a fast method for the determination of statins in human plasma using microextraction by packed sorbent (MEPS) followed by ultra-high-performance liquid chromatography-tandem mass spectrometry (UHPLC-MS/MS). *Analytical Methods*, 9(20), 3039-3048.
- Osborn, A. E. (2003). Saponins in cereals. *Phytochemistry*, 62(1), 1-4.
- Ostovari, A., Hoseinie, S. M., Peikari, M., Shadizadeh, S. R., & Hashemi, S. J. (2009). Corrosion inhibition of mild steel in 1 M HCl solution by henna extract: A comparative study of the inhibition by henna and its constituents (Lawson, Gallic acid, α -d-Glucose and Tannic acid). *Corrosion Science*, 51(9), 1935-1949.
- Owczarek, A., Magiera, A., Matczak, M., Piotrowska, D. G., Olszewska, M. A., & Marchelak, A. (2017). Optimisation of preparative HPLC separation of four isomeric kaempferol diglycosides from *Prunus spinosa* L. by application of the response surface methodology. *Phytochemistry Letters*, 20, 415-424.
- Özer, E. Ö., Platin, S., Akman, U., & Hortaçsu, Ö. (1996). Supercritical Carbon Dioxide Extraction of Spearmint Oil from Mint-Plant Leaves. *Canadian Journal of Chemical Engineering*, 74(6), 920–928. <https://doi.org/10.1002/cjce.5450740615>
- Palin, C. T. (1880). A list of plants of Kutch. *Bombay gazetteer*, 5.
- Pan, J. Y., Chen, S. L., Yang, M. H., Wu, J., Sinkkonen, J., & Zou, K. (2009). An update on lignans: Natural products and synthesis. *Natural Product Reports*, 26(10), 1251–1292. <https://doi.org/10.1039/b910940d>

- Pan, S. Y., Pan, S., Yu, Z. L., Ma, D. L., Chen, S. B., Fong, W. F., ... & Ko, K. M. (2010). New perspectives on innovative drug discovery: an overview. *Journal of Pharmacy & Pharmaceutical Sciences*, 13(3), 450-471.
- Pandey, A. K. (2007). Anti-staphylococcal activity of a pan-tropical aggressive and obnoxious weed *Parthenium hysterophorus*: an in vitro study. *National Academy Science Letters*, 30(11/12), 383-386.
- Parvaiz, M., Hussain, K., Khalid, S., Hussnain, N., Iram, N., Hussain, Z., & Ali, M. A. (2014). A review: Medicinal importance of *Glycyrrhiza glabra* L. (Fabaceae Family). *Global Journal of Pharmacology*, 8(1), 8–13. <https://doi.org/10.5829/idosi.gjp.2014.8.1>.
- Patel, P. (2019). Agricultural Revision in Drought Prone Arid Region of Kutch: People Led, Market Oriented Growth Under Adverse Climatic Conditions. 3rd World Irrigation Forum, Bali, Indonesia.
- Patil D, Patil A, Vadera K and Ansari A (2015): Standardization and quality control parameters of aerial parts (leaves and stem) of *Trigonella foenum-graecum* L. An important medicinal plant. *Journal of Chemical and Pharmaceutical Research*, 7(3): 163-170.
- Patil, A. G., Koli, S. P., & Patil, D. A. (2010). Pharmacognostical standardization and HPTLC fingerprint of *Crataeva tapia* linn. ssp. *odora* (jacob.) almeida leaves. *International Journal of Pharma and Bio Sciences*, 1, 2.
- Patil, A. R., Ghagare, P. M., Deshmane, B. J., & Kondawar, M. S. (2020). Review on Chromatography Principal Types and its Application. *Research Journal of Pharmaceutical Dosage Forms and Technology*, 12(1), 27-32.
- Pereira, D. M., Valentão, P., Pereira, J. A., & Andrade, P. B. (2009). Phenolics: From chemistry to biology. *Molecules*, 14(6), 2202-2211.
- Perur, NG., Subramanian, CK., Muhr, GR., & Roy, HE. (1973). Soil fertility Evaluation- To serve Indian Farmers. Mysore Department of Agriculture, Mysore University of Agricultural Sciences and U.S. *Agency for International development*, Bangalore.p.40,43, 44.
- Pharmacopoeia of the People's Republic of China. (2000). The Pharmacopeia Commission of PRC, Beijing.
- Piironen, V., Toivo, J., Puupponen-Pimiä, R., & Lampi, A. (2003). Plant sterols in vegetables, fruits and berries. *J. Sci. Food Agric.* 83, 330–337. 10.1002/jsfa.1316

- Pizzi, A. (2006). Recent developments in eco-efficient bio-based adhesives for wood bonding: opportunities and issues. *Journal of adhesion science and technology*, 20(8), 829-846.
- Pizzi, A. (2008). Tannins: major sources, properties and applications. In *Monomers, polymers and composites from renewable resources* (pp. 179-199). Elsevier
- Plant of the world online – [Taverniera Plants of the World Online | Kew Science](#)
- Pongsuwan, W., Bamba, T., Harada, K., Yonetani, T., Kobayashi, A., & Fukusaki, E. (2008). High-throughput technique for comprehensive analysis of Japanese green tea quality assessment using ultra-performance liquid chromatography with time-of-flight mass spectrometry (UPLC/TOF MS). *Journal of agricultural and food chemistry*, 56(22), 10705-10708.
- Popova, I. E., Hall, C., & Kubátová, A. (2009). Determination of lignans in flaxseed using liquid chromatography with time-of-flight mass spectrometry. *Journal of Chromatography A*, 1216(2), 217–229. <https://doi.org/10.1016/j.chroma.2008.11.063>
- Prajapati, N. D., Purohit, S. S., Sharma, A. K., & Kumar, T. (2003). A handbook of medicinal plants: A complete source book. In *A handbook of medicinal plants: a complete source book* (pp. 554-554).
- Prajapati, S. M., & Patel, B. R. (2015). A comparative clinical study of Jethimala (*Taverniera nummularia* Baker.) and Yashtimadhu (*Glycyrrhiza glabra* Linn.) in the management of Amlapitta. *Ayu*, 36(2), 157.
- Prajapati, S. M., & Patel, B. R. (2015). A comparative clinical study of Jethimala (*Taverniera nummularia* Baker.) and Yashtimadhu (*Glycyrrhiza glabra* Linn.) in the management of Amlapitta. *Ayu*, 36(2), 157.
- Prajapati, S., Notani, B., Harisha, C. R., & Patel, B. (2014). Pharmacognostic Evaluation of Stem and Root of *Taverniera nummularia* Baker. *Annals of Ayurvedic Medicine*.
- Price, K. R., Johnson, I. T., Fenwick, G. R., & Malinow, M. R. (1987). The chemistry and biological significance of saponins in foods and feedingstuffs. *Critical Reviews in Food Science & Nutrition*, 26(1), 27-135.
- Quinn, P. J., & Williams, W. P. (1978). Plant lipids and their Role in membrane function. 34(c), 109–173.

- Raghuveer, I., Anurag, K., Anumalik, Y., Nitika, G., Swadesh, K., Nikhil, G., ... & Himanshu, G. (2015). Metabolites in plants and its classification. *World Journal of Pharmacy and Pharmaceutical Sciences (WJPPS)*, 4(1), 287-305.
- Rahman, S., & Sultana, S. (2006). Chemopreventive activity of glycyrrhizin on lead acetate mediated hepatic oxidative stress and its hyperproliferative activity in Wistar rats. *Chemico-biological interactions*, 160(1), 61–69. <https://doi.org/10.1016/j.cbi.2005.12.003>
- Rajagopal, S. S., Vazhayil, B. K., Varghese, L., & Nanjaian, M. (2017). Development and validation of RP-HPLC method for simultaneous determination of apigenin and luteolin in ethanol extract of *Clerodendrum serratum* (Linn.) leaves. *Asian Journal of Applied Sciences*, 5(1).
- Ramya, V., Dhayalan, V. D., & Umamaheswari, S. (2010). In vitro studies on antibacterial activity and separation of active compounds of selected flower extracts by HPTLC. *Journal of Chemical and Pharmaceutical Research*, 2(6), 86-91.
- Rao, C. H. B. (2011). LC/MS: an essential tool in drug development. *Int. J. Adv. Pharm. Anal*, 1(2), 24-37.
- Rao, K. T., Krishna, I. S. R., & Reddy, C. S. (2007). Biodiversity of Nallamalai Hill Ranges, Eastern Ghats, India. In *Proceedings of the National Seminar on Conservation of Eastern Ghats* (pp. 214-221).
- Razack, S., Hemanth Kumar, K., Nallamuthu, I., Naika, M., & Khanum, F. (2015). Antioxidant, biomolecule oxidation protective activities of *Nardostachys jatamansi* DC and its phytochemical analysis by RP-HPLC and GC-MS. *Antioxidants*, 4(1), 185-203.
- Rhodes, D., Nadolska-Orczyk, A., & Rich, P. J. (2002). Salinity, osmolytes and compatible solutes. In *Salinity: Environment-plants-molecules* (pp. 181-204). Springer, Dordrecht.
- Ribeiro, I. A., & Ribeiro, M. H. (2008). Naringin and naringenin determination and control in grapefruit juice by a validated HPLC method. *Food Control*, 19(4), 432-438.
- “Rigveda,” http://www.indianetzone.com/2/rig_veda.htm.
- Riswanto, F. D. O., Desra, A., Sari, R. M., Thomas, V., Rohman, A., Pramono, S., & Martono, S. (2020). Employing an R software package rsm for optimizing of genistein, daidzein, and glycitein separation and its application for soymilk analysis by HPLC method. *Indonesian Journal of Chemistry*, 20(5), 1184-1198.

- Riviere, C., Pawlus, A. D., & Merillon, J. M. (2012). Natural stilbenoids: distribution in the plant kingdom and chemotaxonomic interest in Vitaceae. *Natural product reports*, 29(11), 1317-1333. (CrossRef)
- Rocha, RP., Melo, EC., & Radunz, LL. (2011). Influence of drying process on the quality of medicinal plants: A review. *J Med Plants Res.*, 5:7076-84.
- Rolland, F., Baena-Gonzalez, E., & Sheen, J. (2006). Sugar sensing and signaling in plants: Conserved and novel mechanisms. *Annual Review of Plant Biology*, 57, 675–709. <https://doi.org/10.1146/annurev.arplant.57.032905.105441>
- Rolland, F., Moore, B., & Sheen, J. (2002). Sugar sensing and signaling in plants. *Plant Cell*, 185–202.
- Rosales, P. F., Bordin, G. S., Gower, A. E., & Moura, S. (2020). Indole alkaloids: 2012 until now, highlighting the new chemical structures and biological activities. *Fitoterapia*, 143, 104558.
- Roxburgh, W. (1832). *Flora Indica* 3. W. Thacker & Co. Calcutta, Serampore. 875 pp.
- Rupasinghe, H. V., Nair, S. V., & Robinson, R. A. (2014). Chemo-preventive properties of fruit phenolic compounds and their possible mode of actions. *Studies in natural products chemistry*, 42, 229-266.
- Russell, W., & Duthie, G. (2011). Plant secondary metabolites and gut health: the case for phenolic acids. *Proceedings of the Nutrition Society*, 70(3), 389-396.
- SAIJO, R., NONAKA, G. I., & NISHIOKA, I. (1989). Tannins and related compounds. Lxxxiv.: Isolation and characterization of five new hydrolyzable tannins from the bark of *mallotus japonicus*. *Chemical and pharmaceutical bulletin*, 37(8), 2063-2070.
- Sakodinskii, K., & Chmutov, K. (1972). M. S. Tswett and Chromatography. *Chromatographia*, 5(8), 471–476. <https://doi.org/10.1007/bf02284393>
- Sami, F., Yusuf, M., Faizan, M., Faraz, A., & Hayat, S. (2016). Role of sugars under abiotic stress. *Plant Physiology and Biochemistry*, 109, 54-61.
- Sanghavi, N., Bhosale, SD., & Malode, S. (2014). RP-HPLC method development and validation of quercetin isolated from the plant *Tridax procumbens* L. *Journal of Scientific and Innovative Research*, 3:594-7.

- Sanjappa, M. 1992. Legumes of India. Bishen Singh Mahendra Pal Singh, Dehra Dun. pp. 175–179.
- Santapau, H. (1953). Plants of Saurashtra, a preliminary list. Saur. Res. Soc, Rajkot.
- Santapau, H. (1962). The flora of Saurashtra Part-1. (Ranunculaceae to Rubiaceae) Saurashtra Research Society, Rajkot:270.
- Sarwar, GR. (2002). Flora of Pakistan, University of Karachi, Missouri Botanical Press.
- Sasidharan, S., Chen, Y., Saravanan, D., Sundram, K. M., & Latha, L. Y. (2011). Extraction, isolation and characterization of bioactive compounds from plants' extracts. *African journal of traditional, complementary and alternative medicines*, 8(1).
- Satpathy, S., Patra, A., Hussain, M. D., & Ahirwar, B. (2017). Simultaneous estimation of genistein and daidzein in Pueraria tuberosa (Willd.) DC by validated high-performance thin-layer chromatography (HPTLC) densitometry method. *Journal of Liquid Chromatography & Related Technologies*, 40(10), 499-505.
- Savic, I. M., Nikolic, V. D., Savic, I. M., Nikolic, L. B., & Stankovic, M. Z. (2013). Development and validation of a new RP-HPLC method for determination of quercetin in green tea. *Journal of analytical chemistry*, 68(10), 906-911.
- Seasotiya L, Siwach P, Malik A, Bai S, Bharti P and Dalal S (2014): Phytochemical evaluation and HPTLC fingerprint profile of Cassia fistula. *International Journal of Advances in Pharmacy, Biology and Chemistry*, 3(3): 604-611.
- Seo, C. S., Lee, M. Y., Lim, H. S., Kim, S. J., Ha, H., Lee, J. A., & Shin, H. K. (2012). Determination of 5-hydroxymethyl-2-furfural, albiflorin, paeoniflorin, liquiritin, ferulic acid, nodakenin, and glycyrrhizin by HPLC-PDA, and evaluation of the cytotoxicity of Palmul-tang, a traditional Korean herbal medicine. *Archives of pharmacal research*, 35, 101-108.
- Sewelam, N., Jaspert, N., Van Der Kelen, K., Tognetti, V.B., Scmitz, J., Frerigmann, H., Stahl, E., Zeier, J., Van Breusege, F., & Maurino, V.G. (2014). Spatial H₂O₂ signaling specificity: H₂O₂ from chloroplasts and peroxisomes modulates the plant transcriptome differentially. *Mol. Plant*, 7, 1191–1210.
- Shah, GL. (1978). Flora of Gujarat State, Published by Sardar Patel University in Vallabh Vidyanagar, Vol.1.

- Shanker, K., Fatima, A., Negi, A. S., Gupta, V. K., Darokar, M. P., Gupta, M. M., & Khanuja, S. P. S. (2007). RP-HPLC method for the quantitation of glabridin in Yashti-madhu (*Glycyrrhiza glabra*). *Chromatographia*, 65(11), 771-774.
- Sharkey, T. D. (2021). Pentose phosphate pathway reactions in photosynthesizing cells. *Cells*, 10(6), 1547. (CrossRef)
- Sharma, A., Shanker, C., Tyagi, L. K., Singh, M., & Rao, C. V. (2008). Herbal medicine for market potential in India: an overview. *Acad J Plant Sci*, 1(2), 26-36.
- Sharma, BK. (2006) Instrumental methods of chemical analysis. (25th edn.); Goel publishing house, pp. 286-385.
- Sharma, K. P. (2019). Tannin degradation by phytopathogen's tannase: A Plant's defense perspective. *Bio catalysis and Agricultural Biotechnology*, 21, 101342.
- Sharma, O. P., Kumar, N., Singh, B., & Bhat, T. K. (2012). An improved method for thin layer chromatographic analysis of saponins. *Food chemistry*, 132(1), 671-674.
- Sharma, S. S., & Dietz, K. J. (2006). The significance of amino acids and amino acid-derived molecules in plant responses and adaptation to heavy metal stress. *Journal of experimental botany*, 57(4), 711-726.
- Sheen, J., Zhou, L., & Jang, J. C. (1999). Sugars as signaling molecules. *Current opinion in plant biology*, 2(5), 410-418.
- Shetti, P., & Jalalpure, S. S. (2021). A single robust stability-indicating RP-HPLC analytical tool for apigenin quantification in bulk powder and in nanoliposomes: a novel approach. *Future Journal of Pharmaceutical Sciences*, 7(1), 1-9.
- Shettigar, P. S. P. R. (2010). An advancement of analytical techniques in herbal research. *Journal of Advanced Scientific Research.*, 1(1), 8–14.
- Shetty, BV., & Singh, V. (1987). Flora of Rajasthan Vol.1.
- Shouqin, Z., Junjie, Z., & Changzhen, W. (2004). Novel high pressure extraction technology. *International Journal of Pharmaceutics*, 278(2), 471–474. <https://doi.org/10.1016/j.ijpharm.2004.02.029>
- Singh NP, 1988, Flora of Eastern Karnataka, Published by Mittal Publications, Jai Mata Market, Trinagar, Delhi: 257.

- Singh, P.S., Bal, M., Mukhtar, H.M., & Shah, G. (2011). Standardization and pharmacological investigation on leaves on *Ficus bengalensis*. *International Journal of Research in Pharmacy and Chemistry*, 1:891-903.
- Sipahimalani, J.L. (2002). Indian Herbal Pharmacopoeia. Mumbai: Indian Drug Manufacturers' Association.
- Sladkovský, R., Solich, P., & Opletal, L. (2001). Simultaneous determination of quercetin, kaempferol and (E)-cinnamic acid in vegetative organs of *Schisandra chinensis* Baill. by HPLC. *Journal of pharmaceutical and biomedical analysis*, 24(5-6), 1049-1054.
- Smeekens, S., Ma, J., Hanson, J., & Rolland, F. (2010). Sugar signals and molecular networks controlling plant growth. *Current Opinion in Plant Biology*, 13(3), 273–278. <https://doi.org/10.1016/j.pbi.2009.12.002>
- Sowmya, N., Haraprasad, N., & Hema, B. P. (2019). Exploring the total flavonoid content of peels of *Citrus aurantium*, *Citrus maxima* and *Citrus sinensis* using different solvents and HPLC-analysis of flavonones—Naringin and Naringenin in peels of *Citrus maxima*. *Pharma Innov*, 8, 12-17.
- Sparg, S., Light, M. E., & Van Staden, J. (2004). Biological activities and distribution of plant saponins. *Journal of ethnopharmacology*, 94(2-3), 219-243.
- Sriraman, S., Ramanujam, G. M., Ramasamy, M., & Dubey, G. P. (2015). Identification of beta-sitosterol and stigmasterol in *Bambusa bambos* (L.) Voss leaf extract using HPLC and its estrogenic effect in vitro. *Journal of Pharmaceutical and Biomedical Analysis*, 115, 55-61.
- Srivastava, S., & Srivastava, A. K. (2013). Biotechnology and genetic engineering for alkaloid production. *Natural Products: Phytochemistry, Botany and Metabolism of Alkaloids, Phenolics and Terpenes*; Ramawat, G.K., Mérillon, J.-M., Eds, 213-250.
- Stadler, M., Dagne, E., & Anke, H. (1994). Nematicidal activities of two phytoalexins from *Taverniera abyssinica*. *Planta medica*, 60(06), 550-552.
- Sticher, O. (2008). Natural product isolation. *Natural Product Reports*, 25(3), 517–554. <https://doi.org/10.1039/b700306b>
- Suga, T., Ohta, S., Munesada, K., Ide, N., Kurokawa, M., Shimizu, M., & Ohta, E. (1993). Endogenous pine wood nematicidal substances in pines, *Pinus massoniana*, *P. strobus* and *P. palustris*. *Phytochemistry*, 33(6), 1395-1401.

- Sulistyowati, E., Martono, S., Riyanto, S., Lukitaningsih, E., & Rohman, A. (2020). Rapid quantitative analysis of daidzein and genistein in soybeans (*Glycine max* (L). Merr.) using FTIR spectroscopy and multivariate calibration. *Journal of Applied Pharmaceutical Science*, 10(11), 117-123.
- Suman, A., Ali, M., Alam, P. (2009). New prenylated isoflavanones from the roots of *Glycyrrhiza glabra*. *Chem Nat Comp.*, 45(4):487–491. doi: 10.1007/s10600-009-9403-1.
- SUN, L. C., LI, S. Y., WANG, F. Z., & XIN, F. J. (2017). Research progresses in the synthetic biology of terpenoids. *Biotechnology Bulletin*, 33(1), 64.
- Suzuki, J., & El-Haddad, S. (2017). A review: fentanyl and non-pharmaceutical fentanyls. *Drug and alcohol dependence*, 171, 107-116.
- Taiwan Herbal Pharmacopeia (2019). 3rd ed. Taiwan Republic of China: Ministry of Health and Welfare.
- Tang, W., Sun, B., & Zhao, Y. (2014). Preparative separation and purification of rosmarinic acid from perilla seed meal via combined column chromatography. *Journal of Chromatography B*, 947, 41-48.
- Tang, Z. H., Li, T., Tong, Y. G., Chen, X. J., Chen, X. P., Wang, Y. T., & Lu, J. J. (2015). A systematic review of the anticancer properties of compounds isolated from Licorice (*Gancao*). *Planta medica*, 81(18), 1670-1687.
- Tatke, P. and Jaiswal, Y. (2011). An overview of microwave assisted extraction and its applications in herbal drug research. *Res. J Med. Plant.*, 5, 21–31.
- Ten Kate, K., & Laird, S. A. (2019). *The commercial use of biodiversity: access to genetic resources and benefit-sharing*. Routledge.
- Thaker, J. I. (1910). *Vanaspati shastra. Flora of Bardahill) Reprint, Pravin Pustak, Bhandar, Rajkot*, 34
- Ninh The Son. (2019). Secondar Metabolites of Genus *Pandanus*: An Aspect of Phytochemistry. *Mini-Reviews in Organic Chemistry*, 16(7), 689-710.
- Thakur, P. S., & Rai, V. K. (1982). Dynamics of amino acid accumulation of two differentially drought resistant *Zea mays* cultivars in response to osmotic stress. *Environmental and Experimental Botany*, 22(2), 221-226.

- The Society of Japanese Pharmacopeia. (2001). The Japanese Pharmacopeia. *Fourteenth* Ed., JP XIII, .
- Thulin, M. (1985). *Acta Univ. Upsalienses: Symb. Bot. Ups.*, 25:44-95.
- Tian, M., Yan, H., & Row, K. H. (2008). Simultaneous extraction and separation of liquiritin, glycyrrhizic acid, and glabridin from licorice root with analytical and preparative chromatography. *Biotechnology and bioprocess engineering*, 13(6), 671-676.
- Tomak, E. D., & Gonultas, O. (2018). The wood preservative potentials of valonia, chestnut, tara and sulphited oak tannins. *Journal of Wood Chemistry and Technology*, 38(3), 183-197.
- Tondi, G., Zhao, W., Pizzi, A., Du, G., Fierro, V., & Celzard, A. (2009). Tannin-based rigid foams: a survey of chemical and physical properties. *Bioresource technology*, 100(21), 5162-5169.
- Torres, P., Avila, J. G., de Vivar, A. R., García, A. M., Marín, J. C., Aranda, E., & Céspedes, C. L. (2003). Antioxidant and insect growth regulatory activities of stilbenes and extracts from *Yucca periculosa*. *Phytochemistry*, 64(2), 463-473.
- Trusheva, B., Trunkova, D., & Bankova, V. (2007). Different extraction methods of biologically active components from propolis; a preliminary study. *Chemistry Central Journal*, 1(1), 1–4. <https://doi.org/10.1186/1752-153X-1-13>
- USP Herbal Medicine Compendium (2021). General-Noticesresources. Available at: <https://hmc.usp.org/about/general-noticesresources> (Accessed August 26, 2021).
- USP. (2002). United States Pharmacopeia and National Formulary, USP.
- USP44-NF39 (2021b). General Chapters, General Test and Assay, Articles of Botanical Origin. Available at: https://online.uspnf.com/uspnf/document/1_GUID-E8A1366F-9657-41FC-9EDC-C20F4BE473B6_5_en-US (Accessed August 26, 2021).1–15.
- Valitova, J.N., Sulkarnayeva, A.G., & Minibayeva, F.V. (2016). Plant sterols: Diversity, biosynthesis, and physiological functions. *Biochemistry*, 81, 1050–1068.
- Vanhercke, T., Dyer, J. M., Mullen, R. T., Kilaru, A., Rahman, M. M., Petrie, J. R., ... & Singh, S. P. (2019). Metabolic engineering for enhanced oil in biomass. *Progress in lipid research*, 74, 103-129. (crossref)

Vermeris, W., & Nicholson, R. (2007). *Phenolic compound biochemistry*. Springer Science & Business Media.

Versluys, M., Tarkowski, Ł. P., & Van den Ende, W. (2017). Fructans as DAMPs or MAMPs: evolutionary prospects, cross-tolerance, and multistress resistance potential. *Frontiers in Plant Science*, 7, 2061.

Vickers, N. J. (2017). Animal communication: when i'm calling you, will you answer too? *Current biology*, 27(14), R713-R715.

Vidyadhar, S., Saidulu, M., Gopal, T. K., Chamundeeswari, D., Rao, U., & Banji, D. (2010). In vitro anthelmintic activity of the whole plant of *enicostemma littorale* by using various extracts.

Vijayalakshmi, R. A., & Ravindhran, R. (2012). Comparative fingerprint and extraction yield of *Diospyros ferrea* (willd.) Bakh. root with phenol compounds (gallic acid), as determined by uv-vis and ft-ir spectroscopy. *Asian Pacific Journal of Tropical Biomedicine*, 2(3), S1367-S1371.

Vijayanarayana, K., Rodrigues, R. S., Chandrashekhar, K. S., & Subrahmanyam, E. V. S. (2007). Evaluation of estrogenic activity of alcoholic extract of rhizomes of *Curculigo orchoides*. *Journal of Ethnopharmacology*, 114(2), 241-245.

Vincken, J. P., Heng, L., de Groot, A., & Gruppen, H. (2007). Saponins, classification and occurrence in the plant kingdom. *Phytochemistry*, 68(3), 275-297.

Viswanathan, V., & Mukne, A. P. (2016). Development and Validation of HPLC and HPTLC Methods for Estimation of Glabridin in Extracts of *Glycyrrhiza glabra*. *Journal of AOAC International*, 99(2), 374-379.

Von Rohr, P.R. & Trepp, C. (1996). *High Pressure Chemical Engineering*. Elsevier, Amsterdam.

Wang, H., Haibin, S. U., & Liu, H. (2004). Clinical research on treatment on several kinds of liver hepatitis with diammonium glycyrrhizinate. *Chinese Journal of Infectious Diseases*, (02), 113-115.

Wang, J., Chen, X., Wang, W., Zhang, Y., Yang, Z., Jin, Y., Ge, H. M., Li, E., & Yang, G. (2013). Glycyrrhizic acid as the antiviral component of *Glycyrrhiza uralensis* Fisch. against coxsackievirus A16 and enterovirus 71 of hand foot and mouth disease. *Journal of*
251

- ethnopharmacology*, 147(1), 114–121. <https://doi.org/10.1016/j.jep.2013.02.017>
- Wang, L., & Weller, C. L. (2006). Recent advances in extraction of nutraceuticals from plants. *Trends in Food Science and Technology*, 17(6), 300–312. <https://doi.org/10.1016/j.tifs.2005.12.004>
- Wang, L., Xu, R., Hu, B., Li, W., Sun, Y., Tu, Y., & Zeng, X. (2010). Analysis of free amino acids in Chinese teas and flower of tea plant by high performance liquid chromatography combined with solid-phase extraction. *Food Chemistry*, 123(4), 1259-1266.
- Wang, L., Yang, R., Yuan, B., Liu, Y., & Liu, C. (2015). The antiviral and antimicrobial activities of licorice, a widely-used Chinese herb. *Acta pharmaceutica sinica B*, 5(4), 310-315.
- Wang, Q., Qian, Y., Wang, Q., Yang, Y. F., Ji, S., Song, W., ... & Ye, M. (2015). Metabolites identification of bioactive licorice compounds in rats. *Journal of Pharmaceutical and Biomedical Analysis*, 115, 515-522.
- Wang, X., Usmanovich, K. R., Luo, L., Xu, W. J., & Wu, J. H. (2019). Salvation of rare and endangered medicinal plants. In *Molecular Pharmacognosy* (pp. 103-143). Springer, Singapore.
- Wang, Y. C., & Yang, Y. S. (2007). Simultaneous quantification of flavonoids and triterpenoids in licorice using HPLC. *Journal of Chromatography B*, 850(1-2), 392-399.
- WEINGES, K., & PLIENINGER, P. (1999). Erinnerungen a Karl Johann Freudenberg (1886 – 1983). *Eur. J. Org. Chem.*, 3, 707–736.
- Whistler, R. L., & Corbett, W. M. (1966). *The Encyclopedia of Chemistry*. Eds. G.L. Clark and G.G. Hawley, Reinhold Publishing Corp: New York.
- Wink, M. (2003). Evolution of secondary metabolites from an ecological and molecular phylogenetic perspective. *Phytochemistry*, 64(1), 3-19.
- Wink, M. (2010). “Introduction: biochemistry, physiology and ecological functions of secondary metabolites” in *Annual plant reviews: Biochemistry of plant secondary metabolism*. ed. M. Wink, Vol. 40 (Wiley-Blackwell), 1–19.
- World Health Organization Programme on Traditional Medicine. *Guidelines for the Assessment of Herbal Medicines*. 1991. [(accessed on 28 February 2022)]. Available

online: https://apps.who.int/iris/bitstream/handle/10665/58865/WHO_TRM_91.4.pdf?sequence=1&isAllowed=y

Wu, G., Bazer, F. W., Dai, Z., Li, D., Wang, J., & Wu, Z. (2014). Amino acid nutrition in animals: protein synthesis and beyond. *Annu Rev Anim Biosci*, 2(1), 387-417.

Wu, X. Q., Li, W., Chen, J. X., Zhai, J. W., Xu, H. Y., Ni, L., & Wu, S. S. (2019). Chemical constituents and biological activity profiles on *Pleione* (Orchidaceae). *Molecules*, 24(17). <https://doi.org/10.3390/molecules24173195>

Würdig, G., & Woller, R. (1989). *Chemie des Weins* Eugen Ulmer GmbH & Co. Stuttgart, 34(61), 201.

X. Zhang et al. / *Steroids* 70 (2005) 886–895.

Xu, K., Chu, F., Li, G., Xu, X., Wang, P., Song, J., Zhou, S., Lei, H. (2014). Oleanolic acid synthetic oligo glycosides: a review on recent progress in biological activities. *Pharmazie*, 69, 483- 495.

Yamada, M., Kita, Y., Kohira, T., Yoshida, K., Hamano, F., Tokuoka, S. M., & Shimizu, T. (2015). A comprehensive quantification method for eicosanoids and related compounds by using liquid chromatography/mass spectrometry with high speed continuous ionization polarity switching. *Journal of Chromatography B*, 995, 74-84.

Yamunadevi, M., Wesely, E. G., & Johnson, M. (2011). A Chromatographic Study on the Glycosides of *Aerva lanata* L. *Chinese Journal of Natural Medicines*, 9(3), 210-214.

Yang, C. W., Lin, C. C., & Kao, C. H. (2000). Proline, ornithine, arginine and glutamic acid contents in detached rice leaves. *Biologia plantarum*, 43(2), 305-307

Yang, Y., & Benning, C. (2018). Functions of triacylglycerols during plant development and stress. *Current Opinion in Biotechnology*, 49, 191-198. (crossref)

Yasui, S., Fujiwara, K., Tawada, A., Fukuda, Y., Nakano, M., & Yokosuka, O. (2011). Efficacy of intravenous glycyrrhizin in the early stage of acute onset autoimmune hepatitis. *Digestive diseases and sciences*, 56(12), 3638–3647. <https://doi.org/10.1007/s10620-011-1789-5>.

Ye, Z., Dai, J. R., Zhang, C. G., Lu, Y., Wu, L. L., Gong, A. G., ... & Wang, Z. T. (2017). Chemical differentiation of *Dendrobium officinale* and *Dendrobium devonianum* by using

HPLC fingerprints, HPLC-ESI-MS, and HPTLC analyses. *Evidence-Based Complementary and Alternative Medicine*, 2017.

Yi Sheng Xiao-Bin Chen, (2009). Vol.1, No.3, 203-206. doi:10.4236/health.2009.13034

Yi, Y., Zhang, Q. W., Li, S. L., Wang, Y., Ye, W. C., Zhao, J., & Wang, Y. T. (2012). Simultaneous quantification of major flavonoids in “Bawanghua”, the edible flower of *Hylocereus undatus* using pressurised liquid extraction and high performance liquid chromatography. *Food chemistry*, 135(2), 528-533.

Yoshida, T., Nakata, F., Hosotani, K., Nitta, A., & Okudat, T. (1992). Dimeric hydrolysable tannins from *Melastoma malabathricum*. *Phytochemistry*, 31(8), 2829-2833.

Yu, B. W., Chen, J. Y., Wang, Y. P., Cheng, K. F., Li, X. Y., & Qin, G. W. (2002). Alkaloids from *Menispermum dauricum*. *Phytochemistry*, 61(4), 439-442.

Yusop, Z., Jaafar, J., Aris, A. B., Majid, Z. A., Umar, K., & Talib, J. (2016). Development and validation of a selective, sensitive and stability indicating UPLC–MS/MS method for rapid, simultaneous determination of six process related impurities in darunavir drug substance. *Journal of pharmaceutical and biomedical analysis*, 128, 141-148.

Zarnowski, R., & Suzuki, Y. (2004). Expedient Soxhlet extraction of resorcinolic lipids from wheat grains. *Journal of Food Composition and Analysis*, 17(5), 649–663. <https://doi.org/10.1016/j.jfca.2003.09.007>

Zhang, H. F., Yang, X. H., & Wang, Y. (2011). Microwave assisted extraction of secondary metabolites from plants: Current status and future directions. *Trends in Food Science and Technology*, 22(12), 672–688. <https://doi.org/10.1016/j.tifs.2011.07.003>

Zhang, H. F., Yang, X. H., Zhao, L. D., & Wang, Y. (2009). Ultrasonic-assisted extraction of epimedin C from fresh leaves of *Epimedium* and extraction mechanism. *Innovative Food Science and Emerging Technologies*, 10(1), 54–60. <https://doi.org/10.1016/j.ifset.2008.09.007>

Zhao, T., Dong, Y., & Zhao, M (2013). Gas chromatography-mass spectrometry analysis of volatiles components obtained from the *Glycyrrhiza glabra* L. leaf and root. *Science and Technology of Food Industry*, 34, 96–99. (In Chinese)

Zheng, L., & Zheng, F. (2015). Development and validation of an LC-APCI-MS/MS method for the determination of phenethyl isothiocyanate in human plasma. *Biomedical Chromatography*, 29(4), 619-625.

- Zhou, W., Li, J., & Yue, T. (2020). *Remote Sensing Monitoring and Evaluation of Degraded Grassland in China*. Springer Science and Business Media LLC.
- Zhou, Y. Q., Zhang, Q. W., Li, S. L., Yin, Z. Q., Zhang, X. Q., & Ye, W. C. (2012). Quality evaluation of semen oroxyli through simultaneous quantification of 13 components by high performance liquid chromatography. *Current Pharmaceutical Analysis*, 8(2), 206-213.
- Zhu, H., Wang, Y., Liu, Y., Xia, Y., & Tang, T. (2010). Analysis of flavonoids in *Portulaca oleracea* L. by UV-vis spectrophotometry with comparative study on different extraction technologies. *Food Analytical Methods*, 3(2), 90–97. <https://doi.org/10.1007/s12161-009-9091-2>
- Zore, G. B., Winston, U. B., Surwase, B. S., Meshram, N. S., Sangle, V. D., Kulkarni, S. S., & Mohan Karuppayil, S. (2008). Chemoprofile and bioactivities of *Taverniera cuneifolia* (Roth) Arn.: a wild relative and possible substitute of *Glycyrrhiza glabra* L. *Phytomedicine: international journal of phytotherapy and phytopharmacology*, 15(4), 292–300. <https://doi.org/10.1016/j.phymed.2007.01.006>
- Zu, Y., Li, C., Fu, Y., & Zhao, C. (2006). Simultaneous determination of catechin, rutin, quercetin kaempferol and isorhamnetin in the extract of sea buckthorn (*Hippophae rhamnoides* L.) leaves by RP-HPLC with DAD. *Journal of pharmaceutical and biomedical analysis*, 41(3), 714-719.
- Zwenger Chhandak Basu, S., Zwenger, S., & Basu, C. (2008). School of Biological Sciences Faculty Publications. 4. *Biotechnology and Molecular Biology Reviews*, 3(1), 1–007.