#### **List of Tables**

# Chapter I

Table Number	Page No.
Table 1.1: System Suitability Test Parameters and Recommendations	42
Table 1.2: Important validation parameters suggested by regulatory	
agencies	43
Table 1.3: Last 5 years major licorice export data	48
Table 1.4: Last 5 years major licorice import data	50

# Chapter II

Table Number	Page No.
Table 2.1 Taxonomical status of Taverniera cuneifolia	54
Table 2.2: Geographical distribution of different species of <i>Taverniera</i> genus	56
Table 2.3: Naturally occurring primary metabolites, reported from genus	
Taverniera	62
Table 2.4: Phenolic acid and Lignans reported from genus Taverniera	67
Table 2.5: Naturally occurring secondary metabolites, reported from genus	
Taverniera	68
Table 2.6: Terpenoids reported from genus Taverniera	71
Table 2.7: Sterols reported from genus Taverniera	73
Table 2.8: Summary of some reported pharmacological studies on Taverniera	
genus	78

# Chapter III

Table Number	Page No.
Table 3.13: Instrument parameters for Method Validation	98
Table 3.14: Gradient Mobile Phase	98
Table 15.3: LC data acquisition parameters	101
Table 3.16: MS data acquisition parameters	102
Table 3.17: Parameters for Liquid Chromatography analysis	104
Table 3.18: Parameters for QToF instrument	104

### Chapter IV

Table Number	Page No.
Table 4.1: Different biogeographical zones with GPS location	109
Table 4.2: Morphological & Phytochemical observations of different locations	112
Table 4.3: Organoleptic characters of powders of <i>T. cuneifolia</i> roots	112
Table 4.4: Geographical parameters of Gujarat	114
Table 4.5: Macro and micronutrient analysis of Soil	114
Table 4.6: Identification of various sugar markers in the sample <i>T. cuneifolia</i>	
and G. glabra	128
Table 4.7: The chromatographic and calibration parameters, LOD and LOQ	132
Table 4.8: Precision of the method (intra-day)	132
Table 4.9: Precision of the method (inter-day)	133
Table 4.10: Recovery of glucose S8	133
Table 4.11: Recovery of fructose S8	133
Table 4.12: Recovery of sucrose S8	134
Table 4.13: Recovery of sucrose in Rajkot sample	134
Table 4.14: Recovery of fructose in Rajkot sample	135
Table 4.15: Recovery of sucrose in Kutch sample	135
Table 4.16: Results from the quantitative analysis of sugars detected in <i>T</i> .	
cuneifolia (Bagodara)	135
Table 4.17: Results from the quantitative analysis of sugars detected in <i>T</i> .	
cuneifolia (Rajkot)	136
Table 4.18: Results from the quantitative analysis of sugars detected in T.	426
<i>cuneifolia</i> (Rajkot)  Table 4.19: Identification of amino acid in <i>T. cuneifolia</i> in comparison with <i>G</i> .	136
glabra	141
Table 4.20: The chromatographic and calibration parameters, LOD and LOQ	143
Table 4.21: Precision of the method (intra-day)	144
Table 4.22: Precision of the method (inter-day)	144
Table 4.23: Recovery of Proline & Arginine	144
Table 4.24: Results from the quantitative analysis of amino acid detected in <i>T</i> .	144
cuneifolia and G. glabra	144
Table 4.25: Parameters of Glycyrrhizin	150
Table 4.26: Applicability of the developed method for the determination of	
Glycyrrhizin in Glycyrrhiza glabra (GG) and Taverniera cuneifolia (TC) sample	151
Table 4.27: Summary of some reported phytochemical studies on Apigenin	
estimation using HPLC	153
Table 4.28: Summary of some reported phytochemical studies on Glabridin	450
estimation using HPLC	153

Figure 4.29: Summary of some reported phytochemical studies on Genistein	
estimation using HPLC	154
Table 4.30: Summary of some reported phytochemical studies on Glycyrrhizin	
estimation using HPLC	154
Table 4.31: Summary of some reported phytochemical studies on Kaempferol	
estimation using HPLC	154
Table 4.32: Summary of some reported phytochemical studies on Liquiritigenin	
estimation using HPLC	155
Table 4.33: Summary of some reported phytochemical studies on Naringenin	
estimation using HPLC	155
Table 4.34: Summary of some reported phytochemical studies on Quercetin	
estimation using HPLC	155
Table 4.35: Summary of some reported phytochemical studies on Stigmasterol	
estimation using HPLC	157
Table 4.36: Summary of some reported phytochemical studies on β- sitosterol	
estimation using HPLC	157
Table 4.37: Identification result of selected standards in plant sample	158
Table 4.38: Results of system suitability of HPLC methodology for	
simultaneous method development of standards liquiritigenin, apigenin,	
genistein, glabridin and glycyrrhizin	163
Table 4.39: Results of system suitability of HPLC methodology for	
simultaneous method development of standards β-sitosterol and stigmasterol	163
Table 4.40: The chromatographic and calibration parameters, LOD and LOQ	164
Table 4.41: Recovery of quality control samples of standards from <i>G. glabra</i> via	104
HPLC techniques	168
Table 4.42: Recovery of quality control samples of standards from <i>T. cuneifolia</i>	100
chloroform extract via HPLC techniques	169
Table 4.43: Recovery of quality control samples of standards from <i>T. cuneifolia</i>	
methanol extract via HPLC techniques	170
Table 4.44: Results of intermediate precision of standards by HPLC technique	1,0
(Day 1,	171
Day 2 & Day 3)	171
Table 4.45: Results of intraday precision of standards by HPLC technique	172
Table 4.46: Results from the quantitative analysis of phytocomponents detected	470
in T. cuneifolia & G. glabra	173
Table 4.47: Characterization of standard compounds by LC-MS/MS-Q-ToF	174
Table 4.48: Result of the characterisation of aforementioned compounds in the	
plant samples	177
Table 4.49: Gradient solvent system used in column-chromatography for the	
	180
isolation of compounds from extract.	100
isolation of compounds from extract.  Table 4.50: List of saturated fatty acids identified in GCMS analysis of column	
isolation of compounds from extract.  Table 4.50: List of saturated fatty acids identified in GCMS analysis of column fraction, leaf oil, leaf oil residue, seed oil, and seed oil residue	183
isolation of compounds from extract.  Table 4.50: List of saturated fatty acids identified in GCMS analysis of column fraction, leaf oil, leaf oil residue, seed oil, and seed oil residue  Table 4.51: List of unsaturated fatty acids identified in GCMS analysis of	183
isolation of compounds from extract.  Table 4.50: List of saturated fatty acids identified in GCMS analysis of column fraction, leaf oil, leaf oil residue, seed oil, and seed oil residue  Table 4.51: List of unsaturated fatty acids identified in GCMS analysis of column fraction, leaf oil, leaf oil residue, seed oil, and seed oil residue	
isolation of compounds from extract.  Table 4.50: List of saturated fatty acids identified in GCMS analysis of column fraction, leaf oil, leaf oil residue, seed oil, and seed oil residue  Table 4.51: List of unsaturated fatty acids identified in GCMS analysis of	183
isolation of compounds from extract.  Table 4.50: List of saturated fatty acids identified in GCMS analysis of column fraction, leaf oil, leaf oil residue, seed oil, and seed oil residue  Table 4.51: List of unsaturated fatty acids identified in GCMS analysis of column fraction, leaf oil, leaf oil residue, seed oil, and seed oil residue	183 184
isolation of compounds from extract.  Table 4.50: List of saturated fatty acids identified in GCMS analysis of column fraction, leaf oil, leaf oil residue, seed oil, and seed oil residue  Table 4.51: List of unsaturated fatty acids identified in GCMS analysis of column fraction, leaf oil, leaf oil residue, seed oil, and seed oil residue  Table 4.52: Characterization of standard compounds by LC-MS/MS-Q-ToF	183 184 190

	1
Table 4.56: The Phenols identified through LC-MS-Q-ToF analysis	202
Table 4.57: The Terpenol glycosides identified through LC-MS-Q-ToF analysis	203
Table 4.58: The Alkaloids identified through LC-MS-Q-ToF analysis	203
able 4.59: The Terpenes identified through LC-MS-Q-ToF analysis	204
Table 4.60: The Sterols identified through LC-MS-Q-ToF analysis	206