

List of Figures

Figure 2.1: A schematic illustration of the concept of pseudopotential.	31
Figure 3.1: Top (a) and side view (b) of α -SiX (X=N, P, As, Sb, Bi) monolayers.	43
Figure 3.2: Band structure and PDOS of α -SiN (a and b), α -SiP (c and d), α -SiAs (e and f), α -SiSb (g and h) and α -SiBi (i and j) monolayers, respectively.	45
Figure 3.3: Structural geometry (a), band structure (b) and PDOS (c) of HGY monolayer, respectively.	46
Figure 3.4: Structural geometry of Ni@ α -SiN (a), Ni@ α -SiP (b), Ni@ α -SiAs (c), Ni@ α -SiSb (d) and Ni@ α -SiBi (e) SACs. The green sphere represents the Ni atom.	49
Figure 3.5: Structural geometry of Zr@HGY (a), Y@HGY (b), V@HGY (c), Sc@HGY (d), Mn@HGY (e), Co@HGY (f), Fe@HGY (g) and Cr@HGY (h) SACs, respectively.	50
Figure 3.6: Band structure of Ni@ α -SiN (a), Ni@ α -SiP (b), Ni@ α -SiAs (c), Ni@ α -SiSb (d) and Ni@ α -SiBi (e) SACs, respectively.	52
Figure 3.7: Band structure of Zr@HGY (a), Y@HGY (b), V@HGY (c), Sc@HGY (d), Mn@HGY (e), Co@HGY (f), Fe@HGY (g) and Cr@HGY (h) SACs, respectively.	55
Figure 3.8: PDOS of Ni@ α -SiN (a), Ni@ α -SiP (b), Ni@ α -SiAs (c), Ni@ α -SiSb (d) and Ni@ α -SiBi (e) SACs, respectively.	56
Figure 3.9: PDOS analysis of Zr@HGY (a, b), Y@HGY (c, d), V@HGY (e, f), Sc@HGY (g, h), Mn@HGY (i, j), Co@HGY (k, l), Fe@HGY (m, n) and Cr@HGY (o, p) SACs, respectively.	59
Figure 3.10: Structural geometry of the H adsorbed over Ni@ α -SiN (a), Ni@ α -SiP (b), Ni@ α -SiAs (c), Ni@ α -SiSb (d) and Ni@ α -SiBi (e) SACs, respectively.	62
Figure 3.11: PDOS and band structure of the H adsorbed over Ni@ α -SiN (a), Ni@ α -SiP (b), Ni@ α -SiAs (c), Ni@ α -SiSb (d) and Ni@ α -SiBi (e) SACs, respectively.	63
Figure 3.12: Structural geometry of the H adsorbed over Zr@HGY (a), Y@HGY (b), V@HGY (c), Sc@HGY (d), Mn@HGY (e), Co@HGY (f), Fe@HGY (g) and Cr@HGY (h) SACs, respectively.	66
Figure 3.13: PDOS and band structure of the H adsorbed over Zr@HGY (a1-a4), Y@HGY (b1-b4), V@HGY (c1-c4), Sc@HGY (d1-d4), Mn@HGY (e1-e4), Co@HGY (f1-f4), Fe@HGY (g1-g4) and Cr@HGY (h1-h4) SACs, respectively.	67
Figure 3.14: Structural geometry of Co@HGY (a), Fe@HGY (b), Cr@HGY (c) and Ni@ α -SiN (d) SACs at 0 K and 300 K, respectively.	71

Figure 3.15: The reaction coordinates plot for HER activity of pristine α -SiX (X=N, P, As, Sb, Bi), Ni anchored α -SiX (X=N, P, As, Sb, Bi) SACs, pristine HGY and transition metals (Zr, Y, V, Sc, Mn, Co, Fe, Cr) anchored HGY SACs, respectively.	72
Figure 4.1: The structural geometry (a), band structure (b), PDOS (c) and charge density contour plot (d) of 2D o-B ₂ N ₂	79
Figure 4.2: The structural geometry of B (a), N (b) and BN (c) defected 2D o-B ₂ N ₂ , respectively.	81
Figure 4.3: Band structure and PDOS of the B (a and d), N (b and e) and BN (c and f) defected 2D o-B ₂ N ₂ , respectively.	82
Figure 4.4: The structural geometry of the H adsorbed over B defected (a), N defected (b) and BN defected (c) 2D o-B ₂ N ₂ , respectively.	84
Figure 4.5: PDOS and band structure of the H adsorbed over B defected (a and b), N defected (c and d) and BN defected (e and f) 2D o-B ₂ N ₂ , respectively.	84
Figure 4.6: A charge density difference illustrations of the H adsorbed over BN defected (a), C doped at B site (b) and C doped at N site (c) of the 2D o-B ₂ N ₂ , respectively.	88
Figure 4.7: The structural geometry of C doped at B site (a) and at N site (b) of 2D o-B ₂ N ₂	89
Figure 4.8: PDOS and band structure of the C doped at B site (a and b) and at N site (c and d) of 2D o-B ₂ N ₂ , respectively.	89
Figure 4.9: The structural geometry of the H adsorbed over C doped at B site (a) and at N site (b) of 2D o-B ₂ N ₂ , respectively.	91
Figure 4.10: PDOS and band structure of the H adsorbed over C doped at B site (a and b) and at N site (c and d) of 2D o-B ₂ N ₂ , respectively.	91
Figure 4.11: The comparison of the structure plots at 0 K and 300 K of BN defected (a and b), C doped at B (c and d) and N sites (e and f) of the 2D o-B ₂ N ₂ , respectively.	95
Figure 4.12: Reaction coordinates of pristine, defected and C doped 2D o-B ₂ N ₂ for HER. ...	96
Figure 5.1: The structural geometry (a) and band structure (b) of the 2D o-B ₂ N ₂	103
Figure 5.2: The structural geometry of the Na (a), K (b), Be (c), Mg (d) and Ca (e) decorated 2D o-B ₂ N ₂ , respectively.	106
Figure 5.3: The structural geometry of the B (a), N (b) and BN (c) defected 2D o-B ₂ N ₂ , respectively.	107
Figure 5.4: The optimized geometry of the Sc (a), Ti (b), Y (c) and Zr (e) decorated 2D o-B ₂ N ₂ , respectively.	109

Figure 5.5: Band structure (a), charge density difference plot (b) and PDOS (e and f) of Ti decorated 2D o-B ₂ N ₂ ; PDOS of the pristine 2D o-B ₂ N ₂ (c) and isolated Ti atom (d), respectively.	110
Figure 5.6: Band structure (a), PDOS (b) and charge density difference plot (c) of the hydrogen molecule adsorbed over Ti decorated 2D o-B ₂ N ₂ , respectively.	112
Figure 5.7: The structural geometry of one (a), two (b), five (c), ten (d) and fourteen (e) hydrogen molecules adsorbed Ti decorated 2D o-B ₂ N ₂ , respectively.	114
Figure 5.8: The structural geometry of two Ti decorated 2D o-B ₂ N ₂ (a); and six (b), eighteen (c) and thirty-four (d) hydrogen molecules adsorbed on two Ti decorated 2D o-B ₂ N ₂ , respectively.	115
Figure 5.9: The results of AIMD simulation: The energy vs time step plot (a), top and side view of structural geometry (b) and variation in bond distances (Ti1-B (c) and Ti2-B (d)) plots of 2Ti/o-B ₂ N ₂ at 396 K, respectively.....	117