

REFERENCES

- *Alexopoulos CJ and Beneke ES (1955) Laboratory manual for introductory mycology. Bourgess Pub. Co. Minneapolis. 3.
- *Ivanova M and Petrova K (1995) Somatic embryogenesis and regeneration from traditional and mutant lines of maize on nutrient medium with NaCl. Rasteniev' dni-Nauki 32: 124-126.
- *Reinert J (1958) Morphogenese and ihre kontrolle an Gewebekulturen and karotten-Naturwissen chaften 45: 344-345.
- *Uddin MR, Dinus RJ, Webb DT (1990) Interactions of carbohydrates and Abscisic acid on maturation of *Pinus taeda* somatic embryos, Presented at 5th Conifer Biotechnology Working Group, Sittingbourne UK.
- Abe T and Futsuhara Y (1986) Genotypic variability for callus formation and plant regeneration in rice (*Oryza sativa* L.). Theor. Appl. Genet. 72: 3-10.
- Abe T and Futsuhara Y (1991) Diallel analysis of callus growth and plant regeneration in rice. Rice Genetic II. International Rice Research Institute, Manila, Phillippines.
- Abrol IP (1986) Salt affected soils: An overview. In VL Chopra, RS Paroda, eds

 Approaches for incorporating drought and salinity resistance in crop plants,

 Oxford and IBH Publishing Company, New Delhi, pp 1-23.
- Alia and Paradhasaradhi P (1993) Suppression in mitochondrial electron transport is the prime cause behind stress induced proline accumulation. Biochem. Biophys. Res. Comm. 193: 54-58.

- Ammirato PV (1983) Embryogenesis. In: DA Evans, WR Sharp, PV Ammirato and Y Yoshida (Eds.) Handbook of Plant Cell Culture, Mac Millan, New York Vol. 1, pp. 82-123.
- Ammirato PV and Steward FC (1971) Some effects of the environment on the development of embryos from cultured free cells. Bot. Gaz. 132: 149-158.
- Anbazhagan RV and Ganapathy A (1999) Somatic embryogenesis in cell suspension cultures of pigeon pea (*Cajanus cajan*). Plant Cell. Tiss Org. Cuture 56: 179-184.
- Apelbaum A and Canallakis ZN, Applewhite, Kaur-Sawhney R, Galston AW (1988)

 Binding of Spermidine to a unique protein in thin-layer tobacco tissue culture.

 Plant. Physiol. 88: 996-998.
- Aribaud M, Kevers C, Tanguy MJ, Gaspar T (1999) Low activity of amine-oxidase and accumulation of conjugated polyamines in disfavour of organogenic programmes in *Chrysanthemum* leaf disc explants. Plant Cell. Tiss. Org. Cult. 55: 85-94.
- Armstrong CL and Green CE (1985) Establishment and maintenance of friable embryogenic maize callus and the involvement of L-proline. Planta 164: 207-214.
- Attree SM and Fowke LC (1993) Somatic embryogenesis and synthetic seeds of conifers. Plant. Cell. Tiss. Org. Cult.35: 1-35.
- Attree SM, Dunstan DI, Fowke LC (1991a) Biotechnology in Agriculture and Forestry.

 Trees III. Vol 16 pp 423-445.
- Avjioglu A and Knox RD (1989) Storage lipids accumulation of zygotic and somatic embryos in culture. Ann. Bot. 63: 409-420.

- Bagni N and Mengoli M (1985) Characterization of carrot callus line resistant to high concentrations of putrescine. Plant Growth Reg. 3: 371-380.
- Bapat VA and Rao PS (1988) Sandal wood plantlets from Synthetic seeds. Plant Cell Rep 7: 434-436.
- Bargawa S and Chandra N (1989) Effect of sodium chloride on tolerant callus cultures of Moth bean (*Vigna aconitifolia* Jacq) Marechal CVIPCMO 926. Ind. J. Expt. Biol 27: 83-84.
- Barwale BR (1986) Food security in India In Hybrid Rice MacMillan India Ltd., Madras, pp 2-5.
- Basu S, Gangopadhyay G, Gupta S, Mukherjee BB (1996) Screening for cross tolerance against related osmotic stress in adapted calli of salt sensitive and tolerant varieties of rice. Phytomorphology 46(4): 357-364.
- Basu S, Gangopadhyay G, Mukherjee BB (1997) Isoenzyme of peroxidase and esterases as osmotic-stress markers in rice callus cultures. Ind. J. Expt. Biol. 35: 1359-1364.
- Basu S, Gangopadhyay G, Mukherjee BB, Gupta S (1997a) Plant regeneration of salt adapted callus of indica rice (Var. Basmati 370) in saline conditions. Plant. Cell. Tiss.Org. Cult. 50: 153-159.
- Bates LS, Waldren RP, Teare ID (1973) Rapid determination of free proline for water stress studies. Plant and Soil 39: 205-207.
- Ben Amer MI and Boner A (1997) The relationship between green spot initiation and plantlet regeneration of wheat callus grown under short term conditions. Plant Cell. Tiss. Org. Cult. 50: 67-69.
- Bernfeld P (1955) Amylases, α and β ; Meth. Enzym. 1: 149-155.

- Bernier G, Havelange A, Houssa C, Petitjean A, Lejeune P (1993) Physiological signals that induce flowering. Plant Cell 5: 1147-1155.
- Bertossi AF, Noirot M, Duval Y (1999) BA enhances the germination of oil palm somatic embryos derived from embryogenic suspension cutlures. Plant Cell. Tiss. Org. Cult. 56: 53-57.
- Bhaskaran S and Smith RH (1990) Regeneration in cereal tissue culture: a review.

 Crop. Sci. 30: 1328-1337.
- Bhaskaran S, Smith RH, Schertz K (1983) Sodium chloride tolerant callus of *Sorghum bicolor* (L) moench. Z. Planzenphysiol. 112: 459-463.
- Bhattacharaya S. (1991) Differential responses of rice (*Oryza sativa* L.) callus derived from three cultivars on saline environment. Phytomorphology 41: 138-137.
- Binh DQ, Heszky LE, Gyulai G, Csillag A (1992) Plant regeneration of NaCl pretreated cells from long-term suspension culture of rice (*Oryza sativa* L.) in high saline conditions. Plant. Cell. Tiss. Org. Cult. 29: 75-82.
- Binzel LM and Hasegawa MP (1987) Solute accumulation in tobacco cells adapted to NaCl. Plant. Physiol. 84: 1408-1415.
- Bradford MM (1976) A rapid and sensitive method of quantification of microgram quantities of protein utilizing the principle of protein dye binding.

 Anal.Biochem. 72: 248.
- Brady CJ, Gibson TS, Barlow EWR, Spiers J, Wyn Jones RG (1984) Salt tolerance in Plants. I. Ions, Compatible solutes and stability of plant ribosomes. Plant Cell. Environ. 7: 571-578.
- Bray E (1993) Molecular responses to water defecit. Plant . Physiol. 103: 1035-1040.

- Brettel RIS, Wirnickle W, Thames E (1980) Embryogenesis from cultured immature inflorescence of *Sorghum bicolor*. Protoplasma 104: 141-148.
- Bright SWJ and Jones MGK (1985) Advances in Agricultural Biotechnology. *In*: Cereal tissue and cell culture. Martinus Nijhoff (eds) Dr. W. Jurk Publishers pp 175-180.
- Bronsema FBF, van Oostreen FJW, van Lammeren MAA (1997) Comparative analysis of callus formation and regeneration on cultured maize embryos of the inbred lines A188 and A632. Plant Cell. Tiss. Org. Cult. 50: 57-65.
- Brown C, Brooks FJ, Pearson D, Mathias JR (1989) Control of embryogenesis and organogenesis in immature wheat embryo callus using increased medium osmolarity by abscic acid. J. Plant. Physiol. 133: 727-733.
- Brown DCW and Thorpe TA (1995) Crop improvement through tissue culture.

 W.J.Micro, Biol. Biotech. 11: 409-415.
- Burtin D, Tanguy JM, Paynot M, Rossini N (1989) Effects of the suicide inhibitors of arginine and ornithine decarboxylase activities on organogenesis, growth, free polyamines and hydroxylcinnamoyl putrescine levels in leaf explants of *Nicotiana xanthi* n.c. cultivated *in vitro* in a medium producing callus formation. Plant. Physiol. 89: 104-110.
- Cano EA, Perez-Alfocea F, Moreno V, Caro M, Bolarin MC (1998) Evaluation of salt tolerance in cultivated and wild tomato spices through *in vitro* shoot apex culture. Plant Cell. Tiss. Org. Cult. 53: 19-26.
- Centeno LM, Fernandez B, Feito I, Rodriguez A (1999) Uptake, distribution and Metabolism of 1-Napthaleneacetic acid and Indole-3-Acetic Acid during callus Initiation from *Actimidia deliciosa* Tissues. J. Plant. Growth. Reg. 18: 81-88.

- Chapman VJ (1975) The salinity problem in general, its importance and distribution with special reference to natural halophytes. In A Polijakoff Mayber, J Gale (eds.) Plants in saline environments. Springer-Verlag, Berlin, Heidelberg, New York, pp 7-24.
- Chauhan VA (1996) Isolation of hydroxyproline resistant cell lines of rice. Ph.D. thesis submitted to M.S. University of Baroda.
- Chauhan VA and Prathapasenan G (1998) Rice callus growth, proline content and activity of proline and IAA oxidase under the influence of hydroxyproline and NaCl. Acta. Physiol. Plant. 20(2): 197-200.
- Chauhan VA and Prathapasenan G (1999) Enzymatic studies in salt tolerant and salt sensitive rice cultivars under the influence of hydroxyproline and NaCl. Acta. Agr. Hung 47(2): 117-125.
- Chen TH, Lan L, Chen SS (1988) Somatic embryogenesis and plant regeneration from cultured young infloresence of *Oryza sativa* L. (Rice). Plant Cell. Tiss. Org. Cult. 4: 51-54.
- Choi EY, Yang CCD, Choi TK (1998) Induction of somatic embryos by macrosalt from mature zygotic embryos of *Panax ginseng*. Plant. Cell. Tiss. Org. Cult. 52: 177, 181.
- Choi Y, Kim WJ, Yoon SE (1999) High Frequency Plant Production via somatic embryogenesis from callus or cell suspension cultures in *Eleutherococcus senticosus*. Ann. Bot. 83: 309-314.
- Choudhary MT, Wainwright SJ, Merret MJ, Alam M (1994) Salt tolerant plants of Lucerne (*Medicago media* Pers.) regenerated from salt selected suspension cultures. Pl. Sci. 98: 97-102.

- Chowdhary NC, Tyagi AK, Maheswari N, Maheswari CS (1993) Effect of L-Proline and L-Tryptophan on somatic embryogenesis and plantlet regeneration of rice (*Oryza sativa* L V. Pusa 169). Plant Cell Tiss. Org. Cult. 32: 357-361.
- Christianson ML (1987) In Green CE, Sommers DA, Hackett WP and Biesboer DD (Eds) Plant Tissue and cell culture. Proc. Vth Instl. Cong. Plant Tissue and Cell Culture, Minnesota, Alan R. Liss. New York, pp 45-66.
- Christianson ML and Warnick DA (1988) Organogenesis in vitro as a Developmental Process. Hortsci. 23: 518-519.
- Close TJ and Bray EA (1993). Plant responses to cellular dehydration during environmental stresses. Amer. Soc. Plant Physiol. Rockville, MD, 104-108.
- Cornu D and Geoffrion (1990) Euk. Soci. Bot. Fdr. 137: 25-34.
- Cushman JC, De Rocher EJ, Bohnert HJ (1990) Gene expression during adaptation to salt stress. In Environmental Injury to Plants (eds.) F. Katlerman, SanDiego, Academic Press, pp 173-203.
- Cvikrova M, Binarova P, Cenklova V, Eder J, Machackova I (1999) Reinitiation of cell division and polyamine and aromatic monoamine levels in alfalfa explants during induction of somatic embryogenesis. Physiol. Plant 105: 330-337.
- Deane RC, Fuller MP, Dix PJ (1995) Selection of hydroxyproline resistant proline accumulating mutants of cauliflower (*Brassica oleracea* var. botrytis) Euphytica 85: 329-334.
- Dhed'a D, Dumortier F, Panis B, Vuylsteke D, De Langhe E (1991) Plant regeneration in cell suspension cultures of the cooking banana (V Bluggoe (Musa spp. ABB group) Fruits 46: 125-135.

- Dix PJ (1986) Cell line selection. In Plant Cell Culture Technology, Yeomann (ed.),M.M. Blackwell Scientific Publications, Oxford, pp 143-201.
- Dix PJ and Street HE (1975) Sodium chloride resistant cultured cell lines from Nicotana sylvestris and Capsicum annum. Plant. Sci. Lett. 5: 231-237.
- Dix PJ, Mclysaght VA, Pearce RS (1984) The potential of cell cultures for the production of salt tolerant cultivars. In W Lange, AC Zeven, NG Hogenboom, (eds.), Proceedings 10th Congress of Eucarpia. Pudoc, Wageningen.
- Downton WJS (1984) Salt tolerance of food crops. Prospectives for improvements.

 CRC Critical Reviews in Plant Sciences 1: 183-201.
- Druart P and Dewulf O (1993) Activated charcol catalyses sucrose hydrolysis during autoclaving. Plant. Cell. Tiss. Org. Cult. 32: 97-99.
- Dubey RS and Rani M (1990) Influence of NaCl salinity on the behaviour of protease, aminopeptidase and carboxyl peptidase in rice seedlings in relation to salt stress.

 Aust. J. Plant. Physiol. 17: 215-224.
- Duncan DB (1955) Multiple range and Multiple F-tests. Biometrics 11: 1-42.
- Eapen S and Rao PS (1995) Plant regeneration from immature inflorescence callus cultures of wheat, rye and triticale. Euphytica 34: 153-159.
- Emons AMC, Samallo-Droppers A, Van Der Toorn C (1993) The influence of sucrose, mannitol, L-Proline, abscic acid and gibberallic acid on the maturation of somatic embryos of *Zea mays* L. from suspension cultures. J. Plant. Physiol. 142: 597-604.
- Emons and Kieft U (1991) Histological comparison of single somatic embryos of maize from suspension culture with somatic embryos attached to callus cells. Plant Cell. Rep. 10: 485-488.

- Epstein E (1980) Response of plants to saline environment. *In* DW Rains, RC Valentine, A Hollander, eds, Genetic engineering of osmoregulation: Impact on plant productivity for food, chemical and energy. Plenum Press, New York, pp. 7-21.
- Evans PT and Malmmberg RI (1989) Do polyamines have a role in plant development?

 Ann. Rev. Plant.Physiol. Plant Mol. Biol. 40: 235-269.
- Faure O, Dewitte W, Nougarede A, Oncklen VH (1998). Precociously germinating somatic embryos of *Vitis vinifera* have lower ABA and IAA levels than their germinating zygotic counterparts. Physiol. Plant 102: 591-595.
- Fienberg AA, Choi JH, Lubich WP, Sugh ZR (1984) Developmental regulation of polyamine metabolism in growth and metabolism of carrot culture. Planta 162: 532-539.
- Finkelstein RR. and Crouch ML. (1986) Rapeseed embryo development in culture on high osmoticum is similar to that in seeds. Plant Physiol. 81: 907-912.
- Flores HE and Galston AW (1982) Polyamines and plant stress: activation of putrescine biosynthesis by osmotic shock. Science: 217: 1259-1261.
- Flowers TJ and Yeo AR (1981) Variability in the resistance of sodium chloride salinity with rice (*Oryza sativa* L) varieties. New Phytol. 88: 295-373.
- Flowers TJ, Troke PF, Yeo AR (1977) The mechanism of salt tolerance in halophytes.

 Ann. Rev. Plant. Physiol. 28: 89-121.
- Fujii AJ, Slade D, Redenbaugh K (1993) Planting artificial seeds and somatic embryos.

 In Synthetic seed technology application of synthetic seeds to crop improvement

 CRC Press, Boca Raton, Fla, pp 183-202.

- Galiba G and Yamada Y (1988) A novel method for increasing the frequency of somatic embryogenesis in wheat tissue culture by NaCl and KCl supplementation. Plant Cell Rep. 7: 53-58.
- Galston AW (1983) Polyamines as modulators of plant development. Bioscience 33: 382-388.
- Galston AW, and Kaur-Sawhney R (1987) Polyamines as endogenous growth regulator.

 In PJ Davies, ed. Plant Hormones and Their Role in Plant Growth and Development Martinus Nijhoff, Boston, pp. 280-295.
- Gangopadhyay G, Basu S, Gupta S (1997) In vitro selection and physiological characterization of NaCl and mannitol adapted callus lines of Brassica juncea.

 Plant Cell Tiss. Org. Cult. 50: 161-169.
- Gangopadhyay G, Basu S, Mukherjee BB, Gupta S (1997) Effects of salt and osmotic shocks on unadapted and adapted callus lines of tobacco. Plant Cell. Tiss. Org. Cult. 49: 45-52.
- Garg BK and Gupta, K (1997) Saline Waste Land Environment and Plant Growth, Scientific Publishers, Jodhpur, India pp 242.
- Garg BK, Vyas SP, Kathju S, Lahiri AN, Meli PC, Sharma PC (1995) Salinity-fertility interaction on growth, mineral composition and nitrogen metabolism of Indian Mustard J. Plant. Nutr. 16: 1637-1650.
- Gartner A and Lorz H (1996) *In vitro* regeneration systems of barley (*Hordeum vulgare*L.) Plant Tiss. Cult. Biotech. 2: 11-23.
- Gasper TH, Kevers C, Hausman JF (1997a) Indissociable chief factors in the inductive phase of adventitious rooting. *In* Biology of root formation and development Altman & Y, Waisel (eds.), Plennum Press, New York, pp 55-64.

- Gasper TH, Kevers C. Penel C, Greppin H, Reid D, Thrope A (1996b) Plant hormones and plant growth regulators in plant tissue culture. *In vitro* cell. Dev. Biol. Plant 32: 272-289.
- George EF and Sherington FF (1984) Plant propagation by tissue culture. Handbook and directory of commercial laboratories. Exergetics Ltd, Eversley, Basingstoke, Hans, England.
- George L and Eapen S (1988) Plant regeneration from somatic embryogenesis from immature inflorescence culture of *Sorghum almus*. Ann. Bot. 61: 589-591.
- George L and Eapen S (1994) Organogenesis and embryogenesis from diverse explants in pigeon pea (*Cajanus cajan* L.) Plant Cell Rep. 13(7): 417-420.
- George L and Eapen S (1995) Encapsulation of somatic embryos of finger millet, *Eleusine coraceana* Gaertn Ind. J. Expt. Biol. 33(4): 291-293.
- George L, Eapen S, Rao PS (1989) High frequency somatic embryogenesis and plant regeneration from immature inflorescense cultures of two Indian cultivars of sorghum (Sorghum bicolor L. Moeneh) Proc. Indian Acad. Sci. 99(5) 405-410.
- Ghanem SA (1995) *In vitro* embryogenesis of lentil under saline conditions. Bull. Facult. Agr. Uni. Cairo 46: 113-125.
- Goebel-Tourand I, Mauro MC, Sossountzov L, Miginiac E, Deloire A (1993) Arrest of somatic embryo development in grapevine: Histological characterization and the effect of ABA, BAP and Zeatin in stimulating plantlet development. Plant Cell. Tiss. Org. Cult. 33: 91-103.
- Gordon SA and Weber RP (1951) Calorimetric estimation of indole acetic acid. Plant Physiol. 26: 192-195.

- Gossett RD, Millhollon, Lucas CM, Banks WS, Marney MM (1994) The effect of NaCl on antioxidant enzyme activities in callus tissue of salt-tolerant and salt sensitive cotton cultivars (*Gossypium hirsutum* L.) Plant Cell. Rep. 13: 498-503.
- Greenway H and Munns R (1980) Mechanism of salt tolerance in non-halophytes.

 Annu.Rev. Plant Physiol 31: 149-190.
- Grey DJ and Purohit A (1991) Somatic embryogenesis and development of synthetic seed technology. Crit. Rev. Plant. Sci. 10: 33-61.
- Gulati A and Jaiswal PK (1992) Comparative salt response of callus cultures of *Vigna* radiata (L) Wilezek to various osmotic and ionic stresses. J. Plant. Physiol. 14: 120-124.
- Guohua M (1998) Effects of cytokinins and ausins on Cassava shoot organogenesis and somatic embryogenesis from somatic embryo explants. Plant. Cell. Tiss. Org. Cult. 54: 1-7.
- Gupta US (1997) Cell and Tissue culture. In Crop Improvement Vol. 2, Stress Tolerance in plants. Oxford and IBH Publishing Company, New Delhi pp 18-24.
- Handa S, Handa AK, Hasegawa PM, Bressan RA (1986) Proline accumulation and adaptation of cultured plant cells to water stress. Plant Physiol 80: 938-945.
- Harrington HM and Alm DM (1988) Interaction of heat and salt shock in cultured tobacco cells. Plant. Physiol. 88: 611-625.
- Henke RR, Mansur A, Constantin JM (1978) Organogenesis and plantlet formation from organ-seedling derived calli of Rice (*Oryza satīva*) Physiol. Plant 44: 11-14.

- Holme BI, Krogstrup P, Hansen J (1997) Embryogenic callus formation, growth and regeneration in callus and suspension cultures of *Miscanthus ogiformis* Honda x Guganteus as affected by proline. Plant. Cell. Tiss. Org. Cult. 50: 203-210.
- Hutchison JM, Senaratna T, Tsujita, Saxena KP (1997) Somatic embryogenesis in liquid cultures of a tetraploid Alstroemeira. Plant Cell. Tiss. Org. Cult. 47: 293-297.
- International Rice Research Notes (1993) March, 18(1):58.
- Islam AS (1996) Plant Tissue Culture Studies in Cereal Biotechnology and Molecular Biology A progress report with special reference to Rice and Wheat, Oxford and IBH publishing company, New Delhi pp. 111-123.
- Jain RK (1997) Effect of some factors on plant regeneration from indica rice cells and protoplasts. A review Ind. J. Expt. Biol. 35: 323-331.
- Jain RK, Jain S, Choudhary JB (1991b) In vitro selection for salt tolerance in Brassica juncea L. using cotyledon explants callus and cell suspension cultures. Ann. Bot. 67: 517-519.
- Jain S, Nainawatee SH, Jain RK (1994) Photosynthetic response to NaCl salinity in in vitro salt tolerant somaclones and parent cv. prakash of Brassica junea Ind. J. Expt. Biol. 32: 588-593.
- Jana MM, Nadgauda SR, Rajmohan K, Mascarenhas FA (1994) Rapid somatic embryogenesis from nucelli of monoembryonic mango varieties. In Vitro Cell. Dev. Biol. 301: 55-57.
- Janeiro LV, Ballester A, Vieitez AM (1997) *In vitro* response of encapsulated somatic embryos of *Camellia*. Plant Cell. Tiss. Org. Cult. 51: 119-125.

- Jarvis BC, Shanon PRM, Yasmin S (1983) Involvement of polyamines with adventious root development in stem cuttings of mung bean. Plant Cell Physiol 24: 677-683.
- Jasrai YT, Bhatt PN, Mehta AR (1988) Endogeneous IAA content and epiphyllus bud regeneration in *Kalanchoe mortagei*, Abstract No. 8.33 "International Congress of Plant Physiology" IARI, New Delhi.
- Jia JF, Lu W, Zhao XS (1993). Selection of NaCl tolerant variant of Seteria italica via tissue culture technique and its physiological and biochemical characteristics. Proceedings of the First Asia-Pacific Conference in Agricultural Biotechnology, Beijing, China. In, Current Plant and Biotechnology in Agriculture (CB You, Zi Chen, YJ Ding, eds), Vol.15, Dordrecht, Netherlands, Kluwer Academic Publishers, pp. 309-312.
- John A and Prathapasenan G (1999) High frequency plantlet regeneration from root explants of rice (*Oryza sativa* L. var. CSR-10). Phytomorphology 49(2): 165-169.
- John A, Chauhan VA, Prathapasenan G (1997) Regeneration of plants from NaCl adapted callus cultures of rice. In proceedings of volume of National Symposium, S.P. University, Anand, Gujarat. Plant. Struct. Morph. 111-114.
- Kackar and Shekhawat SN (1991) Plant regeneration from cultured immature inflorescence of *Cenchrus setigerus* and *C. ciliaris*. Ind. J. Expt. Biol. 29: 62-64.
- Katiyar S and Chandel G (1998) High frequency somatic embryogenesis from immature Inflorescence of *Coix aquatica* Roxb. Plant. Tiss. Cult. 8(2): 131-137.

- Kaur- Sawhney R, Shih LM, Flore HE, Galston AW (1982) Relation of polyamine synthesis and titre to aging and senescence in oat leaves. Plant Physiol 69: 405-410.
- Kaur-Sawhney R, Dai YR, Galston AW (1985) Effect of inhibitors of polyamine biosynthesis on gibberellin-induced internode growth in light-grown dwarf peas.

 Plant Cell Physiol. 27: 253-260.
- Kavikishor PB and Mehta AR (1989) Carbohydrate oxidation during organogenesis in callus culture of Tobacco. Ind. J. Expt. Biol. 27: 124-127.
- Kavikishor PB, Sangam S, Naidu RK (1999) Sodium, Potassium, Sugar alcohol and proline mediated somatic embryogenesis and plant regeneration in recalcitrant rice callus. In: Plant Tissue Culture and Biotechnology: Emerging Trends. PB Kavikishor (ed) University Press Hyderabad, pp 78-85.
- Ketchum REB, Warren RC, Klima LJ, Lopez-Gutierrez F, Nabors MW (1991) The mechanism and regeneration of proline accumulation in suspension cultures of halophytic grass *Distichlis spicata* L. J. Plant. Physiol. 137: 368-374.
- Khush GS (1997) Origin, dispersal, cultivation and variation of rice. Plant Mol. Biol. 35: 25-34.
- Kim YH and Janick J (1991) Abscisic acid and proline improve desication tolerance and increase fatty acid content of celery somatic embryogenesis. Plant Cell. Tiss. Org. Cult. 24: 83-89.
- Kinizios SE, Barberaki M, Aivalikis G, Drossopoulos J, Holevas DC (1997) *In vitro* morphogenetic response of wheat embryos to different NaCl concentrations and growth regulator treatment. Plant. Breed 116: 113-118.

- Lambe P, Hity SN, Mutambel, Deltour R, Duinant M (1999) Somatic embryogenesis in pearl millet (*Pennisetum glaucum*): strategies to reduce genotype limitation and to maintain long-term totipotency. Plant Cell. Tiss. Org. Cult. 55: 23-29.
- Lebrun L, Rajasekaran K, Mullins MG (1985) Selection *in vitro* for NaCl-tolerance in *Vitts rupestris* Scheele. Ann. Bot. 56: 733-739.
- Lee CW and Thomas JC (1985) Jojoba embryo culture and oil production. Hortsci. 20: 762-764.
- Leone A, Costa A, Tucci M, Grillo S (1994) Adaptation versus shock response to polyethylene glycol-induced low water potential in cultured potato cells. Physiol. Plant 92: 21-30.
- Levitt J (1980) Responses of plants to environmental stresses. Vol. II Water, Radiation, salt and other stresses. Academy Press, New York pp. 607.
- Li HJ (1990) Study on selection of salt resistant variants of rice. Journal of Shenyang Agricultural University 21 (1): 53-59.
- Linsmaeir EM and Skoog F (1965) Organic growth factor requirements of tobacco tissue culture Physiol. Plant 18: 100-127.
- Litz RE and Gray DJ (1995) Somatic embryogenesis for agricultural improvement W.J. Micro. Biotech. 11: 416-425.
- Litz RE. and Gray DJ. (1992) Organogenesis and somatic embryogenesis. *In*Biotechnology of Perennial Fruit crops, Hammerschlag FA. & Litz RE (eds)

 CAB Int., Wallingfora, UK, pp. 3-34.
- Lu W and Jia JF (1994) Selection of a salt tolerant cell line from embyogenic callus of millet and studies on its physiological and biochemical characteristics. Acta Agro. Sini. 20 (2): 241-247.

- Lusardi MC, Locatelli F, Stadler J, Lupotto E. (1991) *In vitro* characterization of salt-selected maize genotypes. J. Gen. Breed. 45: 285-291.
- Lutt S, Kinet JM, Bouharmont J (1999) Improvement of rice callus regeneration in the presence of NaCl. Plant. Cell. Tiss. Org. Cult. 57: 3-11.
- Lutts S, Kinet JM, Bouharmont J (1996) Effect of salt stress on growth, mineral nutrition and proline accumulation in relation to osmotic adjustment in rice (*Oryza sativa* L.) cultivars differing in salinity resistance. Plant Growth Reg. 19: 207-218.
- Maeda E and Saka H (1973) Light microscopy of cell organelles in the shooting rice callus tissue. Plant Physiol. 37: 51-58.
- Malfatti H, Vallee JC, Perdrizet E, Carre M, Martin C (1983) Acids amines libres D. explantes foliares de *Nicotiana tobaccum*, cultivars *in vitro* sur des milieux induisant la thizogenese ou la caulogenese. Physiol. Plant 57: 492-498.
- Manetas Y, Petropoulu Y, Karabourniotis G (1986) Compactable solutes and their effects on phosphoenol pyruvate carboxylase of C₄-halophytes. Plant. Cell. Environ. 9: 145-151.
- Martin-Tanguy J, Aribaund M, Carre M, Gasper TH (1997) ODC mediated biosynthesis and DAO-mediated catabolism of Putrescine involved rooting by leaf explants of *Chrysanthemum morifolium* Ramat *in vitro*. Plant. Physiol. Biochem. 35: 595-602.
- Mascarenhas AF, Pathak M, Hendre RR, Jaganathan V (1975). Tissue cultures of maize, wheat rice and sorghum: Part I Initiation of viable callus and root culture. Ind. J. Expt. Biol. 13: 103-107.

- Mathur AK, Ganapathy PS, Johri BM (1980) Isolation of sodium-chloride-tolerant plantlets of *Kiockxia ramisissima* under *in vitro* conditions. Z. Pflanzenphysiol 99: 287-294.
- Mathur KC, Reddy PR, Rajamani S, Moorthy STB (1999). Integrated pest management in rice for improve productivity and sustainability. Oryza 36(3): 195-207.
- McHughen and Swartz M (1984) A tissue culture raised salt-tolerant line of flax (*Linum usitassimum*) J. Plant. Physiol. 117: 109-117.
- Meredith CP (1984) Selecting better crops from cultured cells. In JP Gustafson, (eds),

 Gene manipulation in plant improvement 16th Stadler Genetics Symposium.

 New York, Plenum Press, pp. 503-528.
- Mohamand AS (1993) Tissue culture variability in wheat germplasm: callus initiation and long term plant regeneration and maintenance. Pak. J. Sci. Ind. Res. 36: 306-309.
- Mohanty KH, Mallik S, Grover A (2000) Prospects of improving flooding tolerance in low land rice varieties of conventional breeding and genetic engineering. Curr. Sci. 78(2): 132-137.
- Mooney H, Winner WE, Pell EJ (1991) In Response of plants to multiple stresses, (eds)

 Academic Press, San Diego.
- Muralidharan EM and Mascarenhas AF (1995) Somatic embryogenesis in Eucalyptus.

 In Somatic embryogenesis in woody plants. Vol 2, Angiosperm; Jain S, Gupta P,

 Newton R (eds). Kluwer Academic Publishers, Dordrecht, pp 23-40.
- Muralitharan MS, Van Steveninck RFM, Chandler SF (1990) Growth characteristics and ion contents of non-selected and salt-selected callus lines of high bush

- blueberry (*Vaccinium corymbosum*) cultivars Blue Crop and Denise Blue. Plant Cell Rep. 9: 151-155.
- Murashige T and Skoog F (1962) A revised medium for rapid growth and bioassay with tobacco tissue cultures. Physiol. Plant. 15: 473-497.
- Nabors MW, Daniels A, Nodolny L, Brown C (1975) Sodium chloride tolerant lines of tobacco cells. Plant. Sci. Lett. 4: 155-159.
- Nambisan P and Chopra VL (1992) Somatic embryogenesis in *Triticum aestivum* L. Morphological observations on germination. Ind. J. Exp. Biol. 30: 12-14.
- Nogaard JV and Krogstrup P (1990) Somatic embryogenesis of *Abies nordmanniana*LK controlled by cytokinin, presented at 5th Conifer Biotechnology Working

 Group, Sittingbourne, UK.
- Norlyn JD (1980) Breeding salt tolerant crop plants. *In* Genetic Engineering of Osmoregulation. Impact on Plant Productivity for Food, Chemical and Energy; D.W. Rains, R.C. Valentine and A. Hollander (eds.) Plennum Press, New York, pp 293-309.
- Norstog IK (1970) Induction of embryo like structures by kinetin in cultured Barley embryos. Dev. Biol. 23: 665-670.
- O'Neill MC, Arthur EA, Mathias JR (1996) The effects of proline, thioproline and methylglyoxalbis-(guanylhydrozone) on shoot regeneration frequencies from stem explants of *B. napus*. Plant Cell. Rep. 15: 695-698.
- Ochatt SJ, Marconi PL, Arnozis PA, Caro OH (1999) *In vitro* recurrent selection of potato: Production and characterization of salt tolerant cell lines and plants.

 Plant. Cell. Tiss. Org. Cult. 55: 1-18.

- Ozawa K and Komamine A (1988) Establishment of a system of high-frequency embryogenesis from long-term cell suspension cultures of rice (*Oryza sativa* L.).

 Theor. Appl. Genet 77: 205-211.
- Padmaja G, Reddy VD, Reddy GM (1992) Somatic embryogenesis and plant regeneration from mature embryo callus cultures of triticale. Ind. J. Expt. Biol. 30: 181-184.
- Paleg LG, Douglas TJ, van Daal A, Keech D (1981). Proline, betaine and other organic solutes protects enzymes against heat inactivation. Aust. J. Plant. Physiol. 8: 107-114.
- Paleg LG, Steward GR, Bradbeer JW (1984) Proline and glycine betaine influence on protein solvation., Plant Physiol 75: 974-978.
- Pandey R and Ganapathy PS (1985). The proline enigma: NaCl-tolerant and NaCl-sensitive callus lines of *Cicer arietinum* (L.). Plant Sci. 40: 13-17.
- Pardhasaradhi P, Alia, Vani B (1993) Inhibition of mitochondrial electron transport is the prime cause behind proline accumulation during mineral deficiency in *Oryza sativa*. Plant and Soil 155/156: 465-468.
- Pareek A, Singla S, Grover A (1998) Proteins alterations associated with salinity, dessication, high and low temperature stresses and abscisic acid application in seedlings of Pusa 169, a high yielding rice (*Oryza sativa* L.) cultivar. Curr. Sci. 75(10):1023-1035.
- Pessarakli M. (1994) Hand book of plant and crop stresses. Marcel Dekker, Inc. New York pp. 697.
- Piqueras A and Hellin E (1992) Selection and characterization of NaCl-tolerant cell line of *Citrus limonum*. Suelo. Y. Planta 2: 629-640.

- Piqueras A, Jose A, Hernandez, Olmos E, Hellin E, Sevilla F (1996) Changes in antioxidant enzymes and organic solutes associated with adaptation of citrus cells to salt stress. Plant Cell. Tiss. Org Cult. 45: 58-60.
- Prakash NS and Sarin NB (1993) Isolation of salt tolerant cell lines of *Cajanus cajan* and comparative study of their physiological responses with intact plants under salt stress. Plant Physiol. Biochem. 20(1): 11-15.
- Pua EC and Thorpe TA (1986). Differential response of non-selected and Na₂SO₄ selected callus cultures of *Beta vulgaris* L. to salt stress. J. Plant Physiol. 123: 241-248.
- Purohit SD and Kukda G (1996) Somatic embryogenesis and plantlet regeneration in Heteropogon contortus (L) P. Beavu. J. Ind. Bot. Soc. 75: 145-146.
- Purohit SD, Kukda G, Sekhawat R (1992) Plant regeneration in *Apluda mutica* L. through somatic embryogenesis. Ind. J. Expt. Biol. 30: 1154-1157.
- Radojevic L (1995) Somatic embryogenesis in woody plants. *In Jains S*, Gupta P and Newton R (eds) Vol.2, Kluwer Academic Publishers, Netherlands, pp 409-444.
- Rains DW, Croughan SS, Croughan TP (1986). Isolation and characterization of mutant cell lines and plants: salt tolerance. *In* Cell Culture and somatic cell genetics of plants, Vol.3, IK Vasil, (ed.) FL, Academic Press, pp. 537-547.
- Rains DW, Croughan TP, Stavarek SJ (1980) Selection of salt tolerant plants using tissue culture, *In* Genetic Engineering of Osmoregulation. Impact on plant productivity for food, chemicals and energy. DW Rains, RC Valentine, A Hollaender (eds.) Plenum Press, New York, pp. 279-292.
- Rajam MV (1997) Polyamines, *In Plant Ecophysiology*, Prasad MNV (ed.) John Wiley and Sons, Inc., New York 343-374.

- Ramagopal S (1986) Protein synthesis in a maize callus exposed to NaCl and mannitol.

 Plant Cell Rep. 5: 430-434.
- Rana RS (1986) Breeding crop varieties for salt affected soils. In Approaches for incorporating Drought and Salinity Resistance in crop plants. VL Chopra and RS Paroda (eds.), Oxford and IBH Publishing Co., New Delhi pp. 24-55.
- Rangan TS and Vasil IK (1983) Sodium Chloride Tolerant Embryogenic Cell Lines of Pennisetum americanum (L.) K. Schum. Ann. Bot 52: 59-64.
- Rani UT and Reddy GM (1996) Regeneration of Indica rice (*Oryza sativa* L) cultivars from embryogenegic suspension culture. Ind. J. Expt. Biol. 34: 239-242.
- Rao GG, Polra NV, Babu RV, Girdhar KI (1999) Growth and development of Salvadora persica on highly saline black soils: Salt tolerance during immature phase. Ind. J. Plant. Physiol. 4(3): 152-156.
- Rao GG, Singh R, Barghava GP (1994) Species diversity on salt affected soils under canal command areas and in Bhal region in Gujarat State. Ind. Forest. 121: 1143-1150.
- Raval M and Chatoo BB (1993) Role of media constituents and proline in callus growth, somatic embryogenesis and regeneration of *Oryza sativa* cv indica. Ind. J. Expt. Biol. 31: 600-603.
- Reddy PJ and Vaidyanath K (1986) *In vitro* characterization of salt stress effects and the selection of salt tolerant plants in rice (*Oryza sativa* L.) Theor. Appl. Gent. 71: 757-760.
- Redenbaugh K (1993) Syn seeds: Applications of synthetic seeds to crop improvement CRC Press, Boca Raton.

- Redenbaugh K, Fujii JA, Slade D (1991) Synthetic seed technology. *In* Scale-up and Automation in Plant Propagation Cell Culture and Somatic Cell Genetics of Plants. Vol. 8, Vasil KI (ed.) Academic Press. Inc. New York, pp 35-74.
- Sabu A, Sheeja TE, Nambisan P (1995) Comparison of proline accumulation in callus and seedlings of two cultivars of *Oryza sativa* L. differing in salt tolerance. Ind. J. Expt. Biol. 33: 139-141.
- Sanho M, de Forchelti MS, Pilego F, Valpuesta V, Quesada M (1996) Peroxidase activity and isoemzymes in the culture medium of NaCl adapted tomato suspension cells. Plant Cell. Tiss. Org. Cult. 44: 161-167.
- Sankhla N and Upadhyaya A (1988) Polyamines and adventious root formation. *In*Adventious Root formation in Cuttings Davis TD, Haissig BE, Sankla N. (eds.)

 Discorides Press, Oregon, USA pp. 202-213.
- Santarius KA (1992) Freezing of isolated thylakoid membranes in complex media. VIII.

 Differential cryoprotection by sucrose, proline and glycerol. Physiol. Plant 84:

 87-93.
- Santoro MM, Liu Y, Khan SMA, Hou LX, Bolen DW (1992) Increased thermal stability of proteins in the presence of naturally occurring osmolytes. Biochemistry 31: 5278-5283.
- Santos MA, Camara T, Rodriguez P, Claparols I, Torne MJ (1996) Influence of exogeneous proline on embryogenic and organogenic maize callus subjected to salt stress. Plant. Cell. Tiss. Org. Cutlure 47: 59-65.
- Sengupta ND, Banerjee K, Rouf A (1999) Abiotic stress responsive Elements, their Binding Factors and their utilization for the Improvement of Tolerance to Abiotic stress in Plants. J. Plant Biol. 26(3): 179-185.

- Serrano R and Gaxiola R (1994) Microbial models and salt stress tolerance in plants.

 CRC Crit. Rev. Plant Sci. 13: 121-128.
- Shannon MC (1985) Principles and strategies on breeding for higher salt tolerance.

 Plant. Soil. 89: 587-589.
- Shanon MC (1982) Genetics of salt tolerance: New Challenges. *In* Biosaline research: A look to the future. A. San Pietro, (ed.). Plenum Press. New York, 271-278.
- Sharma SK and Gupta IC (1986) Saline Environment and Plant Growth. Agro Botanical Publishers, Bikaner. pp. 172.
- Sharp WR, Sondanl MR, Caldas LS, Maraffa SB (1980). The physiology of in vitro asexual embryogenesis. Horticultural Reviews. Vol. 2 Janick J. (ed.) AVI Publishing Co. Inc. Westport CN, Hort. Rev. 2: 268-310.
- Shayakhmetov IF (1996) Development of somatic embryios in wheat cell suspension culture in presence of ABA. J. Bul. Soc. Pl. Physio. 102(2): 130-133.
- Shevyakova NI., Ro shchupkin BV. Paramonova NV, Kuznetsov VV (1994) Stress responses in *Nicotiana sylvestris* cells to salinity and high temperature:

 Accumulation of proline, polyamines, betaines and sugars. Russian. J. Plant Physiol. 41: 490-496.
- Shi DC, Xi T, Zhu TC (1989). Somatic embryogenesis from mature embryo of tall wheatgrass (Agropyron elongatum) and effects of NaCl on induction and growth callus from its different explants. Proceedings of the XVI International Grassland Congress, Nice, France. 443-444.
- Shoemaker RC and Hammond EG (1988) Fatty acid composition in soybean (Glycene max. L.) somatic embryogenesis. In vitro. 24: 829-832.

- Singh NK, Handa AK, Hasegaura PM, Bressen RA (1985). Protein associated with adaptation of cultured tobacco cells to NaCl. Plant. Physiol. 79: 126-137.
- Singh RR, Kemp AJ, Kollmorgen FJ, Qureshi AJ, Fincher G (1997) Fertile plant regeneration from cell suspension and protoplast cultures of barley (*Hordeum vulgare* cv Schnoar) Plant Cell. Tiss. Org. Cult. 49: 121-127.
- Singha KM, Sachdev A, Johri RP, Mehta SL (1999) Stress induced Polypeptrides in Lathyrus sativus. J. Plant. Bioch. Biotech. 8: 47-51.
- Singhla SL, Pareek A, Grover A (1997) Plant Ecophysiology Prasad MNV(ed.), John Wiley and Sons, New York, 107-127.
- Skoog F and Miller CO (1957). Chemical; regulation of growth and organ formation.

 Symp. Soc. Exp. Biol. 11: 118-131.
- Smirnoff N (1995) Metabolic flexibility in relation to the environment. *In* Environment and Plant Metabolism flexibility and acclimation. Bios Scientific Publishers, Oxford OXI 1SJ-UK, pp 270.
- Smith RH and Bhaskaran S (1986) Sorghum. In Biotechnology in Agriculture and Forestry; YPS Bajaj (ed.) Springer-Verlag Berlin pp 220-233.
- Steward FC (1958) Growth and development of cultivated cells. III. Interpretation of the growth from free cells to carrot plant. Am. J. Bot. 45: 709-713.
- Steward FC, Mapes MO, Mears K (1958) Growth and organised development of cultured cells. II. Organization in cultures grown from freely suspended cells.

 Am. J. Bot. 45: 705-708.
- Subhashini K and Reddy GM (1990) Effect of salt stress on enzyme activities in callus cultures of tolerant and susceptible rice cultivars. Ind. J. Exp. Biol. 28: 277-279.

- Subhashini K and Reddy GM (1991). Role of proline in callus growth and plant regeneration under salt stress in rice. Proc. Indian. Nat. Sci.Acad. B 57(1): 81-84.
- Sundaresan S and Sudhakaran RP(1996) Effect of water and deficit stress on proline accumulation in two varieties of Cassava (*Manihot esculenta* Cranz.) differing in their tolerance to draught. Ind. J. Exp. Biol. 159-162.
- Swarup A (1997) Integrated soil nutrient supply and management strategies for sustainability of rice-wheat system on salt affected soils. Ind. Farm. 46(10): 4-11.
- Tal M (1994) *In vitro* selection for salt tolerance in crop plants: theoretical and practical considerations. In vitro cell. Dev. Biol. Plant 30: 175-180.
- Tanji KK (1995) Crop salt tolerance. In Agricultural Salinity Assessment and Management (eds.) Scientific Publishers, Jodhpur, Rajasthan 261-325.
- Thomas JC, Armond RL, Bohnert HJ (1992) Influence of Na Cl on growth, proline and phosphoenol pyruvate carboxylase levels in *Membranthem crystallinum* suspension cultures. Plant. Physiol. 98: 626-631.
- Thorpe TA and Gasper T (1978) Changes in isoperoxidase during shoot formation in tobacco callus cultures. Physiol. Plant 27: 365-369.
- Tiburcio A, Figueras X, Claparols I, Santos M, Torne J (1991) Improved plant regeneration in maize callus culture after pretreatment with DL-alpha-difluromethylarginine. Plant Cell. Tiss. Org. Cult. 27: 27-32.
- Torrigiani P, Altamura MA, Scavamagh S, Captiam F, Talasca G, Bagni N (1993)

 Regulation of Rhizogenesis by Polyamines in Tobacco Thin Layers. J. Plant.

 Phusiol. 142: 81-87.

- Trigiano RN and Conger BV (1987) Regulation of growth and somatic embryogenesis by proline and serine in suspension cultures of *Dactylis glomerata*. J. Plant. Physiol. 130: 49-55.
- Tsukahara M, Hirosawa T, Murayama H (1996) Effect of culture methods oin the regeneration of albino rice (*Oryza sativa* L.) plantlets. Plant. Cell. Rep. 15: 597-600.
- Tyagi RK and Rangaswamy NS (1997) *In vitro* induction and selection for salt tolerance in oil-seed brassicas and regneration of the salt tolerant somaclones. Phytomorphology 47: 209-220.
- Unnikrishnan K, Prakash L, Josekutty PC, Bhatt, PN, Mehta AR (1991) Effect of NaCl Salinity on Somatic Embryo Development in *Sapindus trifoliatus* L. J. Expt. Bot. 42(236): 401-406.
- Van Swaaiji AC, Jacobsen E, Kiel JAKW, Feenstra WA (1986) Selection characterisation and regeneration of hydroxyproline-resistant cell lines of *Solanum tuberosum*: Tolerance to NaCl and freezing stress. Physiol Plant 68: 359-366.
- Vasil IK and Vasil V (1981) Somatic embryogenesis and plant regeneration from suspension cultures of pearl millet (*Pennisetum americanum*) Ann. Bot. 47: 667-668.
- Vasil IK and Vasil V (1994) *In vitro* culture of cereals and grasses. *In* Plant cell and Tissue Culture. Vasil IK and Thorpe TA (eds.). Dodrecht. Netherlands. Kluwer Academic Publishers, 293-312.

- Vasil V and Vasil IK (1982) Characterization of embryogeneic cells suspension culture derived from cultured young inflorescence of *Pennisetum americanum* (Pearl millet). Am. J. Bot. 69: 1441-1449.
- Vasil V and Vasil IK (1986) Plant regeneration from friable embryogenic callus and cell suspension cultures of *Zea mays* L. J. Plant. Physiol. 124: 399-408.
- Vazquez Flota FA, Loyola Vargas VM. (1994) A Catharanthus roseus salt tolerant line

 I. Selection and characterization. J. Plant Physiol. 144: 116-120.
- Vincent AK, Bejoy M, Kavikishor, Hariharan M (1992). Changes in enzyme activities in organ forming and non-organ forming callus cultures of *Kaempferia galanga*L. Phytomorphology 42(3&4): 241-244.
- Voetberg GS and Shrip RE (1991) Growth of the maize primary root at low water potentials. III. Role of increased proline deposition in osmotic adjustment. Plant. Physiol. 96: 1125-1130.
- Wang YC and Creelman RA (1986) In vitro production of jojoba liquid wax by zygotic and somatic embryos. J. Am. Soc. Hort. 111: 798-806.
- Williams EG (1987) Somatic embryogenesis as a Tool in Plant Improvement. In Biotechnology in Agriculture Natesh S, Chopra VL, Ramachandra C (eds.)

 Oxford and IBH Publishing Co. Pvt. Ltd., New Delhi, pp 179-201.
- Williams EG and Maheswaran G (1986). Somatic Emrgyogenesis: Factors influencing coordinated behaviour of cells as an embryogenic group. Ann. Bot. 57: 443-462.
- Wu L and Li HW (1971) Induction of callus tissue initiation from different somatic organs of rice plant by various concentrations of 2,4-dichlorophenoxy acetic acid. Cytologia. 36: 411-416.

- Wyan Jones RG and Gorham J (1983) Aspects of salt and drought tolerance in higher plants. *In* Genetic Engineering of Plants. An Agricultural Prespective T. Kosuge, CP Meredith and A Hollandar (eds.). Plennum Press, New York, pp. 355-369.
- Xiao XG and Branchard M (1993) Emrbyogenesis and plant regeneration of spinach (Spinacia oleracea L.) from hypocotyl segments. Plant Cell. Rep. 13: 69-71.
- Yadav JS and Rajam VM(1997) Spatial distribution of free and conjugated polyamines in leaves of *Solanum melanogena* L. associated with differential morphogenetic capacity. efficient somatic embryogenesis with putrescine. J. Exp. Bot. 48(313): 1537-1545.
- Yancey PH (1994) Compatible and counteracting solutes. In: Cellular and Molecular Physiology of Cell Volume Regulation (ed. K. Strange). CRC Press, Boca Raton, FL, pp 81-109.
- Yang YW., Newton RJ, Miller FR (1990) Salinity tolerance in Sorghum II. Cell culture response to sodium chloride in *S. bicolor* and *S. halepense*. Crop Sci. 30(4): 781-785.
- Yeo A (1998) Molecular Biology of Salt tolerance in the context of whole plant physiology-A Review. J.Exp.Bot. Vol. 49 (323): 915-929.
- Zenk MH (1974) Haploids in physiological and biochemical research. In KJ Kasha, ed. Haploids in higher plants. University of Ghelph, pp 339.
- Zhang BH, Li XL, Li FL (1993). Selection and plant regeneration of NaCl-tolerant variant of cotton callus *in vitro*. Plant. Physiol. Comm. 29: 423-425.
- * Original papers not seen