List of Figures

Figures		Page
1.1	Schematic representation of primary production, new production and regenerated production in the photic zone	4
1.2	Monsoon wind-stress filed over the northern Indian Ocean for July (summer monsoon) and January (winter monsoon)	10
2.1	Study Area for Ocean color studies	22
2.2	Block diagram showing steps involved in estimation of new and regenerated production	23
2.3	The cruise track along which sampling was done during SS-220. PP denotes primary productivity stations	24
2.4	The cruise track along which sampling was done during SK-214. PP denotes primary productivity stations	25
2.5	The cruise track along which the sampling was done during SK-220	27
2.6	The cruise track along which sampling was done for the present study in the Southern Ocean (ABP15)	29
2.7	Go-Flo bottles attached to a CTD rosette used to collect sea-water samples from different depths	30
2.8	Author filtering samples in dark on board R/V Akademik Boris Petrov in the Arabian Sea	34
2.9	Some typical examples of (a.) calibration curve obtained immediately after putting fresh chemicals in the oxidation/reduction chambers and (b.) examples of calibration curve used for the estimation of particulate organic nitrogen in the post incubation sample	39
2.10	Schematic representation of the sampling adopted in the present study for iron enrichment experiment	44
3.1	Monsoon wind stress field from the NCEP climatology for a) January, b) April, c) July and d) November	49
3.2	Satellite derived Chl-a over zone 1	53

Figures		Page
3.3	Satellite derived SST over zone 1	53
3.4	Satellite derived Chl-a over zone 2	54
3.5	Satellite derived SST over zone 2	55
3.6	Satellite derived Chl- <i>a</i> data and SSTs from May to September from the region 52°E to 57°E and 5°S to 10°N	55
3.7	Satellite derived wind speed over zone 1	56
3.8	Satellite derived wind speed over zone 2	57
3.9	Satellite derived Chl-a data and SSTs over Zone 1 in winter (October to March)	57
3.10	Satellite derived Chl-a data and SSTs over Zone 2 in winter (October to March)	58
3.11	Vertical profiles of Chl- <i>a</i> at different stations in the eastern Arabian Sea during late winter monsoon (FebMarch 2004)	60
3.12	Euphotic zone integrated chl-a concentration at all the stations during FebMarch 2004	61
3.13	Vertical profiles of nutrients at different stations during the present study (cruise SS#222)	62
3.14	SST and salinity at different stations in the eastern Arabian Sea during Feb-March 2004	64
3.15	Temperature-depth profiles at different stations obtained using CTD data during FebMarch 2004	65
3.16	Station-wise column-integrated total N-uptake for eastern Arabian Sea during the present study	66
3.17	New production at different stations in the eastern Arabian Sea during the late winter monsoon	68
3.18	Relationship between total N uptake rate and nitrate uptake rate in the eastern Arabian Sea during the late winter monsoon	69
3.19	Ammonium and urea uptake rates at different stations in the eastern Arabian Sea during the late winter monsoon	70
3.20	<i>f</i> -ratios at different stations in the Eastern Arabian Sea during the late winter monsoon.	71

Figures

Page

3.21	Station-wise column integrated chlorophyll concentration during Dec-2004.	73
3.22	Vertical profiles of Chl- <i>a</i> at different stations in the eastern Arabian Sea during the early winter monsoon (Dec - 2004)	74
3.23	Vertical profiles of nitrate at different stations during December-2004	75
3.24	SST and salinity at different stations in the eastern Arabian Sea during early winter monsoon (Dec-2004)	76
3.25	Temperature-depth profiles at different stations obtained using CTD data during December 2004	77
3.26	Total N-uptake rate (new production) at different stations in the eastern Arabian Sea during December 2004	78
3.27	Nitrate uptake rate (new production) at different stations in the eastern Arabian Sea during December 2004	80
3.28	Relationship between total N-uptake during the early winter monsoon (December) in the eastern Arabian Sea	81
3.29	Ammonium and urea uptake rates at different stations in the Eastern Arabian Sea during early winter monsoon	82
3.30	<i>f</i> -ratio at different stations in the eastern Arabian Sea during the early winter monsoon	83
4.1	Figure showing wind stress field for the Indian Ocean during January, April, July and October	90
4.2	SST (°C) at all stations along 77°E and 83°E transects	92
4.3	Salinity at all stations along 77°E and 83°E transects	93
4.4	Photic zone integrated nitrate at different stations in the equatorial Indian Ocean	94
4.5	Depth profiles of nitrate at different stations in the equatorial Indian Ocean during pre-monsoon 2005	95
4.6	Total N-uptake rates at different stations in the equatorial Indian Ocean during pre-monsoon 2005	96
4.7	Mixed layer depth integrated total N-uptake rates at different stations in the equatorial Indian Ocean during pre-monsoon 2005	97

Figures

.

,

Page

,

4.8	Photic zone integrated total nitrate uptake rates (new production) at different stations in the equatorial Indian Ocean during pre-monsoon 2005	98
4.9	Station-wise mixed layer depth integrated nitrate uptake rates (new production) at different stations in the equatorial Indian Ocean during pre-monsoon 2005	99
4.10	Relation between mixed layer integrated total N-uptake rate and nitrate uptake rate in the equatorial Indian Ocean during pre- monsoon 2005	100
4.11	Mixed layer depth integrated ammonium uptake rates at different stations in the equatorial Indian Ocean during pre-monsoon 2005	102
4.12	Mixed layer depth integrated urea uptake rates at different stations in the equatorial Indian Ocean during pre-monsoon 2005	103
4.13	<i>f</i> -ratios at different stations in the equatorial Indian Ocean	104
4.14	The aerial extent of the Southern Ocean (left) and the cartoon of the Antarctic Circumpolar Current (ACC; right)	105
4.15	Vertical profiles of Chl-a (left panel) and column integrated Chl-a (right panel) at different stations in the Southern Indian Ocean	110
4.16	Vertical profiles of ambient nitrate concentrations at different stations in the Southern Indian Ocean	112
4.17	SST (in °C) at different stations in the Southern Indian Ocean	113
4.18	Temperature-depth profiles at different stations in the Southern Indian Ocean. The stations in the south had deeper mixed layers compared to those in the north	114
4.19	Total N-uptake rates at different stations in the Southern Indian Ocean	115
4.20	New production at different stations in the Southern Indian Ocean	118
4.21	Relationship between total N uptake and nitrate uptake in the Southern Indian Ocean	119
4.22	Ammonium and urea uptake rates at different stations in the Southern Indian Ocean	120
4.23	f-ratios at different stations in the Southern Indian Ocean	122