

TABLES

Table No	TABLE DESCRIPTION	Page No.
1	Brief mode of action of rot fungi	13
2	Commercial xylanase preparations available for enzymatic bleaching of pulps	28
3	Different media used for isolation of fungi	32
4	Composition of Czapek dox media	34
5	Composition of Malt extract media	34
6	Composition of different moistening agents	36
7	Composition of DNSA	38
8	Ammonium sulphate precipitation	39
9	Composition of 12% resolving gel	41
10	Composition of stacking gel used	41
11	Composition of treatment buffer	42
12	Composition of glycine SDS buffer	42
13	Composition of CBB staining solution	42
14T	Effect of different substrate on xylanase production	45
15T	Activity of xylanase -substrate reaction at different incubation time	46
16T	Effect of temperature on xylanase activity	47
17T	Effect of moistening agents on xylanase activity	48
18T	Effect of moisture level on xylanase production	49
19T	Effect of carbon source on production of xylanase	50
20T	Effect of metal ions on xylanase activity	52
21C	Effect of different combinations on xylanase production	56
22C	Effect of incubation time (days) on xylanase production	57
23C	Effect of moistening agents on production of xylanase	60
24C	Effect of moisture level on xylanase production	61
25C	Effect of mutagenesis on production of xylanase	65
26C	Purification profile of xylanase	65
27	Fungus grown in EP medium for amylase production	78
28	Ammonium sulphate precipitation table	80
29	Composition of 10% resolving gel	82

30	Composition of stacking gel	82
31	Effect of incubation time on amylase activity	93
32	Effect of metal ions on amylase activity	94
33	Various media used for <i>C. asperatum</i> growth and metabolite production	113
34	Fungi that cause wood deterioration and their characteristics	121
35	Average percent weight loss during each incubation period (60, 90 and 120 days)	126
36	Average percent weight loss during each incubation period (60, 90 and 120 days)	132