BIBLIOGRAPHY

- Adam G and Marquardt V, 1986. Brassinosteroids. Phytochemistry 15: 1987.
- Agnew WS, 1985. Squalene synthetase, In. Methods in enzymology, Vol. 110: 359.
- Akoyunoglou GA and Siegelman HW, 1968. Protochlorophyllide synthesis in dark grown bean seedlings. Plant Physiol. 43: 66.
- Alberts AW, Chen J, Kuron G, Hunt V, Huff J, Hoffman C, Rothrock J, Lopez M, Joshua H, Harris E, Patchet A, Monagan R, Currie S, Stapley E, Hensens O, Liesch J and Springer J, 1980. Mevinolin: a highly polent competative inhibitor of 3-hydroxy 3-methyl glutaryl coenzyme-A reductase and cholesterol lowering agent. Proc. Natl. Acad. Sci. USA, 77: 3957.
- Alexopoulose CJ and Beneke ES,1955. Laboratory mannual for introductory mycology. Bourgess Pub. Co., Minneapolis. pp 3.
- Almarego WLF, Goad LJ and Goodwin TW, 1973. Biosynthesis of α -spinasterol from 2^{14} C (4 R) 4 3 H₁) mevalonic acid by Spinacea oleracea and Medicago sativa. Phytochemistry, 12(2): 181.
- Anastasis P, Freer I, Overton K, Rycroft D and Sing SB,1985.

 Journal of Chemical Society and Chemical Communication,

 148: 149.
- Anonymous, 1988. Simple prevention of water loss from micropropagated plants. Agricell reports, 11(3): 1.

- Arebalo RE and Mitchell Jr. ED, 1984. Cellular distribution of 3-hydroxy 3-methylglutaryl CoA reductase and mevalonate kinase in leaves of Nepeta cataria. Phytochemistry, 23: 13.
- Atal CK and Kapoor M, 1972. Medicinal and aromatic plants, National Press, Jammu-Tawi, India.
- Bach TJ, 1981. Untersuchungen zur charakteriseierung und regulation der 3-hydroxy 3-methylglutaryl CoA acylierend, E.C.1.1.34 in Kemlingen von Raphanus sativus Karstr.Beitr Pflanzenphysiol,10:1.
- Bach TJ, 1985. Selected natural and synthetic enzyme inhibitors of sterol biosynthesis as molecular probes for in vivo studies concerning the regulation of plant growth. Plant Sci.,39: 183.
- Bach TJ, 1986. Hydroxymethylglutaryl CoA reductase, a key enzyme in phytosterol syntehsis. Lipids, 21: 82.
- Bach TJ, 1987. Synthesis and metabolism of mevalonic acid in plants. Plant Physiol.,25(2):163.
- Bach TJ and Lichtenthaler HK, 1982. Mevinolin: a highly specific inhibitor of microsomal 3-hydroxy 3-methylglutaryl coenzyme A reductase of radish plants. Z. Naturforsch, 37.C: 46.
- Bach TJ and Lichtenthaler HK, 1983. Inhibition by mevinolin of plant growth, sterol formation and pigment accumulation. Physiol Plant, 59:50.

- Bach TJ and Lichtenthaler HK, 1983. Mechanism of inhibition by mevinolin (MK-803) of microsome bound <u>Radish</u> and partially purified yeast HMG CoA reductase (E.C.1.1.1.34) Z. Naturforsch, 38.C: 212.
- Bach TJ and Lichtenthaler HK, 1984. Application of modified Line weaver-burk plots of studies of kinetics and regulation of Radish HMG CoA reductase. Biochemica. Biophysica.Acta, 794: 152.
- Bach TJ and Lichtenthaler HK, 1987. Plant growth regulation by mevinolin and other sterol biosynthesis inhibitors In: Ecology and metabolism of plant lipids. (Fuller G and Nes WD eds) A C S symposium series, 235, 199. Washington DC.
- Bach TJ, Lichtenthaler HK and Retty J, 1980. Properties of membrane bound HMG CoA reductase (E.C.1.1.1.34) from Radish seedlings and some aspects of its regulation. In: Biogenesis and functions of plant lipids (Mazliak P, Beneveniste P, Costes C and Douce R eds), Elsvier, Holland and Biomedical Press.
- Bach TJ and Webber T, 1989. Enztmatic synthesis of mevalonate in plants. In: Biological role of plant lipids. (Biacs PA Cruiz K and Kremmer T eds) P.279. Plenum Press, N.Y.
- Bae M and Mercer EI, 1970. The effect of long and short day photoperiods on sterol levels in the leaves of Solanum andigena. Phytochemistry, 9: 63.

- Baker EA, 1974. The influence of environment on leaf wax development in Brassica oleracea. New Phytol, 73: 955.
- Beneveniste P, 1986. Sterol biosynthesis. Ann. Rev. of Plant physiology, 37: 275.
- Berg D 1986. In: Fungicide chemistry: Advances and Practical applications. (Green MB and Spilker DA eds) P.125. ACS Symposium series, 325, Washington D.C.
- Bergman L, 1960. Growth and division of single cells of higher plants in vitro J.Gen. Physiol.,43:841.
- Berlin J, 1980. Paraflurophenylalanine resistant cell lines of tobacco. Z. Pflanzenphysiol. 97: 317.
- Berlin J, 1986. Secondary products from plant cell cultures. In:Biotechnology (Rhem HJ and Reed G eds) 4:630.VCH Publishers.
- Berlin J, Beiber H, Fecker L, Forche E, Noe N, Sasse F and Schiel O, 1985. Conventional and new approaches to increase alkaloid production of plant cell cultures. In: Primary and secondary metabolism of plant cell cultures (Neumann KH, Barz W and Reinhard E eds) P.273 Springer verlag, Berlin and N.Y.
- Berlin J and Sasse F, 1985. Selection and screening techniques for plant cell cultures. In: Advances in Biochemical eng./Biotechnology 31: 99. Springer verlag, Berlin.
- Berlin J, Sieg S, Strack D, Bokmen M and Harms H, 1986.

 Production of betanins by suspension culture of

- <u>Chenopodium rubrum</u> L. Plant Cell tissue and organ culture, 5: 163.
- Berlin J and Wollmer B, 1973. Effect of phenylpropionic acid and phenylalanine metabolism in p-fluorophenylalanine sensitive and resistant tobacco cells Z. Naturforsch, 34 C: 770.
- Berlin J and Widholm JM, 1977. Correlation between phenyl alaminine ammonia lyase activity and phenolic biosynthesis in p-fluorophenylalanine sensitive and resistant tobacco and carrot tissue cultures. Plant physiol. 59: 550.
- Berlin J and Widholm JM, 1978. Metabolism of phenyl alanine and tyrosine in tobacco cell lines resistant and sensitive to p-fluorophenyl alanine. Phytochemistry, 17: 65.
- Berlin J, Witte L, Hammer J, Kuboschke K, Zimmer A and Pape D, 1982. Metabolism of p-fluorophenyl alanine sensitive and resistant tobacco cell cultures. Planta, 155: 244.
- Beytia E Valenzuela P and Cori O, 1969. Arch. Biochem. Biophys 129: 346.
- Bhatt PN and Bhatt DP and Sussex IM, 1979. Organ regeneration from leaf discs of S. dulcamara and S. Khasianum. Z. Pflanzenphysiol, 95: 355.
- Bhatt PN and Bhatt DP, 1983. Studies on some factors affecting solasodine contents in tissue culture of Solanum nigrum. Physiol. Plant.,57:159.

- Bhatt PN and Bhatt DP, 1984 a. Changes in sterol composition during leaf aging and in vitro differentiation in solanum nigrum. J. Natl. Products. 47(3): 426.
- Bhatt PN and Bhatt DP, 1984 b. Regulation of sterol biosynthesis in Solanum species J. Exp. Bot. 35(155): 890.
- Bhatt DP, Bhatt PN and Mehta AR, 1986. Steroid analaysis and plant regeneration from cell suspension cultures of S.dulcamara L. Bietr. Biol. Pflanzen, 61: 203.
- Bhatt DP and Fassuliotis, G, 1981. Plant regeneration from protoplasts of egg plants. Z. Pflanzenphysiol.104:81.
- Birner J, 1969. Determination of total steroid bases in Solanum Spp. J. Pharm. Sci. 58(2): 258.
- Bogorad L, 1969. The biosynthesis of chlorophylls. In: The The chlorophylls (Vernon LP and Seely GR eds) Academic Press, New York.
- Bourgin JP, 1978. Valine resistant plants from <u>in vitro</u> selected tobacco cells. Mole. Gen. Genet.161: 225.
- Bradford MM, 1976. A rapid and sensitive method of quantification of microgram quantities of protein utilizing the principle of prolein dye binding .Anal. Biochem.,72: 248.
- Brain KR and Williams MH, 1983. Evidence for an alternate route from sterol to sapogenin in suspension cultures of Trigonella foenemgracium Plant cell Reports 2:7.
- Braithwaite GD and Goodwin TW, 1980. The incorporation of

- ^{14}C acetate, ^{14}C mevalonate and ^{14}C $^{0}\text{2}$ in to -carotene by the fungus phycomyces blakesleeanus. Biochem J. 76:1
- Brodelius PE, Funk C and Shillito RD, 1988. Plant Cell Reports. 7: 186.
- Brodie JD, Wasson G and Porter JW, 1963. Journal of Biological Chemistry, 238: 723.
- Brooker JD and Russel DW, 1975a. Properties of microsomal 3-hydroxy 3-methylglutaryl-coenzyme A reductase from Pisum sativum seedlings.Arch. Biochem. Biophys, 167:723.
- Brooker JD and Russel DW, 1975b. Subcellular localization of 3-hydroxy 3-methylglutaryl-coenzyme A reductase in Pisum sativum seedlings.Arch. Biochem. Biophys. 167: 730.
- Brooker JD and Russel DA, 1979. Regulation of microsomal 3-hydroxy 3-methyl glutaryl CoA reductase from pea seedlings Rapid post translational phytochrome mediated decrease in activity and in vivo regulation by isoprenoid products.Arch.Biochem Biophys 198:323.
- Brown MS, Faust JR and Goldstein JL, 1978. Induction of HMG CoA activity in human fibroblast incubated with compactin (ML.236.B), a competative inhibitor of reductase. The J.Biol. Chem. 253(4): 1121.
- Brown MS and Goldstein LJ, 1980. Multivalent feedback regulation of HMG CoA reductase, a control mechanism coordinating isoprenoid synthesis and cell growth. J. Lipid Res. 21: 505.

- Brown WE and Rodwell VW, 1980. Hydroxy methylglutaryl CoA reductase In: Dehydrogenases requiring Nicotinamide coenzymes (Jeffery J ed) p.232. Birkhauser verlag, Berlin.
- Burden RS, Clark T and Holloway PJ, 1987. Effects of sterol biosynthesis inhibiting fungicides and plant growth regulators on the sterol composition of barley plants. Pestic. Biochem. Physiol. 27: 289.
- Burden RS, David T Cooke and Carte GA, 1989. Inhibitors of sterol biosynthesis and growth in plants and fungii. Phyto chem., 28(7): 1791.
- Bush PB, Grunwald C and Paris DW, 1971. Changes in sterol composition during greening of etiolated barley seedlings Plant. Physiol, 47: 745.
- Bush PB and Grunwald C, 1973. Effect of light on mevalonic acid incorporation in to the phytosterols of Nicotiana tabacum L. Seedlings. Plant Physiol. 51: 110.
- Camara B, Bardat F, Dogbo O, Brangeon J and Moreger R, 1983.

 Isoprenoid metabolism in plastids. Isolation and biochemical characteristics of Capscium annum chromoplasts. Plant Physiol.,73: 94.
- Campbell CE and Kircher HW, 1980. Phyto Chemistry, 19: 2777.
- Ceccarelli N and Lorenzi R, 1984. Growth inhibition by competative inhibitors of 3-hydroxy 3-methylglutaryl coenzyme A reductase in <u>Helianthus tuberosus</u> tissue explants. Plant Sci. Lett, 34: 269.

- Chaleffe RS, 1981. Genetics of higher Plants: Applications of cell culture. Cambridge Univ. Press, N.Y.
- Chaleffe RS, 1983. Isolation of agronomically useful mutants from plat cell cultures. Science, 219: 676.
- Chandler SF and Dodds JH, 1983a. Solasodine production in rapidly proliferating tissue cultures of Solanum laciniatum Ait Plant Cell Reports, 2: 69.
- Chandler SF and Dodds JH, 1983b. Adventitious shoot initiation in serially subcultured callus cultures of Solanum laciniatum. Z. Pflanzenphysiol, 111: 115.
- Chang TY and Limaneck JS, 1980. J. Biological Chemistry, 255: 7787.
- Chappel J, Lanken CV and Vogeli U and Prashant Bhatt, 1989.

 Sterol and sesquiterpenoid biosynthesis during a growth cycle of tobacco cell suspension cultures. Plant Cell Reports, 8: 48.
- Chappel J and Ross Nabel, 1987. Induction of sesquiterpenoid biosynthesis in tobacco cell suspension cultures by fungal elicitor. Plant Physiol, 85: 469.
- Chen CM and Melitz DK, 1979. FEBS Lett.107:15.
- Chin DJ, Luskey KL, Goldstein LJ and Brown MS, 1982.

 Appearance of crystalloid endoplasmic reticulum in compactin-resistant chinese hamster ovary cells with a 500 fold elevation in 3-hydroxy 3-methyl glutaryl coenzyme A reductase. Proc. Natl.Acad. Sci. USA, 79: 1185.

- Chin DJ, Gill G, Russel DW, Liscum L, Basu KL, Goldstein JL and Brown MS, 1984. Nucleotide sequence of HMG CoA reductase, a glycoprotein of the endoplasmic reticulum. Nature, 308: 613.
- Chiu PL, Bottino PJ and Patterson GW, 1980. Sterol composition of nystatin and amphotericin B resistant tobacco calluses. Lipids, 15:50.
- Conner AJ and Meridith CP, 1989. Genetic manipulation of plant cells. In: The biochemistry of plants. 15: 653. (Stumpf PK and Conn EE eds) Academic Press, London.
- Davies BH, 1977. In: Lipids and Lipid polymers in higher plants. P. 199 (Terini M and Lichtenthaler HK eds)
 Springer Verlag, Berlin.
- Deus-Neumann B and Zenk MH, 1984b. Instability of Indole alkaloid production in <u>Catharanthus</u> roseus cell suspension cultures Planta. Med.,50: 427.
- Dix PJ, Mehysagnt UA and Pearce RS, 1984. The potential of cell cultures for production of salt tolerant cultivars. In: Efficiency in plant breeding. Proc. 10th Cong. Eucarpica. (Lange W, Zenen AC, and Hogenboom NG eds) P 219. Wageningen.
- Doll M, Schindler S, Lichtenthaler HK and Bach TJ,1984.

 Differential inhibition by mevinolin of prenyl lipid accumulation in cell suspension cultures of Sylibum marianum L. In: Structure, functions and metabolism of plant lipids (Seigenthaler PA and Eichenberger eds)

 P.227, Elsevier, Amsterdam.

- Douglas TJ and Paleg GL, 1979. Plant growth retardants as inhibitors of sterol biosynthesis in tobacco seedlings. Plant Physiol.,54: 228.
- Douglas TJ and Paleg LG. 1978. AMO 1618 and sterol biosynthesis in tissues and subcellular fractions of tobacco seedlings. Phytochem. 17: 705.
- Douglas TJ and Paleg, LG, 1981. Inhibition of sterol biosynthesis and stem elongation of tobacco seedlings induced by some hypocholesterolemic agents. J Exptl. Botany, 32 (126):59.
- Driver JA, 1986. Method for acclamatizing and propagating plant tissue culture shoots. United States Patent No.4, 612, 725 : 1.
- Dugan RE, 1981 in Biosynthesis of isoprenoid compunds Vol.1, (Porter JW and Spurgeon SL eds) P.95. Wiley, N.Y.
- Eilert U and Constable F, 1986. Elicitation of Sanguimarine accumulation in <u>Papaver somniferum</u> cells by fungal homogenates An induction process. J. Plant. Physiol, 125: 167.
- Emke and Eilert U, 1986. Steroidal alkaloids in tissue culture and regenerated plants of <u>Solanum dulcamara</u>. Plant Cell Reports 5: 34.
- Erdie L, Stainer BCCE and Kuiper PJC, 1980. The effect of salinity on lipid composition and on activity of ${\rm Ca}^{2+}$ and ${\rm Mg}^{2+}$ stimulated ATPase in salt sensitive and salt tolerant Plantago Sp. Physiol. Plant., 49: 315.

- Evans JL and Gealt MA, 1985. Journal of General microbiology 131: 279.
- Faust JR, Goldstein JL and Brain MS, 1979. Squalene synthetase in human fibroblasts, regulation via the low density lipoprolien receptor. Proc. Natl. Acad.Sci. USA. 76: 5018.
- Fletcher RA, Hofstra G and Gao I, 1986. Comparative fungitoxic and plant growth regulating propertis of triazole derivatives. Plant Cell Physiol., 27(2): 367.
- Flick CE, Evans D and Sharp WR, 1983. Organogenesis. In:
 Handbook of Plant cell culture Vol.1 (Evans, Sharp,
 Ammirato and Yamada eds) P.13. Macmillan Pub. Co., N.Y.
- Fujita Y and Tabata M, 1987. Secondary metabolites from plant cells. Pharmaceutical applications and progress in commercial production In: Plant Biology (Green CE, Somers DA, Hackett WP and Biesboer DD eds) Vol.3, P.169. Alan R.Liss Inc., N.Y.
- Gassmann M and Bogorad L, 1967. Chlorophyll production in rapidly greening bean leaves. Plant Physiol.42:774.
- Ghannoum MA, Taini, G, Khamis L, and Radman SS, 1986. J.Gen. Microbiol.132: 2367.
- Golbeck JH, Maurey KM and Newberry GA, 1985. Detection of 3-hydroxy 3-methyl glutaryl coenzyme A reductase (HMGR) activity in <u>Dunaliella</u> salina plant physiol. 77: (supplement) 48

- Goldstein JL and Brown MS, 1984. Progress in understanding the LDL receptor and HMG-CoA reductase, two membrane proteins that regulate the plasma cholesterol. J.Lipid.Res.25: 1450.
- Goldstein JL and Brown MS, 1990. Regulation of mevalonate pathway. Nature, 343: 425.
- Gonzales RA, Das PK and Widholm JM, 1984. Characterization of cultured tobacco cell lines resistant to methionine analogue. Plant Physiol. 74: 640.
- Gonzales RA and Widholm JM, 1985. Selection of plant cells for desirable characteristics: Inhibitor resistance.In:

 Plant Cell culture a practical approach. (Dixon RA ed.)

 P.67, IRL Press Oxford, Washington, D.C.
- Goodwin TW, 1960. Chemistry, biogenesis and physiology of carotenoids In: (Rhuland W, ed) Encyclopedia of plant physiology. 5(1): 394. Springer: Berlin.
- Goodwin TW, 1980. In: The biochemistry of plants. Stumpf PK and Conn EE eds) 4:485. Academic Press, New York.
- Greabe J, 1967. Phytochemistry 7: 2003.
- Greabe JE, Dennis DT, Upper CD and West CA, 1965.

 Biosynthesis of gibberellins 1. The biosynthesis of

 (-)-Kaurene, (-)-Kaur-19-01 and trans geranylgeraniol in

 endosperm nucellus of <u>Echinocystis macrocarpa</u> Greene. J.

 Biol Chem. 240: 1847.

- Graebe JE and Ropers H, 1978. Gibberellins In: Phytohormones and related compounds A comprehensive treatise Vol. 1 P.107,(Letham DS, Goodwin PB and Higgins TJV eds)Elsevier North Holland biomedical Press, Amsterdam.
- Gray JC, 1987. Control of isoprenoid biosynthesis in plants.

 In: Advances in Botanical Research (Cullow JA ed.)

 Vol.14:25. Academic Press, London.
- Green TR and Baisted DJ, 1971. Biochemical Journal, 125:1145.
- Green TR and Baisted DJ, 1972. Biochemical Journal, 130:983.
- Grogger D, 1988. Isoprenoid and steroid alkaloids In: Cell culture and somatic cell genetics of plants (Vasil IK ed) Vol.5: 435.
- Grossmann K, Weiler EW and Jung J, 1985. Effects of different sterols on the inhibition of cell culture growth caused by the growth retardant tetcyclasis. Planta, 164: 370.
- Grumbach and Bach,1979. The effect of PS II herbicides, Amitrol and SAN 6706 on the activity of 3-hydroxy 3-methylglutaryl CoA reductase and the incorporation of 2^{14} C acetate and 2^{3} H mevalonate in to chloroplast pigments of Radish seedlings Z. Naturforsch. 34C: 941.
- Grunwald C. 1968. Effect of Sterols on the permeability of alcohol treated red beet tissue. Plant Physiol.,43: 484.
- Grunwald C, 1970. Quantitative analaysis of free phytosterols by gas chromatography using stationary phase OV 101. Anal. Biochem; 34: 16.

- Grunwald C, 1971a. Effects of free sterols, steryl esters and steryl glycosides in membrane permeability. Plant Physiol.,48: 653.
- Grunwald C, 1974. Sterol molecular modification influencing membrane permeability. Plant Physiol.,54: 624.
- Grunwald C, 19⁷5. Plant sterols Ann. Rev. of plant physiol. Suppl.,86: 85.
- Grunwald C, 1978. Functions of sterols. Phil.Trans R.Soc. Lond.B., 284: 544.
- Grunwald C, 1986. Steroids in Secondary Plant products. In:
 Encyclopedia of Plant Physiology, New Series Vol.8,
 P.221. (Bell EA and Charlwood BV eds). Springer-Verlag,
 Berlin.
- Hartman CL, Mc Roy TJ, Knows TR, 1984b. Selection of alfalfa (Medicago sativa) cell lines and regeneration of plants resistant to the toxin(s) produced by Fusarium oxysporum f.Sp medicaginis. Plant Sci. Lett. 34: 183.
- Hartmann G and Lynnen F, 1961. In: "The enzymes" (Boyer PD, Lardyy H and Myrbach K eds) 2nd ed., Vol 5, p.381. Academic press, N.Y.
- Hashizune T, Matsubara S and Endo A, 1983. Agricultural biological chemistry, 47: 1401.
- Haughan PH, Lenton JR and Good LJ, 1987. Inhibition of Plant sterol biosynthesis by Palcobutrazol and its effect on cell growth. In: Proc. XIV Intl. Bot. Cong. (Zimmer WUB, and Bohake HD eds) Absts. No.1 P.106. berlin W.Germany.

- Haughan PA, Lenton JR and Goad LJ, 1989. Inhibition of Celery cell growth and sterol biosynthesis by the enantiomaers of Palcobutrazol. Phytochemistry, 28(3):781.
- Heble MR, Narayanaswamy S and Chadha MA, 1968. Solasonine in tissue cultures of Solanum xanthocarpum. Natur. Wissenchaften, 55: 350.
- Hedden P, Mac Millan and Phinney BO, 1978. The metabolism of Gibberellins. Ann.Rev.Plant.Physiol., 29:149.
- Heftmann E, 1971. Functions of steroids in Plants. Lipids,6: 128.
- Heftmann E, 1975. Functions of Steroids in Plants. Phytochemistry, 14: 891.
- Heftman E, 1984. Biogenesis of steroids in Solanaceae. Phytochemistry, 22 (9):1843.
- Helenius A and Simons K, 1975. Biochem. Biophys. Acta.,415:
- Hepper CM and Audley BG, 1969. The biosynthesis of Rubber from 3-hydroxy 3-methyl glutaryl CoA in <u>Hevea braseliensis</u> latex, Biochem. J. 114: 379.
- Heupel RC, Sauvaire Y, Le PH, Parish EJ and Nes WD, 1986.

 Sterol composition and biosynthesis in Sorgham.

 Importance to developmental regulation. Lipids, 21:69.
- Hibberd CA, Walter T, Green CE and Gengenbach BG, 1980. Selection and characterization of Feedback insensitive tissue culture of maize. Planta, 148: 183.
- Higgins MJP and Kekwick RGO, 1973. Biochem.J., 134: 295.

- Holden M, 1965. Chlorophylls. In: Chemistry and Biology of plant pigments (Goodwin TW, ed) P.462.
- Hommes NG, Akiyoshi DE and Morris RO, 1985. Meth Enzymol. 110:310.
- Hosoda N, Ito H and Yatasawa M, 1979. Some accounts on culture conditions of callus tissues of Solanum laciniatum Ait. Agri. Biol.Chem., 43(8) 1745.
- Hosokawa G, Patterson G and Lusby WR, 1984. Effects of triarimol, tridemorph and triparanol on sterol biosynthesis in carnot, tobacco and soybean suspension cultures. Lipids, 19: 449.
- Ingram TJ, Reid JB, Murfet IC, Gaskin P, Willis CL and Mc Mc Millan J, 1984. Internode length in pisum: The Le gene controls the 3 -hydroxylation of gibberellin A20 to gibberellin A1. Planta, 160:455.
- Isa RBM and Sipat PR, 1982. Biochim. Biophys.Res.Commun., 108: 206.
- Ito R, Oba K and Uritani I, 1979. Mechanisms for the induction of 3-hydroxy 3-methylglutaryl-coenzyme A reductase in HgCl₂ treated sweet potato root tissue. Plant Cell Physiol., 20: 867.
- Jacob John S, Subramani J, Bhatt PN and Mehta AR, 1989. A novel approach for the production of guggullipid using immobilized differentiated cells of Commiphora wightii. Elsevier Publication, Amsterdam, p.338.

- Jacobs M, Cattoir-Reynaerts A, Negrutin I, Verbuggen I and Degryse E, 1982. Comparison of selection schemes for the isolation of resistant mutants to asparlate derived amino acids and 5(2-aminoethyl) cystein in various cell cultures and whole plant systems. In: Plant tissue culture (Fujiwara A ed) p.453. Maruzen Co.Ltd, Tokyo.
- Jain SC and Sahoo SL, 1981. Growth and Production of steroidal sapogenins and glycoalkaloids in Solanum jasminoides plant suspension cultures. Agric. Biol.Chem., 45, 2909.
- Johnston JA, Racusen DW and Bonner J, 1954.Proc.Natl.Acad. Sci. USA, 40: 1031.
- Joseph Chappel, Lanken CV, Vogeli U, and Bhatt P 1989.

 Sterol and segguiterpenoid biosynthesis during a growth cycle of tobacco cell suspension cultures. Plant Cell Reports, 8: 48.
- Kawasaki, S, Ramagopal M, Chin J and Bloch K, 1985. Proc. Natl. Acad. Sci. USA, 82, 5715.
- Kita T, Brown MS and Goldstein JL, 1984. Feedback regulation of HMGR in livers of mice treated with mevinolin, a competative inhibitor of the reductase. J. Clin. Invest. 66:1094.
- Kondo K and Obo K 1986. Purification and characterization of HMG CoAR from potato tubers. J. Biol. Chem., 100: 967
- Koorneef M, Rueling G and Karssen CM, 1984. The isolation and characterization of abscisic acid insensitive mutants of Arabidopsis thaliana. Physiol Plant., 61: 377.

- Kreis W and Reinhard E, 1989. Production of secondary metabolites by plant cells cultured in bioreactors. Planta Med., 55: 409.
- Kreuz K and Kleinig H, 1982. Polyprenoid biosynthesis in chloroplasts and chromoplasts. In: Biochemistry and metabolism of plant lipids. (Wintermens JFGM and Kuiper PJC eds) P.507. El-sevier-Biomedical Press, Amsterdam.
- Kreuz K and Kleinig, 1984. Europen Journal of Biological Chemistry. 141: 531.
- Krikorian AD <u>et al</u> 1987. In: Plant hormones and their role in plant growth and development (Davies PJ ed) Martinus Nijhoff, Boston.
- Kuc J and Rush JS, 1985. Phytoalexins.Arch Biochem.Biophys., 236: 455.
- Kuiper PJC, 1984. In: Structure, function and metabolism of plant lipids. (Siegenthaler PA and Eichenberger W eds) P.525. Elsevier, Amsterdam.
- Lancaster EJ, &Mann JD, 1975. Changes in Solasodine content during the development of Solanum laciniatum Ait. NJ2 Agric. Res., 18: 139.
- Langeake, P, Khun PJ and Wade M, 1984. In "Progress in pesticide biochemistry and toxicology" Vol.3 (Huston DH and Roberts TR, eds) P: 1, Wiley, Chichester.
- Lea PJ & Norris RD, 1976. Phytochemistry, 15: 585.
- Lenton JR, 1987. Br. Plant Growth Reg. Group, 9:1
- Lichtenthaler HK, 1979. In "Advances in the Biochemistry and physiology of plant Lipids" (Applequist, L A, Liljenberg C, Eds) Elsevier: Amsterdam.

- Lichtenthaler HK, 1980. In "Biogenesis and Function of Plant Lipids" (Mazliak P, Benveniste P, Coster C, Douce R, Eds) Elsevier Amsterdam pp: 299.
- Lichtenthaler HK, Bach TJ Wellbuen AR, 1982. Cytoplasmic and plastidic isoprenoid compounds of oat seedlings and distinct labelling from ¹⁴C mevalonate In "Biochemistry of metabolism of Plant Lipids", (Wintermans JFGM and Kuiper PJC eds) Elsevier Biomedical Press, London.
- Limonrier CH, Rossignol L, Sihachakr M and Haicur R, 1989.

 Solasodine and pigment contents of plants derived from callus culture of Solanum laciniatum Ait.

 Phytomorphology, 39(1): 93.
- Loffler RST, Butters JA and Hollomon DW, 1984. Proc. Br. Crop. Protect. Conf. Pests. Dis., 911.
- Low C, Rodriguez RJ and Parks LW, 1985. Arch. Biochem. Biophys., 240: 530.
- Lursgen K, 1987. In "Hormone action in plant development" (Hoad GV, Lenton JR, Jackson MB Atkin RK eds) P:133 Butterworths, London.
- Luskey KL, Faust JR, Chin DJ, Brown MS and Goldstein JL, 1983. Amplification of the gene for 3-hydroxy 3-methylglutaryl coenzyme A reductase, but not for the 53-K Da protein, in VT-1 Cells. J.Biol Chem., 258: 8462.
- Lynen F. 1962. Biosynthesis pathways from acetate to natural products. Activity of the enzymes in rubber synthesis. Pure Appl. Chem., 14: 137.

- Mahato SB, Banerjee S and Podder S, 1989. Steroid transformation by microorganisms. Phytochemistry, 28(1):7.
- Maliga P, 1984. Isolation and characterization of mutants in plant cell culture. Ann.Rev.Plant.Physiol. 35: 519.
- Mandava NB, 1988. Plant Growth Promoting Brassinosteroids.

 Ann.Rev. Plant. Physiol. Plant Mol.Biol., 39:23.
- Mann JD 1978. Production of Solasodine for Pharmaceutical industry In " Advances in agronomy Vol. 30. (Brady NC ed) Academic Press Inc. N.Y., London.
- Marshall JG and Staba EJ, 1976. Steroids as artifact from Dioscorea deltoidea. Tissue culture, Lloydia, 39:84.
- Mathews BF, Shey SCH and Widholm JM, 1980. Mechanism of resistance of a selected carrot suspension culture to 5 (2-amino ethyl) -L-cysteine 2-pflanzenphysiol, 96:53.
- Mathis P and Sauer K, 1973. Chlorophyll formation in greening bean leaves during the early stages. Plant Physiol., 51: 115.
- Maurey KM and Golbeck JH, 1985. Modulation of 3-hydroxy 3 methylglutaryl co-enzyme A reductase (HMGR) activity in Ochromonas malhanensis. Plant physiol. 77 (Supplement): 45.
- Maurey K, Fred Wolf and Joha Golbek, 1986. HMGCoA Reductase activity in Ochromonas malhanensis.Plant Physiol.,82:523.
- McGaw BA, 1987. Cytokinin biosynthesis and metabolism. In plant hormones and their role in plant growth and development 76: (Davis PJ ed) Martinas, Nijhoff, Boston.
- Milborrow BV, 1974. The chemistry and physiology of abscisic

- acid. Ann. Rev. Plant. Physiol., 25:259.
- Mothes K, Schutte HR and Lickner, 1985. Biochemistry of alkaloids: 406. VCH Publishers, Florida.
- Mudd JB, 1980. In "The Biochemistry of plants. (stumpf RK ed) 4: 509. Academic Press New York.
- Murashige T and Skoog F, 1962. A revised medium for rapid growth and bioassay with tobacco tissue cultures. Physiologia plantarum, 15: 473.
- Nandi DL and Porter JW, 1964. Arch. Biochem. Biophysics;105:
- Nes WD 1984. Isopentenoids In" Plant Biochemistry and Function" (Nes WD, Fuller G and Tsai L eds): 267. Dekker, New York.
- Nes WD and Bach DTJ, 1985. Evidence for mevalonate shunt in a trachyophyte. Proc. R. Soc. of Lond. B; 225: 425.
- Nes WR and Mckean ML, 1977. Biochemisty of steroids and other isopentanoids. University Park Press, Baltimore, London, Tokyo.
- Nishi A, Yoshida A, Mori M and sugano N, 1974. Isolation of variant carrot cell lines with altered pigmentation. Phytochemistry, 13: 1653.
- Nishi, A and Tsuritani I, 1983. Effect of auxin on the metabolism of mevalonic acid in suspension cultured carrot Cells. Phytochemistry, 22: 399.
- Oba K, Kondo K, Doke N and Vritani I, 1985. Induction of 3-hydroxy 3 methylglutaryl CoA reductase in potato tubers

- after slicing fungal infection or chemical treatment, and same properties of the enzyme. Plant Cell Physiol., 26 (5):873.
- Ogura K, Shinka T and Seto S, 1972. J. Biochem., 72: 1101, Tokyo.
- Paleg LG, 1965. Physiological effects of gibberellins.

 Ann Rev. Plant Physiol., 16: 291.
- Paltauf F, Daun G, Zuder G and Hogenauer G, Schulz G and Siedel G, 1982. Biochem. Biophys. Acta., 712: 268.
- Parks LW, 1978. Metabolism of sterols in yeast. CRC. Crit.Rev. Microbiol. 6: 301.
- Parks LW, Bottana CDK and Rodriguer RJ, 1984. In "Isopentanoids in plants: Biochemistry and functions" (Nes WD, Fuller G and Tsai L, eds) P: 433. Dikker, New York.
- Park, RB, 1965. The chloroplast In" Plant biochemistry (Bonner J and Varner JE, eds) Academic Press, New York.
- Phinney BO & Spreng C, 1982. Chemical genetics and the gibberellin pathway in Zea mays L. In " Plant growth substances" (Wareing PF ed) P. 101: 110. Academic Press, London, New York.
- Pinto WJ, Lozano R and Nes WR, 1985. Inhibition of sterol biosynthesis by ergasterol and cholasterol in Saccharomyces cerevisiae. Biochem Biophys Acta: 836.
- Potty VH and Bruemmer JH, 1970. Phytochemistry, 9:99.
- Quadri LEN and Giuletti AM, 1990. Effect of elicitation on Solasodine production by Solanum eleagrifolium cav.

- immobilized cells. In Procedding of VIIth Intl. Congress on plant tissue and cell culture. Held in Amsterdam, pp: 354.
- Qureshi N & Porter JW, 1981. Conversion of acetyl co-enzyme

 A to isopentyl pyrophosphate. In Biosynthesis of
 isoprenoid compounds. (Porter JW and Purgein SL, eds), 1:
 47. Wiley, New York.
- Rahier A et al 1985. Inhibitors of Δ^8 Δ^7 Sterol isomerase and of cycloencalemol obtusifoliol isomerase by N-Benzyl-8 Aza 4x-10-tans-decal-3 -o1, an analogue of a carbocationic high energy intermediate. Phytochemistry, 24(6): 1223.
- Rahier A, Taton M, Bourier-Nave P, Schmitt P, Benveniste P, Schuber F, Narula AS, Cattle L, Anging C and Place P,1986 Lipids, 21: 52
- Rao PS and Narayanaswamy S, 1968. Induced morphogenesis in tissue culture of Solanum xanthocarpum. Planta, 81: 372.
- Redway LA and Indra K Vasil, 1990. Selection of S-(2-Aminoethyl)-L-cysteine tolerance in embryogenic calli and regenerated plants of pennisetum purpureum Schum. Plant Science, 67: 203.
- Reinert J and Yeoman MM, 1983. Plant Cell and tissue culture
 A laboratory manual Narosa Publishing House, New Delhi
 and Springer Verlag Berlin Heidelberg, New York.
- Ripperger H and Schreiber K, 1981. Solanum Steroid alkaloids. In: "encyclopedia of plant physiology, New

- Series" (Peasson A and Zimmerman MH eds) Vol.8, P.167. A.P., N.Y.
- Robert Duperon, Thiersault M and Duperon P,, 1984. Highlevel of glycosylated sterols in Sp. of Solanum and strol changes during the development of tomato. Phytochemistry, 23(4): 742.
- Rogers LJ, Shah SPJ & Goodwin TW, 1966. Intracellular localization of mevalonate activating enzymes in plantcells. Biochem. J., 99: 381.
- Roth PS, Bach TJ & Thompson MJ 1989. Brassica steroids: potent inhibitors of growth of transformed Tobacco callus cultures. Plant Science, 59: 63.
- Rudney H, 1977. In "Biomedical and clinical aspects of co-enzyme Q (Flokers K and Yamamura Y eds): 29. Elsvier, Amsterdam.
- Russell DW, 1985. 3-Hydroxy-3-methylgluryl CoA Reductases from pea seedlings. Methods in Enzymology, 110: 26.
- Russell DW and Davidson H, 1982. Regulation of cystolic HMG-CoA reductase activity in pea seedlings. Contrasting responses to different hormones and hormone product interaction suggest hormonal modulation of activity. Biochem. Biophys. Res. Commu, 104: 1537.
- Sabine JR, 1983. Monographs on enzyme biology: HMG CoA reductase (Sabine JR ed) CRC Press, Boca Raton.
- Sakuta M and Komamine A, 1987. Cell growth and Accumulation of Secondary metabolites. Cell culture and somatic cell genetics of plants, Vol.4: 97.

- Sasse F, Buchholz M and Berlin J, 1983b. Selection of cell lines of <u>Catharanthus roseus</u> with increased tryptophan decarboxylase activity. Z. Naturforsch., C: Biosci. 38C: 916.
- Saxena PK, Gill R, Rashid A and Maheshwari SC, 1982.

 Plantlets from mesophyll protoplasts of Solanum xanthocarpum. Plant Cell Rep., 1: 219.
- Scheinpflug H and Kuc KH, 1987. In: Modern Selective fungicide. (Lyr H ed.), Verlag, Jana: P 173.
- Short KC and Torrey JG, 1972. Cytokinins in seedling roots roots of pea. Plant Physiol.,49: 155.
- Schreopfer G Jr, 1981. Sterol Biosynthesis. Annu.Rev. Biochem, 50, 585.
- Schulze Siebert D and Schultz G, 1985. Plant Physiol, 77: 1305.
- Selvapandiyan A, Subramani J, Bhatt PN and Mehta AR, 1988. A simple method for direct transplantation of cultured plants to the field. Plant Science, 56: 81.
- Sipat AB, 1982a Hydroxymethylglutaryl CoA reductase (NaDPH) in the latex of <u>Hevea brasiliensis</u>. Phytochemistry, 21: 2613.
- Sipat AB, 1982b. Biochem. Biophys Acta, 705; 284.
- Sipat AB, 1985. Meth. Enzymol., 110: 40.
- Skrukrud C, Taylor SE, Hawkins DR and Calvin M, 1987.

 Triterpenoid biosynthesis in <u>Euphorbia lathyris</u>. In:

 Plant Lipids: biochemistry, structure and function.

- (Steumpf PK, Mudd JB and Nes WD, eds.), Plenum Press, New York, in press.
- Sponsel, VM, 1987. Gibberellin biosynthesis and metabolism.

 In :Plant hormones and their role in plant growth and development.(Davis PJ ed) Martinus Nijholf,

 Boston: 43.
- Spurgeon SL, Sathyamoorthy N and Porter JW, 1984. Arch. Biophys., 230: 446.
- Stermer BA and Bostock RM, 1987. Involvement of HMG CoA reductase in regulation of sesquiterpenoid phytoalexin synthesis in potato. Plant Physiol., 84: 404.
- Stilt M and ap Rees T, 1979. Phytochemistry, 18: 1905.
- Stoessl, A, Slothers JB and Ward EWB, 1976. Sesquiterpenoid stress compounds of the Solanaceae. Phytochemistry, 15: 855.
- Subramani J, 1991. Cellular selection and Immobilization of Solanum cells for the over production of steroids. Ph.D. Thesis, M.S.University, Baroda, India.
- Subramani J, Josekutty PC and Bhatt PN, 1989. Solasodine levels in <u>Solanum sisymbriifolium</u> Lam. Ind. J. of Expl. Biol, 27.
- Suga T and Hirata T, 1990. Biotransformation of exogenous substrates. Phytochemistry, 29: 2393.
- Suzuki H, Oba K and Uritani I, 1974. Improved synthesis of 3-hydroxy 3-methylglutaryl Coenzyme A. Agr. Biol. Chem., 39: 2053.

- Suzuki H, Oba K and Uritani I, 1975a. The occurrence and properties of 3-hydroxy-3-methylglutaryl coenzyme A reductase in sweet potato roots infected by <u>Ceratocystis</u> fimbriata. Physiol. Plant Pathol., 7: 265.
- Suzuki H, Oba K and Uritani I, 1975 b. Improved synthesis of 3-¹⁴C-3-hydroxy-3-methylglutaryl coenzyme A. Agric. Biol.Chenm., 39: 1675.
- Suzuki H, Oba K and Uritani I, 1976. Subcellular localization of 3-hydroxyl-3-methylglutaryl-coenzyme A reductase and other membrane-bound enzymes in sweet potato roots. Plant Cell Physiol.,17: 691.
- Tabata M et al, 1978. In: Frontiers of Plant Tissue Culture Proc. 4th Intl. Cong. Plant Tissue Cell Culture. (Thorpe TA, ed.), Univ. Calgary: P 381.
- Tal M and Nevo Y, 1973. Abnormal stomatal behaviour and root resistance and hormonal imbalance in three wilty mutants of tomato. Biochem. Genet, 8: 291.
 - Taten M, Benveniste P and Rahier A, 1988. Phytochemistry, 26: 385.
 - Tjamose E and Kuc J, 1982. Inhibition of steroid glycoalkaloid accumulation by arachidonic and eicosapenta enoic acids in potato. Science, 217: 542.
 - Uemura M and Yoshida S, 1984. Plant Physiol., 75: 818.
 - Unnikrishnan SK, Prakash L, Josekutty PC, Bhatt PN and Mehta AR,1991. Effect of NaCl salinity of somatic Embryo Development in <u>Sapindus</u> <u>trifoliatus</u> L. Journal of Experimental Botany, 42, (236) 401.

- Vogeli U and Chappel J, 1987. Induction of sesquiterpene cyclase and suppression of squalene synthetase in plant cell cultures treated with fungal elicitor. Plant Physiol,83:1291.
- Walsh RC and Sisler HD, 1982. Pestic. Biochem. Physiol.,18: 122.
- Walton DC, 1987. Abscisic acid biosynthesis and metabolism.

 In: Plant hormones and their role in plant growth and development (Davies PJ ed). Martinus Nijhoff, Boston: 113.
- Wangs Y and Faust M, 1988 Changes of fatty acids and sterols in apple buds during bud break induced by a plant bioregulator, thidiazurom. Physiologia plantarum, 72: 115.
- Weete JD, 1980. In: Lipid Biochemistry of fungi and other organisms. Plenum press, New York, p 224.
- Weete JD, Kulifaj M, Montant C, Nes WR and Sancholle M, 1985. Can. J. Microbiol., 31: 1127.
- Weiler EW, Kruger H and Zenk MH, 1980. Radioimmuno assay for determination of the steroidal alkaloid solasodine and related compounds in living plants and herbarium species. Planta Med., 39: 112.
- Widholm JM, 1972a. Anthranilate synthetase from 5-methyltryptophan susceptible and resistant cultured Daucus carrota cells. Biochem. Biophys. Acta., 279: 48.
- Widholm JM, 1972b. Cultured Nicotiana tabacum cells with an

- altered anthranilate synthetase which is less sensitive to feedback inhibition. Biochem. Biophys. Acta., 261: 52.
- Widholm JM, 1974. Cultured carrot cell mutants: 5 methyl tryptophan resistance trait carried from cell to plant and back. Plant Sci. Let., 3: 323.
- Widholm JM, 1976. Selection and characterization of carrot and tobacco cells resistant to cystine and proline analogues. Canad. J. Bot., 54: 1523.
- Widholm JM, 1977a. Relation between auxin autotrophy and tryptophan accumulation in cultured plant cells. Planta, 89: 103.
- Widholm JM, 1977b. Selection and characterisation of amino acid analogue resistant plant cell culture. Crop. Sci., 17: 597.
- Widholm JM, 1978. Selection and characterisation of <u>Daucus</u>
 <u>carrota</u>
 L. Cell line resistant to four amino acid
 analogues. J. Exp. Bot., 29 (122): 1111.
- Widholm JM, 1984. Isolation and characterisation of mutant plant cell cultures. In: Plant cell culture in crop improvement (Sen SK and Giules KL, eds.) Plenum Press, New York: 71.
- Wilhum WG and Kastens J, 1974. Solanum dulcamara L. Triterpenoide und sterine aus dem. Petrolathore extract der Blatter. Planta Medica 25: 115.
- Williams DH, Stone MJ, Huck PR and Rahman SK, 1989. Why are secondary metabolites (natural products) biosynthesized?

 In: Journal of Natural Products, 52: 1189.

- Wink M, 1987. Physiology of accumulation of secondary metabolites with special reference to Alkaloids. Cell culture and somatic cell genetics of plants, Vol.4:17
- Wititsuwannakul R, 1986. Diurnal variation of 3-hydroxy--3-methylglutaryl Coenzyme A reductase activity in latex of <u>Hevea brasiliensis</u> and its relation to rubber content. Experientia, 42: 44.
- Wong RJ, McCormack DK and Russel DM, 1982. Plastid 3-hydroxy -3-methylglutaryl Coenzyme A reductase has distinctive kinetic and regulatory features: Properties of the enzyme and positive phytochrome control of activity in pea seedlings. Arch. Biochem. Biophys., 216: 631.
- Yabuta T and Hayashi T, 1939. Biochemical studies on "bekane" fungus of rice. II. Isolation of "gibberellin" the active principle which makes the rice seedlings grow slenderly. J. Agric.Chem.Soc. (Japan) 15: 257.
- Yamamoto Y and Mizuguchi R, 1982. Selection of a high yielding and stable pigment producing strain in cultured Euphorbia milli cells. Theor. Appl. Genet. 61:13.
- Yu Ito R, Oba K and Uritani I, 1982. Some problems in assay method of HMG CoA reductase activity in sweet potato in the presence of other HMG CoA utilizing enzymes. Agric. Biol. Chem., 46: 2087.
- Zakelj M and John Goad L, 1982 Observations on the biosynthesis of 24-methyl cholesterol and 24-ethyl cholesterol by Zea mayze Phytochemistry, 22(9):1931.

- Zenk MH, El-shagi H, Arens H, Stockigt J, Weiler EW and Deus B, 1977. Plant tissue culture and Bio-technological application (Barz W, Reinhard E & Zenk MH eds) Springer Verlag, Berlin.
- Zenktler M, 1973. <u>In vitro</u> development of embryos and seedlings from pollen grains of <u>Solanum dulcamara</u> Z. Pflanzenphysiol, 69: 189.
- Ziv M, Shwartz A and Fleminger D 1987. Malfunctioning stomata in viterous leaves of carnation (Dianthus caryophyllus) plants propagated <u>in vitro</u>: implications for hardening Plant Sci. 52: 127.