

## List of Tables

---

<b>Table 3.1: Primer used for 16S rRNA gene amplification.....</b>	28
<b>Table 3.2: Reaction mixture for PCR.....</b>	29
<b>Table 3.3: Conditions of PCR.....</b>	29
<b>Table 3.4: NCBI BLAST result of 16S rRNA gene amplicon product through Sanger Sequencing.....</b>	33
<b>Table 4.1: Reaction system-1 for cDNA synthesis .....</b>	45
<b>Table 4.2: Reaction system-2 for cDNA synthesis .....</b>	45
<b>Table 4.3: Primer used for qRT-PCR for type 4 pili expression .....</b>	46
<b>Table 4.4: Reaction system used for qRT-PCR .....</b>	46
<b>Table 4.5: Condition used for qRT-PCR.....</b>	47
<b>Table 5.1: Antibiotics used in the study.....</b>	71
<b>Table 5.2: Reaction system-1 for cDNA for synthesis .....</b>	74
<b>Table 5.3: Reaction system-2 for cDNA synthesis .....</b>	74
<b>Table 5.4: Primer used for Stringent response and toxin-antitoxin genes.....</b>	75
<b>Table 5.5: Reaction system used for qRT-PCR .....</b>	76
<b>Table 5.6: Condition used for qRT-PCR.....</b>	76
<b>Table 5.7: Primers used for qRT-PCR.....</b>	80
<b>Table 5.8: Reaction system for qRT-PCR .....</b>	81
<b>Table 5.9: Reaction conditions used for qRT-PCR .....</b>	81
<b>Table 5.10: MIC of <i>P. aeruginosa</i> isolates .....</b>	83
<b>Table 5.11: Summary of genetic analysis of mutations in genes in clinical isolates .....</b>	98
<b>Table 5.12: MIC determination .....</b>	101