

LIST OF FIGURES

Figure	Captions	Page no
1.1	Structures of chitin, chitosan and cellulose	2
1.2	Molecular conformation of chitosan at the solid state: (a) two fold helix conformation with side view(above) and a sectional view (below), (b) eight fold helix conformation with side view(left) and a sectional view (right)	9
1.3	Sources of chitin	11
1.4	Flow chart for the manufacturing of chitin and chitosan	13
1.5	Crystalline structure of chitosan	16
2.1	FTIR spectrum of CHT	54
2.2	FTIR spectrum of CHT-MC	55
2.3	¹ H NMR spectrum of CHT	56
2.4	Intrinsic viscosity determination of chitosan	61
2.5	Intrinsic viscosity of different grades of chitosan solutions	66
2.6	FTIR spectra of CHT-MC and its low molecular weight derivative	67
2.7	FTIR spectra of CHT and its low molecular weight derivatives	67
2.8	Aggregation of chitosan molecules as a function of molecular weight and concentration	68
2.9	Viscosity of chitosan (CHT) solution as a function of storage time	71
2.10	Effect of storage time of chitosan (CHT) solution on molecular weight	72
2.11	Viscosity of CHT-D5 solution as a function of storage time	74
2.12	Relative Viscosity of chitosan solution as a function of sodium acetate concentration	76
2.13	Effect of sodium acetate on storage stability of chitosan solution	77
2.14	Scanning electron microphotographs(x1000) of (a) Cotton Fibre (control), (b) CHT-MC treated fibres, (c) CHT treated fibres (d) CHT treated and then prolong boiled cotton fibres, (e) CHT-D4 treated fibres and (f) CHT-D5 treated fibres	80
2.15	Whiteness index (WI) of chitosan treated cotton fabric as a function of molecular weight and concentration	82
2.16	Yellowness index (YI) of chitosan treated cotton fabric as a function of molecular weight and concentration	82

2.17	Brightness index (BI) of chitosan treated cotton fabric as a function of molecular weight and concentration	83
2.18	Stiffness of chitosan treated cotton fabric as a function of molecular weight and concentration	85
2.19	Absorbency of chitosan treated cotton fabric as a function of molecular weight and concentration	87
2.20	C.I.Direct Red 81 uptake of chitosan pretreated cotton fabric as a function of molecular weight and concentration	89
2.21	C.I.Direct Blue 71 uptake of chitosan pretreated cotton fabric as a function of molecular weight and concentration	90
2.22	Effect of electrolyte (sodium sulphate) on dyeing of chitosan (CHT) treated cotton fabric with C.I.Direct Red 81	93
2.23	Effect of electrolyte (sodium sulphate) on dyeing of chitosan (CHT) treated cotton fabric with C.I. Direct Blue 71	93
2.24	Effect of chitosan treatment on colour depth and fastness of direct dyed cotton fabric	96
2.25	Effect of chitosan treatment on colour depth and fastness of direct dyed cotton fabric	97
2.26	Effect of chitosan pretreatment on acid dyeing	98
2.27	Crosslinking of cellulose macromolecules by DMDHEU	99
2.28	Effect chitosan treatment on resistance against microbial attack of cotton fabric	104
2.29	Scanning electron microphotographs chitosan treated samples by pad-dry-alkali method(x1000) (a) Cotton Fibre (control), (b) CHT-MC treated fibres, (c) CHT treated fibres and (d) CHT-D5 treated fibres	107
3.1	Nano size on scale	120
3.2	Dissolution of chitosan in acetic acid/water solvent	126
3.3	Hydrodynamic spheres of chitosan molecules in solution	127
3.4	Chitosan-TPP complex formed as a result of ionic gelation	129
3.5	Schematic presentation of ionic gelation of chitosan with TPP	129
3.6	Stability of nanoparticles due to electrostatic repulsion between the same ionic charges	130
3.7	Experimental set up for the preparation of nano chitosan by ionic gelation method	131
3.8	Size distribution of nano chitosan by intensity : (a) CHTN (319.4 nm), (b) CHT-D2N (271.6 nm), (c) CHT-D3N (231.1 nm), (d) CHT-D4N (195.2 nm)	133

	and (e) CHT-D5N (110.7 nm)	
3.9	Particle size of chitosan as a function of intrinsic viscosity	134
3.10	Particle size distribution of nano chitosan as a function of preparation methods	138
3.11	Effect of TPP concentration on appearance CHTN dispersion	139
3.12	Effect of TPP concentration on particle size of CHTN	140
3.13	Relative viscosity of CHTN dispersion as a function of TPP concentration	140
3.14	Drop in viscosity from parent to nano chitosan solution as a function of molecular weight	142
3.15	Stability study: white globular residue formed by microbial attack on CHTN	143
3.16	Scanning electron micrographs (x1000) of (a) control cotton fibre, (b) CHT treated fibres, (c) CHTN (319.4 nm) treated cotton fibres, (d) CHT-D4N (195.2nm) treated cotton fibres and (e) CHT-D5N (110.7 nm) treated cotton fibres	144
3.17	Effect of particle size of nano chitosan on appearance of treated cotton fabric	146
3.18	Effect of particle size of nano chitosan on absorbency of treated cotton fabric	148
3.19	Effect of particle size of nano-chitosan on dye uptake of treated cotton fabric	150
3.20	Effect of particle size of nano-chitosan on dye uptake of treated cotton fabric	154
3.21	Effect of nano chitosan on resistance towards microbial attack	178
4.1	FTIR spectra of chitosan and trimethyl chitosan chloride	180
4.2	Conductometric titration of TMCHT1 Vs AgNO ₃	181
4.3	Conductometric titration of TMCHT2 Vs AgNO ₃	181
4.4	Conductometric titration of TMCHT3 Vs AgNO ₃	183
4.5	¹ H NMR spectrum of TMCHT 3	197
4.6	FTIR spectra of N- ethyl chitosan derivatives	197
4.7	FTIR spectra of N-butyl chitosan derivatives	198
4.8	FTIR spectra of N-dodecyl chitosan derivatives	199
4.9	¹ HNMR spectrum of N-Ethyl chitosan (1:2)	201
4.10	¹ HNMR spectrum of N- ethyl N, N-dimethyl chitosan chloride (1:2)	204
4.11	¹ HNMR spectrum of N-butyl chitosan (1:2)	205
4.12	¹ HNMR spectrum of N-butyl N, N-dimethyl chitosan chloride (1:2)	205
4.13	¹ HNMR spectrum of N- dodecyl chitosan (1:4)	208

4.14	¹ HNMR spectrum of <i>N</i> - dodecyl <i>N, N</i> -dimethyl chitosan chloride (1:4)	209
4.15	FTIR spectra of <i>N</i> -benzyl <i>N,N</i> -dimethyl chitosan chloride	217
4.16	FTIR spectra of <i>N</i> -(1-naphthyl) methylene <i>N,N</i> -dimethyl chitosan chloride derivative	217
4.17	¹ HNMR spectrum of <i>N</i> -Benzyl chitosan (1:4)	219
4.18	¹ HNMR spectrum of <i>N</i> - benzyl <i>N, N</i> - dimethyl chitosan chloride(1:4)	221
4.19	¹ HNMR spectrum of <i>N</i> -(1- naphthyl) methylene chitosan (1:4)	223
4.20	¹ HNMR spectrum of <i>N</i> -(1- naphthyl) methylene <i>N, N</i> dimethyl chitosan chloride (1:4)	224
4.21	Reduced viscosity (η_{red}) TMCHT solutions in presence of sodium acetate	232
4.22	Reduced viscosity (η_{red}) <i>N</i> - sub CHT solutions in presence of sodium acetate	233
4.23	Reduced viscosity (η_{red}) <i>N</i> - sub CHT solutions in absence of sodium acetate	234
4.24	Effect of quaternization on intrinsic viscosity of CHT derivatives	235
4.25	Polycations chain conformation as a function of polymer concentration and electrolyte	237
4.26	Effect of <i>N</i> -Sub CHT treatment on whiteness of cotton fabric	241
4.27	Effect of <i>N</i> -Sub CHT treatment on yellowness of cotton fabric	241
4.28	Effect of <i>N</i> -Sub CHT treatment on brightness of cotton fabric	242
4.29	Effect of <i>N</i> -Sub CHT treatment on yellowness of cotton fabric due to chlorine retention	246
4.30	Effect of <i>N</i> -Sub CHT treatment on fibre strength of cotton due to chlorine retention	247
4.31	Effect of <i>N</i> -sub CHT treatment on absorbency of cotton fabric	248
4.32	Effect of <i>N</i> -sub CHT treatment on direct dyeing of cotton fabric	251
4.33	Effect of <i>N</i> -sub CHT treatment on direct dyeing of cotton fabric	252
4.34	Effect of dye bath condition on direct dyeing of <i>N</i> -Sub CHT treated cotton fabric	253
4.35	Effect of dye bath condition on direct dyeing of <i>N</i> -Sub CHT treated cotton fabric	254
4.36	Drop in colour value of salt free dyeing from conventional dye bath in direct dyeing of <i>N</i> -Sub Q CHT treatment of cotton fabric	254
4.37	Effect of <i>N</i> -Sub CHT treatment on color depth of direct dyed cotton fabrics	258
4.38	Effect of <i>N</i> -Sub Q CHT derivatives treatment on color depth of direct dyed cotton fabrics	258

4.39	Effect of <i>N</i> -Sub CHT treatment on dyeing with acid dye	261
4.40	Effect of different chitosan and <i>N</i> - Sub CHT treatment on soiling of cotton fabric	264
4.41	Effect of different CHT and <i>N</i> - sub CHT treatment on degree of soiling of cotton	265
4.42	Effect of quaternized <i>N</i> - sub CHT treatment on yellowness index	267
4.43	Release of particulate soil from fibre surface	268
4.44	Effect different <i>N</i> - sub CHT treatments on resistance against microbial attack of cotton fabric (soil burial test)	270
4.45	Effect different <i>N</i> - sub CHT treatment on resistance against microbial attack of dyed cotton fabric (soil burial test)	274
4.46	Effect different quaternized <i>N</i> - sub CHT treatment on resistance against microbial attack of dyed cotton fabric (soil burial test)	274
5.1	Effect of calcium ions in dye bath on colour value of direct dyed cotton fabric	297
5.2	FTIR spectra of CHT and CHT-Ca complex residues: (a) CHT, (b) CHT-Ca, treatment time, 24hrs and (c) CHT-Ca, treatment time 96hrs	300
5.3	Effect of treatment time on extent of chelation of Ca^{+2} ions by different chelating agents	304
5.4	Effect of concentration of chelating agents on the extent of removal of Ca^{+2} ions for 1h treatment	306
5.5	Effect of concentration of chelating agents on the extent of removal of Ca^{+2} ions for 24h treatment	308
5.6	Effect of pH of CHT solution on chelation efficiency measured in terms sorption of Ca^{+2} ions from water	311
5.7	Effect of particle size on chelation efficiency of CHT for calcium ions	314
5.8	Effect of Cu (II) ions in the dye bath on the dyeing of cotton using direct and reactive dyes	317
5.9	FTIR spectrum of CHT	318
5.10	FTIR spectrum of CHT-Cu complex residue	318
5.11	Chelation behaviour of chitosan derivatives for Cu(II) ions	324
5.12	Effect of pH of CHT derivatives solutions on chelation efficiency for Cu(II) ions	327
5.13	Effect of concentration of chelating agents on chelation of Cu(II) ions for 1h treatment	329
5.14	Effect of concentration of chelating agents on chelation of Cu(II) ions for 24h	331

	treatment	
5.15	Effect of particle size of chitosan on extent of chelation Cu(II) ions	334
5.16	Calibration curves for dye solutions	338
5.17	Sorption kinetics of C.I.Direct Red 81 at neutral pH	340
5.18	Sorption kinetics of C.I.Acid Blue158 at neutral pH	342
5.19	Effect of NMP pretreatment on sorption ability of CHT direct and acid dyes at neutral pH	344
5.20	Sorption kinetics of C.I.Direct Red 81 at acidic pH	346
5.21	Sorption kinetics of C.I.Acid Blue158 at acidic pH	348
5.22	Effluents containing C. I. Direct Red 81 treated various adsorbents	348
5.23	Effluents containing C. I. Acid Blue 158 treated various adsorbents	349