

---

## List of Figures

<b>Figure 1.1:</b> A simplified diagram illustrating the plant immune system	9
<b>Figure 1.2:</b> <i>Magnaporthe oryzae</i> distribution map	10
<b>Figure 1.3:</b> Disease cycle of <i>M. oryzae</i>	12
<b>Figure 1.4:</b> Schematic showing the early invasion into host cell by blast fungus	13
<b>Figure 1.5:</b> Diagram depicting the array of tools for examining fungal pathogen genomes	15
<b>Figure 1.6:</b> Species tree of blast fungus showing its divergence into multiple host-specific lineages	18
<b>Figure 1.7:</b> Coinfection on common host plants as one of the potential mechanisms for genetic exchange between distinct <i>M. oryzae</i> lineages	20
<b>Figure 1.8:</b> MAX effectors AVR-PikD and AVR-Pia exhibit the same structural fold	24
<b>Figure 1.9:</b> The evolutionary dynamics of effectors	26
<b>Figure 1.10:</b> Fungal non-proteinaceous effectors (NPEs) illustrating potential effector-like functions at the interface of plant–fungal interactions	28
<b>Figure 1.11:</b> Secondary metabolites are produced via metabolic pathways, encoded by biosynthetic gene clusters	30
<b>Figure 1.12:</b> Population genomics to identify chemical effectors	32
<b>Figure 2.1:</b> Schematic showing the steps of mono-conidial isolation from a leaf infected with blast disease	43
<b>Figure 2.2:</b> Schematic depicting the Rep-PCR methodology	45
<b>Figure 2.3:</b> Schematic representing design of mating crosses between opposite mating type strains of <i>M. oryzae</i>	48
<b>Figure 3.1:</b> Representative images of blast disease infected host tissues	57
<b>Figure 3.2:</b> Sampling sites of blast disease in the current study	58
<b>Figure 3.3:</b> Colony morphology of different strains of <i>M. oryzae</i>	61
<b>Figure 3.4:</b> Colony characteristics of representative strains of <i>M. oryzae</i>	62
<b>Figure 3.5:</b> Molecular fingerprinting based on Pot2 Rep-PCR	64
<b>Figure 3.6:</b> Molecular finger printing based on Mg-SINE Rep-PCR	65
<b>Figure 3.7:</b> PCR amplification of <i>AVR-Pita</i> gene from various rice and non-rice <i>M. oryzae</i> isolates	66

---

---

<b>Figure 3.8:</b> Organization of enzymatic domains in the hybrid PKS-NRPS product of <i>ACE1</i> gene	66
<b>Figure 3.9:</b> PCR amplification of <i>ACE1</i> gene from various rice and non-rice <i>M. oryzae</i> isolates	67
<b>Figure 3.10:</b> PCR amplification of <i>AVR1-CO39</i> gene in different isolates	68
<b>Figure 3.11:</b> PCR amplification of <i>AVR-Pik</i> gene in various isolates	69
<b>Figure 3.12:</b> PCR amplification of <i>AVR-Pizt</i> gene in different <i>M. oryzae</i> strains	69
<b>Figure 3.13:</b> PCR amplification of <i>PWL2</i> gene in different strains	70
<b>Figure 3.14:</b> PCR amplification confirming the mating types in various isolates from rice and non-rice hosts	71
<b>Figure 3.15:</b> Dendrogram generated by taking into account presence/absence of AVR genes and mating types	75
<b>Figure 3.16:</b> Host invasion of representative field strains of <i>M. oryzae</i> on leaf sheaths of four different host plants	76
<b>Figure 3.17:</b> Pathogenicity tests of field strains of <i>M. oryzae</i>	78
<b>Figure 3.18:</b> Mating assay displaying sexual crosses of representative field strains	79
<b>Figure 3.19:</b> Perithecia development as a result of successful mating between opposite mating type strains	80
<b>Figure 3.20:</b> Key factors considered for classification of <i>M. oryzae</i> field strains and subsequent selection for comparative genomics analyses	81
<b>Figure 4.1:</b> Workflow depicting the analyses performed on whole genome sequencing data	86
<b>Figure 4.2:</b> Genetic divergence of Indian population of <i>M. oryzae</i>	89
<b>Figure 4.3:</b> Principal component analysis (PCA) based on total SNPs identified from various <i>M. oryzae</i> strains	90
<b>Figure 4.4:</b> Genomic comparison of total SNPs number in genomes belonging to different lineages	90
<b>Figure 4.5:</b> Whole genome distribution of SNP density shown in the different chromosomes	91
<b>Figure 4.6:</b> Whole genome distribution of Nucleotide diversity measured as Pi per bp, shown along the length of different chromosomes	92

---

---

<b>Figure 4.7:</b> Content of repetitive elements in the de novo assembled genomes from each strain	94
<b>Figure 4.8:</b> Distribution of <i>M. oryzae</i> specific transposable elements in various host-specific lineages	95
<b>Figure 4.9:</b> Proportions of secretory proteins and effectors proteins predicted based on functional annotation	99
<b>Figure 4.10:</b> Venn diagram showing the lineage-wise clustering of proteomes	100
<b>Figure 4.11:</b> Phylogenetic tree based on PAV in accessory genes shows divergence of host-specific lineages of <i>M. oryzae</i>	102
<b>Figure 4.12:</b> Phylogenetic analysis of FAE sequences from host-specific lineages of <i>M. oryzae</i>	103
<b>Figure 4.13:</b> Presence/absence polymorphisms of known effector genes in various lineages of the blast fungus	106
<b>Figure 5.1:</b> <i>M. oryzae</i> worldwide populations are organized in host-specific lineages	110
<b>Figure 5.2:</b> <i>M. oryzae</i> and <i>M. grisea</i> are evolutionarily distinct species adapted to different hosts	111
<b>Figure 5.3:</b> Workflow for exploring fungal biosynthetic diversity	116
<b>Figure 5.4:</b> Occurrence of BGCs associated with specific classes of SM in individual genomes of <i>M. oryzae</i> and <i>M. grisea</i>	117
<b>Figure 5.5:</b> Similarity network analysis of biosynthetic gene clusters (BGCs) from <i>M. oryzae</i> and <i>M. grisea</i>	118
<b>Figure 5.6:</b> Homology of melanin-associated <i>M. oryzae</i> and <i>M. grisea</i> gene cluster family with reference BGCs in MIBiG database	120
<b>Figure 5.7:</b> Analysis of homology between <i>ACE1</i> gene cluster family in <i>M. oryzae</i> and <i>M. grisea</i> and reference BGCs in MIBiG database	121
<b>Figure 5.8:</b> Analysis of homology between putative epipyriculol-associated gene cluster family in <i>M. oryzae</i> and <i>M. grisea</i> and reference BGCs in MIBiG database	122
<b>Figure 5.9:</b> Homology of putative Squalestatin S1-associated <i>M. oryzae</i> and <i>M. grisea</i> gene cluster family with reference BGCs in MIBiG database	123
<b>Figure 5.10:</b> Analysis of homology between Tenuazonic acid-associated gene cluster families in host-adapted <i>M. oryzae</i> and <i>M. grisea</i> isolates	124

---

---

<b>Figure 5.11:</b> BGC-O2 is predominantly present in <i>Oryza</i> -specific lineage of <i>M. oryzae</i>	126
<b>Figure 5.12:</b> BGC-TLE is predominantly present in <i>Triticum</i> , <i>Lolium</i> and <i>Eleusine</i> -specific lineages of <i>M. oryzae</i>	127
<b>Figure 5.13:</b> Comparison of core biosynthetic gene region of BGC-TLE shows the pseudogenization in NRPS gene belonging to <i>Triticum</i> lineage	128
<b>Figure 5.14:</b> BGC-O1 is predominantly present in <i>Oryza</i> lineage of <i>M. oryzae</i>	129
<b>Figure 5.15:</b> Genomic localization of BGC-O1 and synteny between the <i>Oryza</i> and non- <i>Oryza</i> lineages	131
<b>Figure 5.16:</b> Circos plots showing synteny among SM related genes between the representative strains from <i>Oryza</i> and <i>Eleusine</i> lineages	133
<b>Figure 5.17:</b> Evolutionary origin of the reducing polyketide synthase (PKS) MGG_08236 gene	135
<b>Figure 5.18:</b> BGC-O1 genes are specifically expressed during pathogenic development	137

---