

- Acharya, K. V., Dadhaniya, P., Shendage, A., Badne, A., & Chavan, A. (2019) Estuarine Fisheries resource of India. *International Journal of Sciences and Applied Research*, 6(1), 1–12.
- Adélaïde, A., Michaud, E., Stieglitz, T., Fromard, F., Gardel, A., Tavares, M., & Thouzeau, G. (2016). Brachyuran crab community structure and associated sediment reworking activities in pioneer and young mangroves of French Guiana, South America. *Estuarine, Coastal and Shelf Science*, 182, 60–71. <https://doi.org/10.1016/j.ecss.2016.09.003>
- Agardy, T., Alder, J., Dayton, P., Curran, S., Kitchingman, A., Wilson, M., Catenazzi, A., Restrepo, J., Birkeland, C., Blaber, S., Saifullah, S., Branch, G., Boersma, D., Nixon, S., Dugan, P., Davidson, N., & Vörösmarty, C. (2005). Chapter 19: “Coastal systems” . In R. Hasan, R. Scholes, & N. Ash (Eds.), *Ecosystems and Human Well-Being: Current State and Trends: Findings of the Condition and Trends Working Group of the Millennium Ecosystem Assessment*. Washington, DC: Island Press, pp. 513–549.
- Ahyong, S. T., Lowry, J. K., Alonso, M., Bamber, R. N., Boxshall, G. A., Castro, P., Gerken, S., Karaman, G. S., Goy, J. W., Jones, D. S., Meland, K., D. Christopher R., & Jörundur, S. (2011). Subphylum Crustacea Brünnich, 1772. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148(1). <https://doi.org/10.11646/zootaxa.3148.1.33>
- Albouy, C., Guilhaumon, F., Araújo, M. B., Mouillot, D., & Leprieux, F. (2012). Combining projected changes in species richness and composition reveals climate change impacts on coastal Mediterranean fish assemblages. *Global Change Biology*, 18(10), 2995–3003.
- Alcock, A. (1895a). Description of a new species of Oxyrhynch crab of the genus *Parthenope*. *Journal of the Asiatic Society of Bengal, Calcutta*, 64(2), 296–297.

- Alcock, A. (1895b). Materials for a carcinological fauna of India. No. 1. The Brachyura *Oxyrhyncha*. *Journal of the Asiatic Society of Bengal, Calcutta*, 64(2), 157–291.
- Alcock, A. (1896). Materials for a carcinological fauna of India. No. 2. The Brachyura *Oxystoma*. *Journal of the Asiatic Society of Bengal, Calcutta*, 65(2), 134–296.
- Alcock, A. (1898). Materials for a carcinological fauna of India. No. 3. The Brachyura Cyclometopa. Part 1. The family Xanthidae. *Journal of the Asiatic Society of Bengal, Calcutta*, 67(1), 67–233.
- Alcock, A. (1899). Materials for a carcinological fauna of India. No. 4. The Brachyura Cyclometopa. Part II. A revision of the Cyclometopa with an account of the families Portunidae, Cancridae and Corystidae. *Journal of the Asiatic Society of Bengal, Calcutta*, 68(2), 1–104.
- Alcock, A. (1900a). Materials for a carcinological fauna of India. No. 5. Brachyura Primigenia or Dromicea. *Journal of the Asiatic Society of Bengal, Calcutta*, 68(5), 123–169.
- Alcock, A. (1900b). Materials for a carcinological fauna of India. No. 6. Brachyura Catometopa or Grapsoidea. *Journal of the Asiatic Society of Bengal, Calcutta*, 69(3), 279–486.
- Alcock, A. (1901). *Catalogue of the Indian Decapod Crustacea in the collection of the Indian Museum. Part I. Brachyura. Fasciculus I. Introduction and Dromides or Dromiacea (Brachyura Primigenia)*. The Indian museum, Calcutta. pp. 1–80.
- Allen, B. J., Rodgers, B., Tuan, Y., & Levinton, J. S. (2012). Size-dependent temperature and desiccation constraints on performance capacity: Implications for sexual selection in a fiddler crab. *Journal of Experimental Marine Biology and Ecology*, 438, 93–99.
<https://doi.org/10.1016/j.jembe.2012.09.009>

- Allen, C. J., Paterson, G. L., Hawkins, L. E., Hauton, C., Clark, P. F., & Aryuthaka, C. (2010). Zonation on sandy tropical beaches: a case study using *Dotilla intermedia* (Brachyura: Ocypodidae). *Marine Ecology Progress Series*, 408, 97–107.
- Allen, G. W. (1963). Estuarine destruction: A monument to progress. *Transactions of the North American Wildlife and Natural Resources Conference*, 29, 324–331.
- Altevogt, R. (1955). Some studies on two species of Indian fiddler crabs, *Uca marionis nitidus* (Dana) and *U. annulipes* (Latr.). *Journal of the Bombay Natural History Society*, 52(4), 1–15.
<https://cir.nii.ac.jp/crid/1570291224534090624>
- Amouroux, J. M., & Tavares, M. T. (2005). Natural recovery of Amazonian mangrove forest as revealed by Brachyuran crab fauna: preliminary description. *HAL (Le Centre Pour La Communication Scientifique Directe)*, 71–79.
- Amrutha, M. M., & Kumar, S. V. (2017). Characteristics of high monsoon wind-waves observed at multiple stations in the eastern Arabian Sea. *Ocean Science Discussions*, 1–30.
- Ansell, A. D. (1988). Migration or Shelter? Behavioural Options for Deposit Feeding Crabs on Tropical Sandy Shores. *Springer EBooks*, 151, 15–26.
https://doi.org/10.1007/978-1-4899-3737-7_2
- Araújo, J. M. C., Otero, X. L., Marques, A. G. B., Nóbrega, G. N., Silva, J. R. F., & Ferreira, T. O. (2011). Selective geochemistry of iron in mangrove soils in a semiarid tropical climate: effects of the burrowing activity of the crabs *Ucides cordatus* and *Uca maracoani*. *Geo-Marine Letters*, 32(4), 289–300.
<https://doi.org/10.1007/s00367-011-0268-5>
- Atkinson, R. J. A., & Chapman, C. J. (1984). Resin casting: A technique for investigating burrows in sublittoral sediments. *Progress in Underwater Science*, 9, 15–25.

- Atkinson, R. J. A., & Taylor, A. C. (1998). Physiological ecology of burrowing decapods. In *Aspects of Decapod Crustacean Biology: the proceedings of a Symposium held at the Zoological Society of London on 8th and 9th April 1987* (pp. 201–226). Clarendon Press. New York: Oxford University Press, ISBN: 9780198540083.
- Attrill, M. J., & Rundle, S. D. (2002). Ecotone or Ecocline: Ecological Boundaries in Estuaries. *Estuarine, Coastal and Shelf Science*, 55(6), 929–936. <https://doi.org/10.1006/ecss.2002.1036>
- Backwell, P. R. Y., & Passmore, N. I. (1996). Time constraints and multiple-choice criteria in the sampling behaviour and mate choice of the fiddler crab, *Uca annulipes*. *Behavioral Ecology and Sociobiology*, 38(6), 407–416. <https://doi.org/10.1007/s002650050258>
- Bairagi, N. (1995). "Ocypodidae: Decapoda: Crustacea." Hugli Matla Estuary, West Bengal. Estuarine Ecosystem Series. Vol. 2., by Director ZSI, Kolkata, edited by Director ZSI, Kolkata, pp. 263–288.
- Bandekar, P. D., & Kakati, V. S. (2021). Annotated checklist of brachyuran crabs of Karwar mangroves, West Coast of Karnataka, India. *Indian Journal of Ecology*, 47(3), 650–654.
- Barnes, D. (2002). Ecology of subtropical hermit crabs in SW Madagascar: refuge-use and dynamic niche overlap. *Marine Ecology Progress Series*, 238(238), 163–172. <https://doi.org/10.3354/meps238163>
- Barnes, D. K. (1997). Ecology of tropical hermit crabs at Quirimba Island, Mozambique: vertical migration (tree climbing). *Marine Ecology Progress Series*, 158, 233–240.
- Barrass, R. (1963). The Burrows of *Ocypode ceratophthalmus* (Pallas) (Crustacea, Ocypodidae) on a Tidal Wave Beach at Inhaca Island, Mocambique. *The Journal of Animal Ecology*, 32(1), 73–85. <https://doi.org/10.2307/2518>

- Bartolini, F., Cimò, F., Fusi, M., Dahdouh-Guebas, F., Lopes, G. P., & Cannicci, S. (2011). The effect of sewage discharge on the ecosystem engineering activities of two East African fiddler crab species: Consequences for mangrove ecosystem functioning. *Marine Environmental Research*, 71(1), 53–61. <https://doi.org/10.1016/j.marenvres.2010.10.002>
- Batie, R. E. (1982). Population of the crab *Hemigrapsus oregonensis* (Grapsidae) in Yaquina Bay Estuary, Oregon. II. Vertical distribution, biomass, and production. *Northwest Science*, 56, 241–249.
- Baustian, M. M., Hansen, G. J. A., Kluijver, A. de, Robinson, K., Henry, E. N., Knoll, L. B., Rose, K. C., & Carey, C. C. (2014). Chapter 3: Linking the bottom to the top in aquatic ecosystems: mechanisms and stressors of benthic-pelagic coupling. In P. F. Kemp (Ed.), *Eco-DAS X Symposium Proceedings* (pp. 38–60). Association for the Sciences of Limnology and Oceanography. 10.4319/ecodas.2014.978-0-9845591-4-5.25
- Beheshti, K., Endris, C., Goodwin, P., Pavlak, A., & Wasson, K. (2021). Burrowing crabs and physical factors hasten marsh recovery at panne edges. *Plos one*, 17(1), e0249330. [doi: 10.1101/2021.03.17.435836]
- Bertics, V. J., & Ziebis, W. (2009). Biodiversity of benthic microbial communities in bioturbated coastal sediments is controlled by geochemical microniches. *The ISME Journal*, 3(11), 1269–1285. <https://doi.org/10.1038/ismej.2009.62>
- Bertini, G., Fransozo, A., & de Melo, G. A. S. (2004). Biodiversity of brachyuran crabs (Crustacea: Decapoda) from non-consolidated sublittoral bottom on the northern coast of São Paulo State, Brazil. *Biodiversity and Conservation*, 13(12), 2185–2207. <https://doi.org/10.1023/b:bioc.0000047900.96123.34>
- Bertness, M. D., & Miller, T. (1984). The distribution and dynamics of *Uca pugnax* (Smith) burrows in a new England salt marsh. *Journal of Experimental*

Marine Biology and Ecology, 83(3), 211–237.
[https://doi.org/10.1016/s0022-0981\(84\)80002-7](https://doi.org/10.1016/s0022-0981(84)80002-7)

Bhadra, S. (1995). Portunidae: Decapoda: Crustacea. In: *Director ZSI, Kolkata (Ed.), Hugli Matla Estuary, West Bengal. Estuarine Ecosystem Series. Vol. 2. Zoological Survey of India, Kolkata*, 2, pp. 249-262.

Bhadra, S. (1998) Crustacea: Decapoda: Grapsidae. In: *Director ZSI, Kolkata (Ed.), Fauna of West Bengal. State Fauna Series 3. Vol. 10. Zoological Survey of India, Kolkata*, pp. 405-415.

Bhat, M., Patel, K. J., & Trivedi, J. N. (2021). First record of *Metopograpsus cannicci* Innocenti, Schubart and Fratini, 2020 (Crustacea: Decapoda: Grapsidae) from India. *Journal of Animal Diversity*, 3(4), 44–48.
<https://doi.org/10.52547/jad.2021.3.4.6>

Blankensteyn, A. (2006). O uso do caranguejo maria-farinha *Ocypode quadrata* (Fabricius) (Crustacea, Ocypodidae) como indicador de impactos antropogênicos em praias arenosas da Ilha de Santa Catarina, Santa Catarina, Brasil. *Revista Brasileira de Zoologia*, 23(3), 870–876.
<https://doi.org/10.1590/s0101-81752006000300034>

Bolam, S. G. (2003). Spatial patterns of estuarine macrobenthic assemblages: relationship with hydrodynamic regime. Estuarine Process Research Project, CEFAS, pp. 1–27.

Bopp, J. J., Sclafani, M., Frisk, M. G., McKown, K., Ziegler-Fede, C., Smith, D. R., & Cerrato, R. M. (2021). Telemetry reveals migratory drivers and disparate space use across seasons and age-groups in American horseshoe crabs. *Ecosphere*, 12(10), e0381. [doi: 10.1002/ecs2.3811]

Botto, F., & Iribarne, O. (2000). Contrasting Effects of Two Burrowing Crabs (*Chasmagnathus granulata* and *Uca uruguayensis*) on Sediment Composition and Transport in Estuarine Environments. *Estuarine, Coastal and Shelf Science*, 51(2), 141–151.
<https://doi.org/10.1006/ecss.2000.0642>

- Botto, F., Iribarne, O. O., Martínez, M. M., Delhey, K., Carrete, M., & Martinez, M. M. (1998). The Effect of Migratory Shorebirds on the Benthic Species of Three Southwestern Atlantic Argentinean Estuaries. *Estuaries*, 21(4), 700–709. <https://doi.org/10.2307/1353274>
- Botto, F., Iribarne, O., Gutierrez, J., Bava, J., Gagliardini, A., & Valiela, I. (2006). Ecological importance of passive deposition of organic matter into burrows of the SW Atlantic crab *Chasmagnathus granulatus*. *Marine Ecology Progress Series*, 312, 201-210. [doi: <https://doi.org/10.3354/MEPs312201>]
- Botto, L. D., Lisi, A., Robert-Gnansia, E., Erickson, J. D., Vollset, S. E., Mastroiacovo, P., Botting, B., Cocchi, G., de Vigan, C., de Walle, H., Feijoo, M., Irgens, L. M., McDonnell, B., Merlob, P., Ritvanen, A., Scarano, G., Siffel, C., Metneki, J., Stoll, C., & Smithells, R. (2005). International retrospective cohort study of neural tube defects in relation to folic acid recommendations: are the recommendations working? *BMJ (Clinical Research Ed.)*, 330(7491), 571. <https://doi.org/10.1136/bmj.38336.664352.82>
- Bouchet, P. (2006). THE MAGNITUDE OF MARINE BIODIVERSITY. In C. M. Duarte (Ed.), *The Exploration of Marine Biodiversity Scientific and Technological Challenges* (pp. 31–62). Natural History Museum, Paris, France.
- Boxshall, G., & Hayes, P. (2019). Biodiversity and Taxonomy of the Parasitic Crustacea. *Parasitic Crustacea*, 3, 73–134.
- Braga, A. A., Fransozo, A., Bertini, G., & Fumis, P. B. (2007). Bathymetric distribution and recruitment of the spider crab *Libinia spinosa* H. Milne Edwards 1834 in the Ubatuba and Caraguatatuba regions, northern coast of São Paulo, Brazil (Crustacea, Brachyura, Majoidea, Pisidae). *Senckenbergiana biologica*, 7-16.
- Braithwaite, R., & Talbot, M. R. (1972). Crustacean burrows in the Seychelles, Indian Ocean. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 11(4), 265–285. [https://doi.org/10.1016/0031-0182\(72\)90048-x](https://doi.org/10.1016/0031-0182(72)90048-x)

- Brown, A. C., & McLachlan, A. (1990). *The Ecology of Sandy Shores* (pp. 1–387). Elsevier Publishers, Amsterdam.
- Bruno, S. A., Pinheiro, M. A. A., & Colpo, K. D. (2012). Sediment Bioturbation Potential of *Uca rapax* and *Uca uruguayensis* as a Result of Their Feeding Activity. *Journal of Crustacean Biology*, 32(2), 223–229. <https://doi.org/10.1163/193724011x615451>
- Brurford, F. R., McGregor, P. K., & Oliveira, R. F. (2000). The sequence of mudball placement by male fiddler crabs, *Uca tangeri*. *Etología: Revista de La Sociedad Española de Etología*, 8(8), 53–55.
- Buchanan, B. A., & Stoner, A. W. (1988). Distributional patterns of blue crabs (*Callinectes* sp.) in a tropical estuarine lagoon. *Estuaries*, 11, 231–239.
- Buchanan, J. B., Sheader, M., & Kingston, P. F. (1978). Sources of variability in the benthic macrofauna off the south Northumberland coast, 1971–1976. *Journal of the Marine Biological Association of the United Kingdom*, 58(1), 191–209.
- Burggren, W., & McMahon, B. R. (2009). *Biology of the land crabs*. Cambridge University Press.
- Calcagno, J. A., Lovrich, G. A., Thatje, S., Nettelmann, U., & Anger, K. (2005). First year growth in the lithodids *Lithodes santolla* and *Paralomis granulosa* reared at different temperatures. *Journal of Sea Research*, 54(3), 221–230. [doi: <https://doi.org/10.1016/j.seares.2005.04.004>]
- Calcagno, J. A., Thatje, S., Anger, K., Lovrich, G. A., & Kaffenberger, A. (2003). Changes in biomass and chemical composition during lecithotrophic larval development of the southern stone crab *Paralomis granulosa*. *Marine Ecology Progress Series*, 257, 189–196. [doi: <https://doi.org/10.3354/meps257189>]
- Cameron, A. M. (1966). Some Aspects of the Behaviour of the Soldier Crab, *Mictyris longicarpus*. *Pacific Science*, 20, 224–234.

- Candisani, L. C., Sumida, P. Y. G., & Pires-Vanin, A. M. S. (2001). Burrow morphology and mating behaviour of the thalassinidean shrimp *Upogebia noronhensis*. *Journal of the Marine Biological Association of the United Kingdom*, 81(5), 799–803. <https://doi.org/10.1017/s0025315401004611>
- Cannicci, S., Paula, J., & Vannini, M. (1999). Activity pattern and spatial strategy in *Pachygrapsus marmoratus* (Decapoda: Grapsidae) from Mediterranean and Atlantic shores. *Marine Biology*, 133(3), 429–435.
- Canonico, G., Buttigieg, P. L., Montes, E., Muller-Karger, F. E., Stepien, C., Wright, D., Benson, A., Helmuth, B., Costello, M., Sousa-Pinto, I., Saeedi, H., Newton, J., Appeltans, W., Bednaršek, N., Bodrossy, L., Best, B.D., Brandt, A., Goodwin, K.D., Iken, K., Marques, A.C., Miloslavich, P., Ostrowski, M., Turner, W., Achterberg, E.P., Barry, T., Defeo, O., Bigatti, G., Henry, L. A., Ramiro-Sánchez, B., Durán, P., Morato, T., Roberts, J.M., García-Alegre, A., Cuadrado, M.S. and Murton, B. (2019) Global Observational Needs and Resources for Marine Biodiversity. *Frontiers in Marine Science* 6, 367. [doi: 10.3389/fmars.2019.00367]
- Chakrabarti, A. (1981). Burrow patterns of *Ocypode ceratophthalma* (Pallas) and their environmental significance. *Journal of Paleontology*, 187, 113–130.
- Chakrabarti, A., Chakrabarti, R., & Hertweck, G. (2006). Surface traces and bioturbate textures from bubbler crabs: an indicator of subtropical to tropical tidal flat environments. *Senckenbergiana Maritima*, 36(1), 19–27. <https://doi.org/10.1007/bf03043700>
- Chakraborty, S. K., & Choudhury, A. (1992). Ecological studies on the zonation of brachyuran crabs in a virgin mangrove island of Sundarbans, India. *Journal of Marine Biological Association of India*, 34(1-2), 189–194.
- Chan, B. K. K., Chan, K. K. Y., & Leung, P. C. M. (2006). Burrow Architecture of the Ghost Crab *Ocypode ceratophthalma* on a Sandy Shore in Hong Kong. *Hydrobiologia*, 560(1), 43–49. <https://doi.org/10.1007/s10750-005-1088-2>

- Chang, C.-W., & Iizuka, Y. (2012). Estuarine use and movement patterns of seven sympatric Mugilidae fishes: The Tatu Creek estuary, central western Taiwan. *Estuarine, Coastal and Shelf Science*, 106, 121–126. <https://doi.org/10.1016/j.ecss.2012.04.023>
- Chapman, M. G., & Underwood, A. J. (1996). Influences of tidal conditions, temperature and desiccation on patterns of aggregation of the high-shore periwinkle, *Littorina unifasciata*, in New South Wales, Australia. *Journal of Experimental Marine Biology and Ecology*, 196(1-2), 213–237. [https://doi.org/10.1016/0022-0981\(95\)00131-x](https://doi.org/10.1016/0022-0981(95)00131-x)
- Chatterjee, S., Mazumdar, D., & Chakraborty, S. K. (2014). Ecological role of fiddler crabs (*Uca spp.*) through bioturbatory activities in the coastal belt of East Midnapore, West Bengal, India. *Journal of the Marine Biological Association of India*, 56(2), 16–25. <https://doi.org/10.6024/jmbai.2014.56.2.01781-03>
- Cheung, W. W., Lam, V. W., Sarmiento, J. L., Kearney, K., Watson, R., & Pauly, D. (2009). Projecting global marine biodiversity impacts under climate change scenarios. *Fish and fisheries*, 10(3), 235–251.
- Chhapgar, B. F. (1956). On the Breeding Habits and Larva Stages of Some Crabs of Bombay. *Records of the Zoological Survey of India*, 54(1-2), 33–52.
- Chhapgar, B. F., & Borgaonkar, S. S. (1985). Extension of range of the estuarine crab *Ilyoplax gangetica* (Kemp) to the west coast of India. *Journal of the Bombay Natural History Society*, 82(1), 226–228.
- Chhapgar, B. F., & Mundkur, T. (1995). Occurrence of the crab *Euxanthus exsculptus* (Herbst) in Gujarat. *Journal of Bombay Natural History Society*, 92(2), 286.
- Chhapgar, B.F. (1957a). On the Marine crabs (Decapoda: Brachyura) of Bombay State, part 1. *Journal of the Bombay Natural History Society*, 54(2), 399–439.

- Chhapgar, B.F. (1957b). On the Marine crabs (Decapoda: Brachyura) of Bombay State, part 2. *Journal of the Bombay Natural History Society*, 54(3), 503–549.
- Chhapgar, B.F. (1957c). Marine crabs of Bombay State. *Contribution No. 1 of the Taraporevala Marine Biological Station, Department of Fisheries, Bombay*, 89.
- Chhapgar, B.F. (1958). Additions to the crab fauna of Bombay State. *Journal of the Bombay Natural History Society*, 55 (3), 582–585.
- Chhapgar, B.F., Desai, B.G. & Patel, S.J. (2004). On two interesting marine crabs (Decapoda: Brachyura) from Mandvi, Kutch. *Journal of the Bombay Natural History Society*, 101 (1), 184–186.
- Chopra, B. (1933). Further Notes on Crustacea Decapoda in the Indian Museum. III. On the Decapod Crustacea Collected by the Bengal Pilot Service off the Mouth of the River Hughli. Dromiacea and Oxystomata. *Records of the Zoological Survey of India*, 35(1), 25–52.
<https://doi.org/10.26515/rzsi/v35/i1/1933/162600>
- Christy, J. H., & Salmon, M. (1984). Ecology and Evolution of Mating Systems of Fiddler Crabs (Genus *Uca*). *Biological Reviews*, 59(4), 483–509.
<https://doi.org/10.1111/j.1469-185x.1984.tb00412.x>
- Christy, J. H. (1982). Burrow structure and use in the sand fiddler crab, *Uca pugilator* (Bosc). *Animal Behaviour*, 30(3), 687–694.
[https://doi.org/10.1016/s0003-3472\(82\)80139-5](https://doi.org/10.1016/s0003-3472(82)80139-5)
- Christy, J. H. (1987). Female Choice and the Breeding Behavior of the Fiddler Crab *Uca beebei*. *Journal of Crustacean Biology*, 7(4), 624–635.
<https://doi.org/10.2307/1548648>
- Christy, J. H. (2010). Pillar Function in the Fiddler Crab *Uca beebei* (I): Effects on Male Spacing and Aggression. *Ethology*, 78(1), 53–71.
<https://doi.org/10.1111/j.1439-0310.1988.tb00219.x>

- Christy, J. H., & Schober, U. M. (1994). A test for resource-defense mating in the fiddler crab *Uca beebei*. *Animal Behaviour*, 48(4), 795–802. <https://doi.org/10.1006/anbe.1994.1303>
- Clayton, D. A. (1988). Hood construction as a spacing mechanism in *Cleistostoma kuwaitense* (Crustacea: Ocypodidae). *Marine Biology*, 99(1), 57–61. <https://doi.org/10.1007/bf00644977>
- Cloern, J. E., Abreu, P. C., Carstensen, J., Chauvaud, L., Elmgren, R., Grall, J., Greening, H., Johansson, J. O. R., Kahru, M., Sherwood, E. T., Xu, J., & Yin, K. (2016). Human activities and climate variability drive fast-paced change across the world's estuarine-coastal ecosystems. *Global Change Biology*, 22(2), 513–529. <https://doi.org/10.1111/gcb.13059>
- Costello, M. J., & Chaudhary, C. (2017). Marine biodiversity, biogeography, deep-sea gradients, and conservation. *Current Biology*, 27(11), R511–R527. [doi: <https://doi.org/10.1016/j.cub.2017.04.060>]
- Cott, H. B. (1929). Observations on the natural history of the racing-crab *Ocypode ceratophthalma* from Beira. *Proceedings of the Zoological Society of London*, 99, 755–765.
- Crane, J. (1941a). Eastern Pacific Expeditions of the New York Zoological Society. XXVI. Crabs of the Genus *Uca* from the West Coast of Central America. *Zoologica: Scientific Contributions of the New York Zoological Society*, 26(19), 145–208. <https://doi.org/10.5962/p.184677>
- Crane, J. (1941b). Eastern Pacific Expeditions of the New York Zoological Society. XXIX. On the growth and ecology of Brachyuran crabs of the genus *Ocypode*. *Zoologica; Scientific Contributions of the New York Zoological Society*, 26(29). <https://doi.org/10.5962/p.203597>
- Crane, J. (1975). *Fiddler Crabs of the World* OCYPODDAE: GENUS *Uca*. Princeton University Press, Princeton, pp. 736.

- Cranwell, L., & Cranwell, L. (1938). Intertidal Communities of the Poor Knights Islands, New Zealand. *Transactions and Proceedings of the Royal Society of New Zealand*, 67, 375–407.
- Dabhi, N., & Gohil, B. (2022). Checklist of intertidal fauna in muddy-rocky ecosystem, gopnath coast, gulf of Khambhat, India. *International Journal of Zoology Studies*, 7(1), 74–79.
- Das, A.K & Dev Roy, M.K. (1989). A general account of the mangrove fauna of Andaman and Nicobar Islands India. In: *Director ZSI, Kolkata (Ed.), Fauna of Conservation Areas. Vol. 4. Zoological Survey of India, Kolkata*, pp. 1-178.
- Day, J. H. (1981). Summaries of current knowledge of 43 estuaries in southern Africa. In J. H. Day (Ed.), *Estuarine Ecology with Particular Reference to Southern Africa*. (Vol. 23, Issue 3, pp. 251–329). Cape Town: A.A. Balkema, Rotterdam. [https://doi.org/10.1016/0006-3207\(82\)90080-5](https://doi.org/10.1016/0006-3207(82)90080-5)
- de, C. (2000). Neoichnological activities of endobenthic invertebrates in downdrift coastal Ganges delta complex, India: Their significance in trace fossil interpretations and paleoshoreline reconstructions. *Ichnos*, 7(2), 89–113. <https://doi.org/10.1080/10420940009380149>
- Deb, M. (1999). Crustacea: Decapoda: Crabs. In: *Director ZSI, Kolkata (Ed.), Fauna of West Bengal. State Fauna Series. Vol. 10. Part 3. Zoological Survey of India, Kolkata*, 345–403.
- Denny, M. W. (1988). Biology and the mechanics of the wave-swept environment. *Princeton University Press, Paris*, 15.
- Dev Roy, M. K. (2013). Diversity and Distribution of Marine Brachyuran Crab Communities Inhabiting West Coast of India. In K. Venkataraman, C. Sivaperuman, & C. Raghunathan (Eds.), *Ecology and Conservation of Tropical Marine Faunal Communities*. (pp. 147–169). Springer, Berlin, Heidelberg.
- Dev Roy, M. K., Nandi, N. C. & Khan, R. A. (2009). Invertebrate diversity. In: Director ZSI, Kolkata (Ed.), *Faunal Diversity of Vembanad Lake. A Ramsar site in*

- Kerala, India. Wetland Ecosystem Series. Vol. 10. Zoological Survey of India, Kolkata, pp. 69-128.
- Dev Roy, M.K. & Bhadra, S. (2001). Brachyuran crabs (Crustacea: Decapoda: Brachyura). *In: Director ZSI, Kolkata (Ed.), Estuarine Ecosystem Series. Vol. 4. Zoological Survey of India, Kolkata*, 35–54.
- Dev Roy, M.K. & Bhadra, S. (2008). Marine and estuarine crabs (Crustacea: Decapoda: Brachyura). *In: Director ZSI, Kolkata (Ed.), Fauna of Goa. State Fauna Series. Vol 16. Zoological Survey of India, Kolkata*, 109–154.
- Dev Roy, M.K. & Bhadra, S. (2011). Brachyuran crabs (Crustacea: Decapoda: Brachyura). *In: Director ZSI, Kolkata (Ed.), Fauna of Tamil Nadu. State Fauna Series. Vol. 17. Part 2. Zoological Survey of India, Kolkata*, 109–269.
- Dev Roy, M.K. & Das, A.K. (2000). Taxonomy, ecobiology and distribution pattern of the Brachyuran crabs of mangrove ecosystem in Andaman Islands. *Records of Zoological Survey of India, Occasional Paper*, 185, 1–211.
- Dev Roy, M.K. & Nandi, N.C. (2007a). Brachyuran diversity in coastal ecosystems of Tamil Nadu. *Journal of Environment and Sociobiology*, 4 (2), 169–192.
- Dev Roy, M.K. & Nandi, N.C. (2007b). Brachyuran bioresources of coastal Andhra Pradesh. *In: Director ZSI, Kolkata (Ed.), National symposium on Conservation and Validation of Marine Biodiversity. Zoological Survey of India, Kolkata*, 53– 66.
- Dev Roy, M.K. & Nandi, N.C. (2008). Brachyuran biodiversity of some selected brackish water lakes of India. *In: Sengupta, M. & Dalwani, R. (Eds.), Proceedings of Taal 2007, The 12th World Lake Conference*, 496–499.
- Dev Roy, M.K. & Nandi, N.C. (2012). Brachyuran crabs (Crustacea). *In: Director ZSI, Kolkata (Ed.), Fauna of Andaman and Nicobar Islands. State Fauna Series. Vol. 19. Part 1. Zoological Survey of India, Kolkata*, 185–236.
- Dev Roy, M.K. & Rath, S. (2017). An inventory of crustacean fauna from Odisha Coast, India. *Journal of Environment and Sociobiology*, 14 (1), 49–112.

- Dev Roy, M.K. (2008). An annotated checklist of mangrove and coral reef inhabiting Brachyuran crabs of India. *Records of Zoological Survey of India, Occasional Paper*, 289, 1–212.
- Dev Roy, M.K. (2013). Diversity and distribution of marine brachyuran crab communities inhabiting West Coast of India. In: Venkataraman, K., Sivaperuman, C. & Raghunathan, C. (Eds.), *Ecology and Conservation of Tropical Marine Faunal Communities Part 1*. Springer, Berlin & Heidelberg, 147–169.
- Diaz, R. J., & Cutter, G. R. J. (2001). In situ measurement of organism-sediment interaction: rates of burrow formation, abandonment and sediment oxidation, reduction. In *Organism-sediment interactions* (pp. 19–32). University of South Carolina Press, Columbia.
- Diemont, W. H., & Van Wijngaarden, W. (1974). Sedimentation Patterns, Soils, Mangrove Vegetation and Landuse in the Tidal Areas of West-Malaysia. In *Proceedings of the International Symposium on Biology and Management of Mangroves* (pp. 513–528). University Press of Florida, Gainesville, Florida.
- Dittel, A. I., Hines, A. H., Ruiz, G. M., & K. Keith Ruffin. (1995). Effects of shallow water refuge on behavior and density-dependent mortality of juvenile blue crabs in Chesapeake Bay. *Bulletin of Marine Science*, 57(3), 902–916.
- Dixit, A. M., Kumar, P., Kumar, L., Pathak, K. D., & Patel, M. I. (2010). *Economic Valuation of Coral Reef Systems in Gulf of Kachchh*. Gujarat: Gujarat Ecology Commission. Final Report of World Bank Aided Integrated Coastal Zone Management (ICZM) Project.
- Dixon, R. W., Peters, S. L., & Townsend, C. G. (2015). Burrowing preferences of Atlantic ghost crab, *Ocypode quadrata*, in relation to sand compaction in Padre Island National Seashore, Texas. *Physical Geography*, 36(3), 188–201. <https://doi.org/10.1080/02723646.2015.1033182>
- Douglas, E. J., Pilditch, C. A., Lohrer, A. M., Savage, C., Schipper, L. A., & Thrush, S. F. (2018). Sedimentary Environment Influences Ecosystem Response to

- Nutrient Enrichment. *Estuaries and Coasts*, 41(7), 1994–2008.
<https://doi.org/10.1007/s12237-018-0416-5>
- Dubey, S. K., Chakraborty, D. P., Chakraborty, S., & Choudhury, A. (2013). Burrow architecture of red ghost crab *Ocypode macrocera* (H. Milne– Edwards, 1852): a case study in Indian Sundarbans. *Exploratory Animal and Medical Research*, 3(2), 136–144.
- Duineveld, G. C. A., Tselepides, A., Witbaard, R., Bak, R. P. M., Berghuis, E. M., Nieuwland, G., van der Weele, J., & Kok, A. (2000). Benthic–pelagic coupling in the oligotrophic Cretan Sea. *Progress in Oceanography*, 46(2-4), 457–481. [https://doi.org/10.1016/s0079-6611\(00\)00029-x](https://doi.org/10.1016/s0079-6611(00)00029-x)
- Dworschak, P. C. (1988). The Biology of *Upogebia pusilla* (PETAGNA) (Decapoda, Thalassinidea). *Marine Ecology*, 9(1), 51–77.
<https://doi.org/10.1111/j.1439-0485.1988.tb00198.x>
- Dworschak, P. C. (2001). The Burrows of *Callianassa tyrrhena* (Petagna 1792) (Decapoda: Thalassinidea). *Marine Ecology*, 22(1-2), 155–166.
<https://doi.org/10.1046/j.1439-0485.2001.00748.x>
- Edney, E. B. (1961). The Water and Heat Relationships of Fiddler Crabs (*Uca* spp.). *Transactions of the Royal Society of South Africa*, 36(2), 71–91.
<https://doi.org/10.1080/00359196109519035>
- Ellis, D. V. (2003). Rocky shore intertidal zonation as a means of monitoring and assessing shoreline biodiversity recovery. *Marine Pollution Bulletin*, 46(3), 305–307.
- Fabricius, J. C. (1775). Systema Entomologiae, sistens insectorum Classes, Ordines, Genera, Species, Adjectis Synonymies, Locis, Descriptionibus, Observationibus. *Libraria Kortii, Flensburgi and Lipsiae*, 832.
- Fabricius, J. C. (1787). Mantissa Insectorum sistens eorum species nuper detectas adiectis Characteribus genericis, Differentiis specificis, Emendationibus, Observationibus. *Hafniae*, 1, 348.

- Fabricius, J. C. (1793). *Entomologia Systematica Emendata et Aucta, Secundum, Classes, Ordines, Genera, Species, Adjectis Synonimas, Locis, Observationibus, Descriptionibus*. Vol. 2. impensis Christ. Gottl. Proft, Hafniae, 519 pp.
- Fielder, D. R. (1970). The feeding behaviour of the sand crab *Scopimera inflata* (Decapoda, Ocypodidae). *Journal of Zoology*, 160(1), 35–49. <https://doi.org/10.1111/j.1469-7998.1970.tb02896.x>
- Flint, R. W. (1981). Gulf of Mexico Outer Continental Shelf Benthos: Macroinfaunal-Environmental Relationships. *Biological Oceanography*, 1(2), 135-155.
- Flores, A. A., & Paula, J. (2001). Intertidal distribution and species composition of brachyuran crabs at two rocky shores in Central Portugal. In *Advances in Decapod Crustacean Research: Proceedings of the 7th Colloquium Crustacea Decapoda Mediterranea, held at the Faculty of Sciences of the University of Lisbon, Portugal, 6–9 September 1999* (pp. 171-177). Springer Netherlands.
- Flores, A. A., Abrantes, K. G., & Paula, J. (2005). Estimating abundance and spatial distribution patterns of the bubble crab *Dotilla fenestrata* (Crustacea: Brachyura). *Austral ecology*, 30(1), 14-23. [doi: 10.1111/j.1442-9993.2005.01409]
- Fortaleza, O. M., Maria Lemos Girão, M., Franklin Junior, W., Pinto de Lima, J., & De Almeida Rocha-Barreira, C. (2020). Which Moon Phase Do We Find More Ghosts? Effects of The Lunar Cycle on The Ghost Crab *Ocypode quadrata* (Fabricius, 1787). *Arquivos de Ciências Do Mar*, 52(2), 85–97. <https://doi.org/10.32360/acmar.v52i2.42737>
- François, F., Gerino, M., Stora, G., Durbec, J., & Poggiale, J. (2002). Functional approach to sediment reworking by gallery-forming macrobenthic organisms: modeling and application with the polychaete *Nereis diversicolor*. *Marine Ecology Progress Series*, 229, 127–136. <https://doi.org/10.3354/meps229127>

- Fransozo, A., Negreiros-Fransozo, M. L., & Mantelatto, F. L. (1992). Composition and distribution of brachyura (costacea, decapoda) from the unconsolidated sublittoral in Enada da Fortaleza, Ubatuba (SP). *Brazilian Journal of Biology*, 52(4), 667–675.
- Frederich, M., & Pörtner, H. O. (2000). Oxygen limitation of thermal tolerance defined by cardiac and ventilatory performance in spider crab, *Maja squinado*. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 279(5), R1531–R1538. <https://doi.org/10.1152/ajpregu.2000.279.5.r1531>
- George, R. W., & Knott, M. E. (1966). The *Ocypode* ghost crabs of western Australia (Crustacea, Brachyura). *Journal of the Royal Society of Western Australia*, 48, 15–21. [https://doi.org/10.1016/0011-7471\(66\)90061-1](https://doi.org/10.1016/0011-7471(66)90061-1)
- Gherardi, F., & Russo, S. (2001). Burrowing activity in the sand-bubbler crab, *Dotilla fenestrata* (Crustacea, Ocypodidae), inhabiting a mangrove swamp in Kenya. *Journal of Zoology*, 253(2), 211–223. <https://doi.org/10.1017/s0952836901000188>
- Gherardi, F., Russo, S., & Anyona, D. (1999). Burrow-orientated activity in the ocypodid crab, *Dotilla fenestrata*, living in a mangrove swamp. *Journal of the Marine Biological Association of the United Kingdom*, 79(2), 281–293. <https://doi.org/10.1017/s0025315498000319>
- Glémarec, M. (1973). The benthic communities of the European North Atlantic continental shelf. *Oceanography and Marine Biology: an Annual Review*, 11, 263–289.
- Gohil, B., & Kundu, R. (2012). Diversity of the intertidal macrofauna at West Coast of Gujarat, India. *Life Sciences Leaflets*, 12, 135–145.
- Golley, F., Odum, H. T., & Wilson, R. L. (1962). The structure and metabolism of Puertrican red mangrove forest in May. *Ecology*, 43, 9–18.

- Gopi, K. C., & Mishra, S. S. (2015). Diversity of Marine Fish of India. In K. Venkataraman & C. Sivaperuman (Eds.), *Marine Faunal Diversity in India* (pp. 171–193). Academic Press, Elsevier Inc., USA.
<https://doi.org/10.1016/b978-0-12-801948-1.00012-4>
- Graf, G. (1992). Benthic-pelagic coupling - a benthic view. *Oceanography and Marine Biology*, 30, 149–190.
- Granek, E. F., Polasky, S., Kappel, C. V., Reed, D. J., Stoms, D. M., Koch, E. W., Kennedy, C. J., Cramer, L. A., Hacker, S. D., Barbier, E. B., Aswani, S., Ruckelshays, M., Perillo, G. M. E., Silliman, B. R., Muthiga, N., Bael, D., & Wolanski, E. (2010). Ecosystem Services as a Common Language for Coastal Ecosystem-Based Management. *Conservation Biology*, 24(1), 207–216.
<https://doi.org/10.1111/j.1523-1739.2009.01355.x>
- Grave, S. D., Pentcheff, N. D., Ahyong, S. T., Chan, T., Crandall, K. A., Dworschak, P. C., Felder, D. L., Feldmann, R. M., Fransen, C. H. J. M., Goulding, L. Y. D., Lemaitre, R., Low, M. E. Y., Martin, J. W., Peter, Schweitzer, C. E., Tan, S. H., Tshudy, D., & Wetzler, R. (2009). A Classification of Living and Fossil Genera of Decapod Crustaceans. *Raffles Bulletin of Zoology*, 21, 1.
- Gravelly, F. H. (1927). Decapoda and Stomatopoda. In: *The littoral fauna of Krusadai Island in the Gulf of Manaar. Bulletin of the Madras Government Museum, New Series (Natural History)*, 1, 141–155.
- Gregory, M. R., & Ballanc, P. F. (1979). On How Some Rays (Elasmobranchia) Excavate Feeding Depressions by Jetting Water. *SEPM Journal of Sedimentary Research*, Vol. 49(4), 1125–1129.
<https://doi.org/10.1306/212f78c9-2b24-11d7-8648000102c1865d>
- Griffin, D. J. G. (1965). The Behaviour of Shore Crabs. *Australian Natural History*, 15, 87–91.
- Griffin, D. J. G. (1968). Social and maintenance behaviour in two Australian ocypodid crabs (Crustacea: Brachyura). *Journal of Zoology*, 156(3), 291–305. <https://doi.org/10.1111/j.1469-7998.1968.tb04353.x>

- Griffis, R. B., & Chavez, F. L. (1988). Effects of sediment type on burrows of *Callianassa californiensis* Dana and *C. gigas* Dana. *Journal of Experimental Marine Biology and Ecology*, 117(3), 239–253. [https://doi.org/10.1016/0022-0981\(88\)90060-3](https://doi.org/10.1016/0022-0981(88)90060-3)
- Griffis, R. B., & Suchanek, T. H. (1991). A model of burrow architecture and trophic modes in thalassinidean shrimp (Decapoda: Thalassinidea). *Marine Ecology Progress Series*, 79(1), 171–183. <https://doi.org/10.3354/meps079171>
- Groombridge, B., & Jenkins, M. D. (2000). *Global biodiversity: earth's living resources in the 21st century*. (p. 254). World Conservation Press, Cambridge, UK.
- Gross, W. J. (1964). Trends in water and salt regulation among aquatic and amphibious crabs. *The Biological Bulletin*, 127(3), 447–466.
- Gül, M. R., & Griffen, B. D. (2018). Impacts of human disturbance on ghost crab burrow morphology and distribution on sandy shores. *PLOS ONE*, 13(12), e0209977. <https://doi.org/10.1371/journal.pone.0209977>
- Gutiérrez, J. L., Jones, C. G., Groffman, P. M., Findlay, S. E. G., Iribarne, O. O., Ribeiro, P. D., & Bruschetti, C. M. (2006). The Contribution of Crab Burrow Excavation to Carbon Availability in Surficial Salt-marsh Sediments. *Ecosystems*, 9(4), 647–658. <https://doi.org/10.1007/s10021-006-0135-9>
- Hall, S., & Thatje, S. (2011). Temperature-driven biogeography of the deep-sea family Lithodidae (Crustacea: Decapoda: Anomura) in the Southern Ocean. *Polar biology*, 34, 363–370. [doi: <https://doi.org/10.1007/s00300-010-0890-0>]
- Halpern, B. S., Walbridge, S., Selkoe, K. A., Kappel, C. V., Micheli, F., D'Agrosa, C., Bruno, J. F., Casey, K. S., Ebert, C., Fox, H. E., Fujita, R., Heinemann, D., Lenihan, H. S., Madin, E. M. P., Perry, M. T., Selig, E. R., Spalding, M., Steneck, R., & Watson, R. (2008). A Global Map of Human Impact on Marine

Ecosystems. *Science*, 319(5865), 948–952.
<https://doi.org/10.1126/science.1149345>

Hamasaki, K., Matsui, N., & Nogami, M. (2011). Size at sexual maturity and body size composition of mud crabs *Scylla* spp. caught in Don Sak, Bandon Bay, Gulf of Thailand. *Fisheries Science*, 77, 49-57. [doi: 10.1007/s12562-010-0307-6]

Haragi, S.B., Naik, U.G. & Rathod, J.L. (2010). Brachyuran diversity in sub littoral zone of tropical estuary, Karwar, west coast of India. *Lake 2010: Wetlands, Biodiversity and Climate Change*, 1–12.

Harborne, A. R., & Mumby, P. J. (2011). Novel ecosystems: altering fish assemblages in warming waters. *Current Biology*, 21(19), R822-R824.

Hargrave, B. T. (1973). Coupling Carbon Flow Through Some Pelagic and Benthic Communities. *Journal of the Fisheries Research Board of Canada*, 30(9), 1317–1326. <https://doi.org/10.1139/f73-212>

Hartnoll, R. G. (1973). Factors affecting the distribution and behaviour of the crab *Dotilla fenestrata* on East African shores. *Estuarine and Coastal Marine Science*, 1(2), 137–152. [https://doi.org/10.1016/0302-3524\(73\)90066-2](https://doi.org/10.1016/0302-3524(73)90066-2)

Hartnoll, R. G. (1975). The Grapsidae and Ocypodidae (Decapoda: Brachyura) of Tanzania. *Journal of Zoology*, 177(3), 305-328.

Hattab, T., Albouy, C., Lasram, F. B. R., Somot, S., Le Loc'h, F., & Leprieux, F. (2014). Towards a better understanding of potential impacts of climate change on marine species distribution: a multiscale modelling approach. *Global Ecology and Biogeography*, 23(12), 1417-1429.

He, Q., & Cui, B. (2015). Multiple mechanisms sustain a plant-animal facilitation on a coastal ecotone. *Scientific reports*, 5(1), 8612.

Heck, J., Kenneth, L., & Coen, L. D. (1995). Predation and the Abundance of Juvenile Blue Crabs: A Comparison of Selected East and Gulf Coast (USA) Studies. *Bulletin of Marine Science*, 57(3), 877–883.

<https://www.ingentaconnect.com/contentone/umrmsas/bullmar/1995/00000057/00000003/art00025#expand/collapse>

Henderson, J. R. (1893). A contribution to Indian carcinology. *Transactions of the Linnaean Society of London Zoology, Series 2*, 5, 325–458. [doi: <https://doi.org/10.1111/j.1096-3642.1893.tb00653.x>]

Herman, P. M. J., Middelburg, J. J., & Heip, C. H. R. (2001). Benthic community structure and sediment processes on an intertidal flat: results from the ECOFLAT project. *Continental Shelf Research*, 21(18-19), 2055–2071. [https://doi.org/10.1016/s0278-4343\(01\)00042-5](https://doi.org/10.1016/s0278-4343(01)00042-5)

Herman, P. M. J., Middelburg, J. J., Koppel, J. V. D., & Heip, C. H. R. (1999). Ecology of estuarine macrobenthos. *Advances in Ecological Research*, 29, 195–240.

Herman, P. M., Middelburg, J. J., & Heip, C. H. (2001). Benthic community structure and sediment processes on an intertidal flat: results from the ECOFLAT project. *Continental Shelf Research*, 21(18-19), 2055–2071.

Hewitt, J. E., Thrush, S. F., & Dayton, P. D. (2008). Habitat variation, species diversity and ecological functioning in a marine system. *Journal of Experimental Marine Biology and Ecology*, 366(1-2), 116–122. [doi: <https://doi.org/10.1016/j.jembe.2008.07.016>]

Hobbs, C. H., Landry, C. B., & Perry, J. E. (2008). Assessing Anthropogenic and Natural Impacts on Ghost Crabs (*Ocypode quadrata*) at Cape Hatteras National Seashore, North Carolina. *Journal of Coastal Research*, 24(6 (246)), 1450–1458. <https://doi.org/10.2112/07-0856.1>

Hornell, J. (1916). Report to the Government of Baroda on the marine zoology of Okhamandal in Kattiawar, Williams and Norgate.

Hughes, D. A. (1966). Behavioural and ecological investigations of the crab *Ocypode ceratophthalmus* (Crustacea: Ocypodidae). *Journal of Zoology*, 150(1), 129–143. <https://doi.org/10.1111/j.1469-7998.1966.tb03000.x>

- Hughes, D. A. (1973). On Mating and the "Copulation Burrows" of Crabs of the Genus *Ocypode* (Decapoda, Brachyura)1). *Crustaceana*, 24(1), 72–76. <https://doi.org/10.1163/156854073x00074>
- Huhta, V. (2007). The role of soil fauna in ecosystems: A historical review. *Pedobiologia-International Journal of Soil Biology*, 50(6), 489–495. <https://doi.org/10.1016/j.pedobi.2006.08.006>
- Iribarne, O., Bortolus, A., & Botto, F. (1997). Between-habitat differences in burrow characteristics and trophic modes in the southwestern Atlantic burrowing crab *Chasmagnathus granulata*. *Marine Ecology Progress Series*, 155, 137–145. <https://doi.org/10.3354/meps155137>
- Jatav, S. K., Kardani, H., Tandel, L., Kumar, A., Mehta, P., & Yadav, S. R. (2023). Crustaceans Diversity along Mangroves of Sikka Coast, Gulf of Kachchh, Gujarat, India. *Indian Journal of Ecology*, 50(5), 1553–1557. <https://doi.org/10.55362/IJE/2023/4095>
- Johannesson, L. T., Stevens, R. L., & Alexanderson, J. H. (2000). Sediment Character in a Micro-Tidal, Harbor-Estuary Environment, Göteborg, Sweden. *Estuaries*, 23(3), 400–410. <https://doi.org/10.2307/1353332>
- Johnson, D. S., & Skutch, A. F. (1928). Littoral Vegetation on a Headland of Mt. Desert Island, Maine. II. Tide-Pools and the Environment and Classification of Submersible Plant Communities. *Ecology*, 9(3), 307–338. <https://doi.org/10.2307/1932371>
- Johnson, D. S., & York, H. H. (1915). *The relation of plants to tide-levels; a study of factors affecting the distribution of marine plants* (Vol. 206, p. 162). Washington, D.C., Carnegie Institution of Washington.
- Jones, D. A. (1972). Aspects of the ecology and behaviour of *Ocypode ceratophthalmus* (Pallas) and *O. kuhlii* de Haan (Crustacea: Ocypodidae). *Journal of Experimental Marine Biology and Ecology*, 8(1), 31–43.

- Jones, M. B. (1976). Limiting factors in the distribution of intertidal crabs (Crustacea: Decapoda) in the avon-heathcote estuary, christchurch. *New Zealand journal of marine and freshwater research*, 10(4), 577-587.
- Jones, M. B., & Simons, M. J. (1982). Habitat preferences of two estuarine burrowing crabs *Helice crassa* Dana (Grapsidae) and *Macrophthalmus hirtipes* (Jacquinot) (Ocypodidae). *Journal of experimental marine biology and ecology*, 56(1), 49-62.
- Katrak, G., Dittmann, S., & Seuront, L. (2008). Spatial variation in burrow morphology of the mud shore crab *Helograpsus haswellianus* (Brachyura, Grapsidae) in South Australian saltmarshes. *Marine and Freshwater Research*, 59(10), 902–911. <https://doi.org/10.1071/mf08044>
- Kemp, S. (1919a). Notes on Crustacea Decapoda in the Indian Museum. 12. Scopimerinae. *Records of Indian Museum*, 16 (5), 305–348. [doi: <https://doi.org/10.5962/bhl.part.25926>]
- Kemp, S. (1919b). Notes on Crustacea Decapoda in the Indian Museum. 12. The Indian species of *Macrophthalmus*. *Records of Indian Museum*, 16 (5), 383–394. [doi: <https://doi.org/10.5962/bhl.part.25928>]
- Khan, R. A. (2003). Biodiversity of marobenthos on the intertidal flats of Sunderban Estuarine region, India. *Records of Zoological Survey of India*, 101 (3–4), 181–205.
- Khan, S. A., Raffi, S. M., & Lyla, P. S. (2005). Brachyuran crab diversity in natural (Pitchavaram) and artificially developed mangroves (Vellar estuary). *Current Science*, 88(8), 1316–1324.
- Kinoshita, K. (2002). Burrow Structure of the Mud Shrimp *Upogebia major* (Decapoda: Thalassinidea: Upogebiidae). *Journal of Crustacean Biology*, 22(2), 474–480. <https://doi.org/10.1163/20021975-99990255>
- Knox, G. A. (2001). *The Ecology of Seashores* (p. 571). CRC Press.

- Krishnamoorthy, P. (2007). Fauna of Chennai coast. In: *Director ZSI, Kolkata (Ed.), Estuarine Ecosystem Series. Vol. 1. Zoological Survey of India, Kolkata*, 83–109.
- Krishnamoorthy, P. (2009). Brachyuran crabs from the collections of Marine Biological Centre. *Records of Zoological Survey of India, Occasional Paper*, 304, 1–46.
- Kristensen, E. (2008). Mangrove crabs as ecosystem engineers; with emphasis on sediment processes. *Journal of Sea Research*, 59(1-2), 30–43.
<https://doi.org/10.1016/j.seares.2007.05.004>
- Kristensen, E., & Alongi, D. M. (2006). Control by fiddler crabs (*Uca vocans*) and plant roots (*Avicennia marina*) on carbon, iron, and sulfur biogeochemistry in mangrove sediment. *Limnology and Oceanography*, 51(4), 1557–1571.
<https://doi.org/10.4319/lo.2006.51.4.1557>
- Kumar, A., Sushilkumar, M. & Galil, B.S. (2013) Calappid and Leucosiid crabs (Crustacea: Decapoda: Brachyura) from Kerala, India, with the description of a new species of *Mursia Desmarest*, 1823, from the Arabian Sea and redescription of *M. bicristimana* Alcock & Anderson, 1894. *Zootaxa*, 3746 (4), 529–551. [doi: <https://doi.org/10.11646/zootaxa.3746.4.2>]
- Kumar, V. S., & Kumar, K. A. (2010). Waves and currents in tide-dominated location off Dahej, Gulf of Khambhat, India. *Marine Geodesy*, 33(2-3), 218-231.
- Kumaralingam, S., Raghunathan, C., Ajithkumar, T.T. & Chandra K. (2017). Reef associated common crustaceans of Andaman and Nicobar Islands, Lakshadweep, and Gulf of Mannar. *Records of Zoological Survey of India, Occasional Paper*, 380, 1–249.
- Lalli, C. M., & Parsons, T. R. (2012). *Biological oceanography: an introduction* (p. 314). Butterworth-Heinemann, Massachusetts, United States.
- Lalli, C., & Parsons, T. R. (1997). *Biological oceanography: an introduction*. Elsevier.

- Laurent Chauvaud, Jean, F., Olivier Ragueneau, & Gérard Thouzeau. (2000). Long-term variation of the Bay of Brest ecosystem: benthic-pelagic coupling revisited. *Marine Ecology Progress Series*, 200, 35–48. <https://doi.org/10.3354/meps200035>
- Lavajoo, F., Kamrani, E., & Sajjadi, M. (2011). Distribution, population and reproductive biology of the fiddler crab *Uca sindensis* (Crustacea: Ocypodidae) in a subtropical mangrove of Pohl Area. *Journal of the Persian Gulf*, 2(5), 9-16.
- Lee, S. Y. (1998). Ecological role of grapsid crabs in mangrove ecosystems: a review. *Marine and Freshwater Research*, 49(4), 335–343. <https://doi.org/10.1071/mf97179>
- Lee, S. Y. (2008). Mangrove macrobenthos: Assemblages, services, and linkages. *Journal of Sea Research*, 59(1-2), 16–29. <https://doi.org/10.1016/j.seares.2007.05.002>
- Lee, S. Y. (2015). Ecology of brachyura. In P. Castro, P. Davie, D. Guinot, F. Schram, & C. V. V. Klein (Eds.), *Treatise on Zoology - Anatomy, Taxonomy, Biology: The Crustacea Volume 9 Part C (2 vols)*. Brill, Netherlands.
- Lee, S.-L., & Shirley. (2004). Do diameters of burrows and food pellets provide estimates of the size structure of a population of *Dotilla myctiroides* at the sand-flats of Ao Tung Khen, Phuket. *Bulletin of the National Research Centre*, 65.
- Lee, Y. H., & Koh, C. H. (1994). Biogenic sedimentary structures on a Korean mud flat: Spring-neap variations. *Netherlands Journal of Sea Research*, 32(1), 81–90. [https://doi.org/10.1016/0077-7579\(94\)90030-2](https://doi.org/10.1016/0077-7579(94)90030-2)
- Lessard-Pilon, S., Porter, M. D., Cordes, E. E., MacDonald, I., & Fisher, C. R. (2010). Community composition and temporal change at deep Gulf of Mexico cold seeps. *Deep Sea Research Part II: Topical Studies in Oceanography*, 57(21-23), 1891–1903. <https://doi.org/10.1016/j.dsr2.2010.05.012>

- Levinton, J. S. (2001). *Marine Biology: Function, Biodiversity, Ecology* (p. 530). Oxford University Press, Oxford, United Kingdom.
- Lewis, J. R. (1964). The ecology of rocky shores. *English Universities Press Ltd., London, United Kingdom*, 323.
- Lim, S. S. L. (2006). Fiddler crab burrow morphology: How do burrow dimensions and bioturbative activities compare in sympatric populations of *Uca vocans* (Linnaeus, 1758) and *U. annulipes* (H. Milne Edwards, 1837)? *Crustaceana*, 79(5), 525–540. <https://doi.org/10.1163/156854006777584241>
- Lim, S. S. L., & Diong, C. H. (2003). Burrow-Morphological Characters of the Fiddler Crab, *Uca annulipes* (H. Milne Edwards, 1837) and Ecological Correlates in a Lagoonal Beach on Pulau Hantu, Singapore. *Crustaceana*, 76(9), 1055–1069. <https://www.jstor.org/stable/20105648>
- Lima, S. L., & Dill, L. M. (1990). Behavioral decisions made under the risk of predation: a review and prospectus. *Canadian Journal of Zoology*, 68(4), 619–640. <https://doi.org/10.1139/z90-092>
- Linsenmair, L. H. (1967). Konstruktion und Signalfunktion der Sandpyramide der Reiterkrabbe *Ocypode saratan* Forsk. (Decapoda Brachyura Ocypodidae)1)2). *Zeitschrift Für Tierpsychologie*, 24(4), 403–456. <https://doi.org/10.1111/j.1439-0310.1967.tb01238.x>
- Little, C. (2000). *The biology of soft shores and estuaries* (1st ed., p. 252). Oxford University Press, New York.
- Little, C., & Smith, L. P. (1980). Vertical zonation on rocky shores in the Severn Estuary. *Estuarine and Coastal Marine Science*, 11(6), 651–669.
- Litulo, C. (2005). Population structure and reproductive biology of the fiddler crab *Uca inversa* (Hoffman, 1874) (Brachyura: Ocypodidae). *Acta Oecologica*, 27(3), 135–141.
- Lohrer, A. M., Fukui, Y., Wada, K., & Whitlatch, R. B. (2000). Structural complexity and vertical zonation of intertidal crabs, with focus on habitat

- requirements of the invasive Asian shore crab, *Hemigrapsus sanguineus* (de Haan). *Journal of Experimental Marine Biology and Ecology*, 244(2), 203-217.
- Lomovasky, B. J., Casariego, A. M., Brey, T., & Iribarne, O. (2006). The effect of the SW Atlantic burrowing crab *Chasmagnathus granulatus* on the intertidal razor clam *Tagelus plebeius*. *Journal of Experimental Marine Biology and Ecology*, 337(1), 19–29. <https://doi.org/10.1016/j.jembe.2006.05.016>
- Lotze, H. K., Lenihan, H. S., Bourque, B. J., Bradbury, R. H., Cooke, R. G., Kay, M. C., Kidwell, S. M., Kirby, M. X., Peterson, C. H., & Jackson, J. B. C. (2006). Depletion, Degradation, and Recovery Potential of Estuaries and Coastal Seas. *Science*, 312, 1806–1809. <https://doi.org/10.1126/science.1128035>
- Lucrezi, S., Schlacher, T. A., & Walker, S. (2008). Monitoring human impacts on sandy shore ecosystems: a test of ghost crabs (*Ocypode* spp.) as biological indicators on an urban beach. *Environmental Monitoring and Assessment*, 152(1-4), 413–424. <https://doi.org/10.1007/s10661-008-0326-2>
- Lukman Adi Nugroho, Agatha Sih Piranti, & Moh. Husein Sastranegara. (2020). Plankton Community and Water Quality During Maximum Tidal Range in Segara Anakan Cilacap. *IOP Conference Series*, 593(1), 012020–012020. <https://doi.org/10.1088/1755-1315/593/1/012020>
- Macintosh, D. J. (1988). The ecology and physiology of decapods of mangrove swamps. *Aspects of Decapod Crustacean biology: the proceedings of a symposium held at the Zoological Society of London on 8th and 9th April 1987 / edited by A.A. Fincham and P.S. Rainbow*. Clarendon Press; New York: Oxford University Press
- Macnae, W., & Kalk, M. (1962). The Fauna and Flora of Sand Flats at Inhaca Island, Mocambique. *The Journal of Animal Ecology*, 31(1), 93–128. <https://doi.org/10.2307/2334>
- Maheta, N.P., & Vachhrajani, K. D. (2023). Burrow characteristics of the fiddler crab – *Austruca sindensis* (Alcock, 1900) from mudflats of Gulf of Khambhat,

- Gujarat, India. *Arthropods*, 12(1), 37–56.
[http://www.iaees.org/publications/journals/arthropods/articles/2023-12\(1\)/burrow-characteristics-of-fiddler-crab.pdf](http://www.iaees.org/publications/journals/arthropods/articles/2023-12(1)/burrow-characteristics-of-fiddler-crab.pdf)
- Mandal, A.K. & Nandi, N.C. (1989). Fauna of Sundarban Mangrove Ecosystems, West Bengal, India. In: *Director ZSI, Kolkata (Ed.), Fauna of Conservation Areas. Vol. 3. Zoological Survey India, Kolkata*, 1–116.
- Mandal, A.K. & Nandi, N.C. (1989). Fauna of Sundarban Mangrove Ecosystems, West Bengal, India. In: *Director ZSI, Kolkata (Ed.), Fauna of Conservation Areas. Vol. 3. Zoological Survey India, Kolkata*, pp. 1-116.
- Mantelatto, F. L. M., & Fransozo, A. (1999). Characterization of the physical and chemical parameters of Ubatuba Bay, northern coast of São Paulo State, Brazil. *Revista Brasileira de Biologia*, 59, 23-31.
- Mantelatto, F. L. M., & Fransozo, A. (2000). Brachyuran community in Ubatuba bay, northern coast of São Paulo state, Brazil. *Journal of shellfish research*, 19(2), 701-710.
- Mantelatto, F. L. M., Fransozo, A., & Negreiros-Fransozo, M. L. (1995). Distribuição do caranguejo *Hepatus pudibundus* (Herbst, 1785) (Crustácea, Decapoda, Brachyura) na Enseada da Fortaleza, Ubatuba (SP), Brasil Distribution of calico crab *Hepatus Pudibunds* (Herbst, 1785) (Crustacea, Decapoda, Brachyura) in Fortaleza Bay, Ubatuba (SP), Brazil.
- Marcus, N. H., & Boero (1998). Minireview: The importance of benthic-pelagic coupling and the forgotten role of life cycles in coastal aquatic systems. *Limnology and Oceanography*, 43(5), 763–768.
<https://doi.org/10.4319/lo.1998.43.5.0763>
- Martin, J. W., & Davis, G. E. (2001). *An Updated Classification of The Recent Crustacea* (Science Series Natural History Museum of Los Angeles County, Vol. 39). Natural History Museum of Los Angeles County Los Angeles, California, United States of America.

- Matsumasa, M., Takeda, S., Poovachiranon, S., & Poovachiranon, M. (1992). Distribution and Shape of *Dotilla myctiroides* (Brachyura: Ocypodidae) Burrow in the Seagrass *Enhalus acoroides* Zone. *Benthos Research*, 43, 1–9. https://doi.org/10.5179/benthos1990.1992.43_1
- McAllister, D. (1995). Status of the World Ocean and Its Biodiversity. *Sea Wind*, 9(4), 1–14.
- McLachlan, A., & Brown, A. C. (2006). *The Ecology of Sandy Shores* (1st ed.). Elsevier, Amsterdam, 328 pp.
- Menendez, R. J. (1987). Vertical zonation of the xanthid mud crabs *Panopeus obesus* and *Panopeus simpsoni* on oyster reefs. *Bulletin of Marine Science*, 40(1), 73–77.
- Michaels, R. E., & Zieman, J. C. (2013). Fiddler crab (*Uca* spp.) burrows have little effect on surrounding sediment oxygen concentrations. *Journal of Experimental Marine Biology and Ecology*, 448, 104–113. <https://doi.org/10.1016/j.jembe.2013.06.020>
- Miller, L. P., Harley, C. D. G., & Denny, M. W. (2009). The role of temperature and desiccation stress in limiting the local-scale distribution of the owl limpet, *Lottia gigantea*. *Functional Ecology*, 23(4), 756–767. <https://doi.org/10.1111/j.1365-2435.2009.01567.x>
- Milne-Edwards, H. (1834). *Histoire naturelle des Crustacés comprenant l'anatomie, la physiologie et la classification de ces animaux*. (Vols. 1–3, p. 531). Librairie encyclopédique de Roret, Paris.
- Mitra, A., & Zaman, S. (2016). *Basics of Marine and Estuarine Ecology* (1st ed.). Springer New Delhi, India. <https://doi.org/10.1007/978-81-322-2707-6>
- Mitra, S. & Misra, A. (2006). A report on intertidal macrofauna of Talsari (Balasore: Orissa). *Records of Zoological Survey of India*, 106 (3), 131–141.

- Mitra, S., Misra, A. & Pattanayak, J.G. (2010). Intertidal Macrofauna of Subarnarekha Estuary (Balasore: Orissa). *Records of Zoological Survey of India, Occasional Paper*, 313, 1-135.
- Mokhlesi, A., Kamrani, E., Backwell, P., & Sajjadi, M. (2011). Study on the behaviour of two fiddler crabs, *Uca sindensis* and *Uca annulipes* (Decapoda: Ocypodidae), in Bandar Abbas, Iran. *Journal of the Marine Biological Association of the United Kingdom*, 91(1), 245-249. [doi: <https://doi.org/10.1017/S0025315410000172>]
- Mokhtari, M., Ghafar, M. A., Usup, G., & Cob, Z.C. (2015) Determination of key environmental factors responsible for distribution patterns of fiddler crabs in a tropical mangrove ecosystem. *PLOS One*, 10(1). [doi: <https://doi.org/10.1371/journal.pone.0117467>]
- Morrissey, D. J., DeWitt, T. H., Roper, D. S., & Williamson, R. B. (1998). Variation in the depth and morphology of burrows of the mud crab *Helice crassa* among different types of intertidal sediment in New Zealand. *Marine Ecology Progress Series*, 182, 231-242. [doi: <https://doi.org/10.3354/meps182231>]
- Moser, S., Macintosh, D., Laoprasert, S., & Tongdee, N. (2005). Population ecology of the mud crab *Scylla olivacea*: a study in the Ranong mangrove ecosystem, Thailand, with emphasis on juvenile recruitment and mortality. *Fisheries Research*, 71(1), 27–41. <https://doi.org/10.1016/j.fishres.2004.07.008>
- Moulton, C. E., Brady, R. S., & Belthoff, J. R. (2006). Association between Wildlife and Agriculture: Underlying Mechanisms and Implications in Burrowing Owls. *Journal of Wildlife Management*, 70(3), 708–716. [https://doi.org/10.2193/0022-541x\(2006\)70\[708:abwaau\]2.0.co;2](https://doi.org/10.2193/0022-541x(2006)70[708:abwaau]2.0.co;2)
- Mouton, E. C., & Felder, D. L. (1996). Burrow distributions and population estimates for the fiddler crabs *Uca spinicarpa* and *Uca longisignalis* in a Gulf of Mexico salt marsh. *Estuaries*, 19, 51-61.

- Murray, N. J., Phinn, S. R., DeWitt, M., Ferrari, R., Johnston, R., Lyons, M. B., Clinton, N., Thau, D., & Fuller, R. A. (2019). The global distribution and trajectory of tidal flats. *Nature*, 565(7738), 222–225. <https://doi.org/10.1038/s41586-018-0805-8>
- Naderloo, R. (2017). *Atlas of crabs of the Persian Gulf*. Springer, 440.
- Naderloo, R., & Tuerkay, M. (2012). Decapod crustaceans of the littoral and shallow sublittoral Iranian coast of the Persian Gulf: faunistics, biodiversity and zoogeography. *Zootaxa*, 3374(1), 1-67. [doi: <https://doi.org/10.11646/zootaxa.3374.1.1>]
- Naderloo, R., Tuerkay, M., & Apel, M. (2011). Brachyuran crabs of the family Macrophthalmidae Dana, 1851 (Decapoda: Brachyura: Macrophthalmidae) of the Persian Gulf. *Zootaxa*, 2911(1), 1-42. [doi: <https://doi.org/10.11646/zootaxa.2911.1.1>]
- Naeem, S., Thompson, L. J., Lawler, S. P., Lawton, J. H., & Woodfin, R. M. (1994). Declining biodiversity can alter the performance of ecosystems. *Nature*, 368, 734-737.
- Nandi, N. C., & Dev Roy, M. K. (1991). Burrowing activity and distribution of *Scylla serrata* (Forsk.) from Hooghly and Matla estuaries, Sundarban, West Bengal. *Journal of the Bombay Natural History Society*, 88(2), 167–171.
- Naskar, R. (2018). Abundance and distribution pattern of intertidal shore crabs on exposed sandy beaches of Bakhali, Sundarbans. *International Journal of Engineering, Science and Mathematics*, 4(2), 153-159.
- Nayak, S. R., & Sahai, B. (1985). Coastal morphology: a case study of the Gulf of Khambhat (Cambay). *International Journal of Remote Sensing*, 6(3-4), 559–567. <https://doi.org/10.1080/01431168508948478>
- Naylor, E. (1982). Tidal and lunar rhythms in animals and plants. *Biological Timekeeping*, 14, 33–48.

- Naylor, E. (2001). Marine Animal Behaviour In Relation to Lunar Phase. *Earth, Moon, and Planets*, 85/86, 291–302. <https://doi.org/10.1023/a:1017088504226>
- Naylor, E., & Williams, B. G. (1984). Environmental entrainment of tidally rhythmic behaviour in marine animals. *Zoological Journal of the Linnean Society*, 80(2-3), 201–208. <https://doi.org/10.1111/j.1096-3642.1984.tb01973.x>
- Needham, D. M., Korupolu, R., Zanni, J. M., Pradhan, P., Colantuoni, E., Palmer, J. B., Brower, R. G., & Fan, E. (2010). Early physical medicine and rehabilitation for patients with acute respiratory failure: a quality improvement project. *Archives of Physical Medicine and Rehabilitation*, 91(4), 536–542. <https://doi.org/10.1016/j.apmr.2010.01.002>
- Newell, R. C. (1979). Biology of Intertidal Animals. *Marine Ecological Surveys Ltd., Faversham, Kent, United Kingdom*, 781.
- Ng, P. K. L., Guinot, D., & Davie, P. J. F. (2008). Systema Brachyurorum: Part I. Anannotated checklist of extant brachyuran crabs of the world. *The Raffles Bulletin of Zoology*, 17, 1–286.
- Ng, P.K.L. & Kumar, A. B. (2015a). A new species of *Afropinnotheres* Manning, 1993 (Crustacea, Brachyura, Pinnotheridae) from southwestern India, the first record of the genus from the Indian Ocean, with a review of the Pinnotheridae of India and adjacent seas. *Zootaxa*, 3947 (2), 264–274. [doi: <https://doi.org/10.11646/zootaxa.3947.2.8>]
- Ng, P.K.L. & Kumar, A. B. (2015b). The species of *Moloha* Barnard, 1946, from the western Indian Ocean, with the description of a new species from India (Crustacea: Brachyura: Homolidae). *European Journal of Taxonomy*, 166, 1–25. [doi: <https://doi.org/10.5852/ejt.2015.166>]
- Ng, P.K.L. & Kumar, A. B. (2016). *Carcinoplax faciata*, a new species of deep-water goneplacid crab from southwestern India (Crustacea: Decapoda: Brachyura: Goneplacoidea). *Zootaxa*, 4147 (2), 192–200. [doi: <https://doi.org/10.11646/zootaxa.4147.2.6>]

- Ng, P.K.L., Prema, M. & Ravichandran, S. (2018). A new species of deep-water spider crab of the genus *Paramaya* De Haan, 1837 from the Bay of Bengal, India (Crustacea, Brachyura, Majidae). *Zookeys*, 769, 77–88. [doi: <https://doi.org/10.3897/zookeys.769.26152>]
- Ng, P.K.L., Trivedi, J.N. & Vachhrajani, K.D. (2015). *Dentoxanthus iranicus* Stephensen, 1946 (Crustacea: Decapoda: Brachyura: Galenidae): a new record from Gujarat, India, with systematic notes. *Marine Biodiversity Records*, 8, 1–5. [doi: <https://doi.org/10.1017/S1755267215001098>]
- Nishihira, M. (1984). Ecological notes on the rock dwelling ocypodid crab, *Ilyoplax integra* (Tesch), newly recorded from Okinawa, Japan. *Galaxea*, 3, 97–101.
- Nordhaus, I., Diele, K., & Wolff, M. (2009). Activity patterns, feeding and burrowing behaviour of the crab *Ucides cordatus* (Ucididae) in a high intertidal mangrove forest in North Brazil. *Journal of Experimental Marine Biology and Ecology*, 374(2), 104–112. <https://doi.org/10.1016/j.jembe.2009.04.002>
- Nugroho, L. A., Piranti, A. S., & Sastranegara, M. H. (2020). Plankton community and water quality during maximum tidal range in Segara Anakan Cilacap. In *IOP Conference Series: Earth and Environmental Science*, 593(1), 1–14.
- Oliveira, R. F., McGregor, P. K., Burford, F. R. L., Custódio, M. R., & Latruffe, C. (1998). Functions of mudballing behaviour in the European fiddler crab *Uca tangeri*. *Animal Behaviour*, 55(5), 1299–1309. <https://doi.org/10.1006/anbe.1997.0685>
- Padate, V. P., Rivonker, C. U. & Anil, A. C. (2013). A new record of *Scylla olivacea* (Decapoda, Brachyura, Portunidae) from Goa, central west coast of India: A comparative diagnosis. *Indian Journal of Geo-Marine Sciences*, 42 (1), 82–89.
- Padate, V. P., Rivonker, C. U., Anil, A. C., Sawant, S. S. & Krishnamurthy, V. (2010) A new species of portunid crab of the genus *Charybdis* (De Haan, 1833) (Crustacea: Decapoda: Brachyura) from Goa, India. *Marine Biology*

Research, 6 (6), 579–590. [doi: <https://doi.org/10.1080/17451000903505608>]

Pal, S. R. & Khora, S. S. (1999). Studies on the carcinological fauna of Gopalpur coast, Bay of Bengal. 1. Non-Brachyrrhynchan (Decapoda: Brachyura). *Records of Zoological Survey of India*, 97 (4), 5–22.

Palmer, J. D. (1995). *The biological rhythms and clocks of intertidal animals*. <https://api.semanticscholar.org/CorpusID:82706949>

Palumbi, S., Evans, T., Pespeni, M., & Somero, G. (2019). Present and Future Adaptation of Marine Species Assemblages: DNA-Based Insights into Climate Change from Studies of Physiology, Genomics, and Evolution. *Oceanography*, 32(3), 82–93. <https://doi.org/10.5670/oceanog.2019.314>

Pandya P. J., 2011. Benthic community structure of Mahi River estuary with special reference to animal-sediment relationship. Ph. D. Thesis. The Maharaja Sayajirao University of Baroda, Vadodara, Gujarat, India.

Pandya, P. J., & Vachhrajani, K. D. (2013). Brachyuran crab diversity of lower estuarine mud flats of Mahi River with new record of two species from Gujarat, India. *Arthropods*, 2(4), 242-250.

Pandya, P., & Vachhrajani, K. (2010). Spatial Distribution and Substratum Preference of the Brachyuran Crab, *Macrophthalmus depressus* (Decapoda, Ocypodidae) along the Lower Estuarine Mudflat of Mahi River, Gujarat, India. *Crustaceana*, 83(9), 1055–1067. <https://doi.org/10.1163/001121610x521235>

Parisi, B. (1916). I Decapodi Giapponesi del Museo di Milano. IV. Cyclometopa. *Atti Societas Italiano Sciences naturelle*, 55, 153-190. [in Italian]

Pati, S. K., Sahu, K. C., Swain, D., Baliarsingh, S. K. & Sharma, R. M. (2012). Marine crabs (Decapoda: Anomura and Brachyura). In: *Director ZSI, Kolkata (Ed.), Fauna of Maharashtra. State Fauna Series. Vol. 20. Part 2. Zoological Survey of India, Kolkata*, 381–385.

- Patra, B. C., Bhattacharya, M., Kar, A., Das, B. K., Ghosh, S., Parua, S., Patra, S. & Rakshit, S. (2017). Crab diversity of Digha coast, West Bengal, India. *Proceedings of the Zoological Society*. [doi: <https://doi.org/10.1007/s12595-017-0251-x>]
- Paul, J., Mondal, S., Mondal, R., & Sarkar, D. (2019). Burrow morphology of the ocypodid crab *Ocypode ceratophthalma* at Chandipur Coast, Eastern India and its implications. *Current Science*, 117(4), 699–705.
- Pawar, P. R. (2017). Biodiversity of brachyuran crabs (Crustacea: Decapoda) from Uran, Navi Mumbai, west coast of India. *Advances in Environmental Biology*, 11(2), 103–112.
- Pearse, A. S. (1912). The habits of fiddler-crabs. *Philippine Journal of Science-Section D: General Biology, Ethnology and Anthropology*, 7(3), 113–133.
- Penha-Lopes, G., Bartolini, F., Limbu, S., Cannicci, S., Kristensen, E., & Paula, J. (2009). Are fiddler crabs potentially useful ecosystem engineers in mangrove wastewater wetlands? *Marine Pollution Bulletin*, 58(11), 1694–1703. <https://doi.org/10.1016/j.marpolbul.2009.06.015>
- Pombo, M., Oliveira, A. L., Xavier, L. Y., Siegle, E., & Turra, A. (2017). Natural drivers of distribution of ghost crabs *Ocypode quadrata* and the implications of estimates from burrows. *Marine Ecology Progress Series*, 565, 131–147. <https://doi.org/10.3354/meps11991>
- Poore, G. C. (2004). *Marine Decapod Crustacea of Southern Australia* (p. 616). CSIRO PUBLISHING, Clayton, Australia.
- Pörtner, H. (2001). Climate change and temperature-dependent biogeography: oxygen limitation of thermal tolerance in animals. *Naturwissenschaften*, 88, 137–146. [doi: <https://doi.org/10.1007/s001140100216>]
- Powers, L. W., & Cole, J. F. (1976). Temperature variation in fiddler crab microhabitats. *Journal of Experimental Marine Biology and Ecology*, 21(2), 141–157. [https://doi.org/10.1016/0022-0981\(76\)90035-6](https://doi.org/10.1016/0022-0981(76)90035-6)

- Prasad, P. N., & Neelakantan, B. (1988). Food and Feeding of The Mud Crab *Scylla serrata* Forskal (Decapoda : Portunidae) From Karwar Waters. *Indian Journal of Fisheries*, 35(3), 164–170.
<https://epubs.icar.org.in/index.php/IJF/article/view/10693>
- Pratt, D. R., Lohrer, A. M., Pilditch, C. A., & Thrush, S. F. (2013). Changes in Ecosystem Function Across Sedimentary Gradients in Estuaries. *Ecosystems*, 17(1), 182–194. <https://doi.org/10.1007/s10021-013-9716-6>
- Prema, M., Ravichandran, S., & Sivakumar, V. K. (2021). New record of the swimming crab *Charybdis (Archias) hongkongensis* Shen, 1934 (Brachyura, Portunidae) from India. *Crustaceana*, 94(11-12), 1335–1343.
<https://doi.org/10.1163/15685403-bja10167>
- Priyadarshani, S., Jayamanne, S., & Hirimuthugoda, Y. (2008). Diversity of mangrove crabs in Kadalkele, Negombo eatuary, Sri Lanka. *Sri Lanka Journal of Aquatic Sciences*, 13(0), 109–121.
<https://doi.org/10.4038/sljas.v13i0.2210>
- Quang Minh Dinh, Tra, T., Ngo, Dang Hoang Dong, & Lam Trung Hau. (2014). Burrow Configuration and Utilization of the Blue-Spotted Mudskipper, *Boleophthalmus boddarti* Caught in Soc Trang, Vietnam. *Journal of Fisheries and Environment*, 38(2), 1–9. <https://li01.tci-thaijo.org/index.php/JFE/article/view/80572>
- Quinn, N. J., & Kojis, B. L. (1987). Reproductive biology of *Scylla* spp. (Crustacea: Portunidae) from the labu estuary in Papua New Guinea. *Bulletin of Marine Science*, 41(2), 234–241.
- Qureshi, N. A., & Saher, N. U. (2012). Burrow morphology of three species of fiddler crab (*Uca*) along the coast of Pakistan. *The Belgian Journal of Zoology*, 142(2), 114–126.
- Rafaelli, D. & Hawkins, S. (1996). Intertidal Ecology. *Chapman & Hall, London*, 356.

- Raffaelli, D., Bell, E. M., Weithoff, G., Matsumoto, A., Cruz-Motta, J. J., Kershaw, P., Parker, R., Parry, D., & Jones, M. K. (2003). The ups and downs of benthic ecology: considerations of scale, heterogeneity and surveillance for benthic–pelagic coupling. *Journal of Experimental Marine Biology and Ecology*, 285-286, 191–203. [https://doi.org/10.1016/s0022-0981\(02\)00527-0](https://doi.org/10.1016/s0022-0981(02)00527-0)
- Raghunathan, C. (2015). Littoral faunal diversity of Great Nicobar island. *Records of Zoological Survey of India, Occasional Paper*, 369, 1–172.
- Rama Rao, K.V., Nageswara Rao, C.A., Nahar, S.C., Rao, D.V. & Mohapatra, A. (1992). Studies on the ecology and fauna of the Rushikulya Estuary, (Ganjam), Orissa. In: Director ZSI, Kolkata (Ed.), *Estuarine Ecosystem Series*. Vol. 1. Zoological Survey of India, Kolkata, pp. 7-26.
- Ramakrishna, S. J. & Talukdar, S. (2003). Invertebrates of Digha coast and some recommendations on their conservation. *Records of Zoological Survey of India*, 101(3-4), 1–23.
- Ramana Murthy, K.V. & Kondala Rao, B. (1993). UNESCO Curriculum Workshop on Management of Mangrove Ecosystem and Coastal Protection. Andhra University, Visakhapatnam, pp. 6-12.
- Rao, D. V. & Rath, S. (2014). Fauna of Brahmani-Baitarani Estuarine complex, Odisha, Bay of Bengal, w.s.r. to ichthyofauna and crustaceans. In: Director ZSI, Kolkata (Ed.), *Estuarine Ecosystem Series*. Vol. 7. Zoological Survey of India, Kolkata, 1–120.
- Rao, D.V., & Rath, S. (2013). Ichthyofauna and decapod crustacean fauna of the Nuanai Estuary, Odisha. In: Director ZSI, Kolkata (Ed.), *Estuarine Ecosystem Series*. Vol. 8. Zoological Survey of India, Kolkata, pp. 1-64.
- Rao, G. S., Suseelan, C. & Kathirvel, M. (1989). Crustacean resources of the Lakshadweep Islands. In: Suseelan, C. (Ed.), *Marine living resources of the union territory of Lakshadweep Islands: An Indicative Survey with*

Suggestions for Development. Central Marine Fisheries Research Institute, Cochin, 72–76.

- Rath, S. & Dev Roy, M. K. (2009). Brachyuran crabs (Crustacea: Decapoda: Brachyura). In: Director ZSI, Kolkata (Ed.), *Estuarine Ecosystem Series. Vol. 5. Fauna of Krishna Estuary. Zoological Survey of India, Kolkata*, 43–81.
- Rath, S. & Dev Roy, M. K. (2010). Brachyuran crabs (Crustacea: Decapoda: Brachyura). In: Director ZSI, Kolkata (Ed.), *Estuarine Ecosystem Series. Vol. 6. Vamsadhara and Nagavali Estuary, Srikakulam, Andhra Pradesh. Zoological Survey of India, Kolkata*, 23–45.
- Rath, S., & Nandi, N.C. (2017). A note on Portunid crabs of Odisha Coast, India. *Oceanography & Fisheries*, 1(3), 1-3.
- Rath, S., & Dev Roy, M. K. (2011). Crabs and Prawns (Crustacea: Decapoda) of Bahuda Estuary, Ganjam, Orissa. *Records of the Zoological Survey of India*, 111(1), 47–61. <https://doi.org/10.26515/rzsi/v111/i1/2011/158891>
- Raval, J. V., Chauhan, S. K., Kardani, H. K., & Vadher, P. J. (2020). Diversity of marine brachyuran crabs from the Sikka coast, Gujarat, India. *Journal of Experimental Zoology, India*, 23(2), 1463–1471.
- Ravichandran, S. & Kannupandi, T. (2007). Biodiversity of crabs in Pichavaram mangrove environment. In: Director ZSI, Kolkata (Ed.), *National Symposium on Conservation and Validation of Marine Biodiversity. Zoological Survey of India, Kolkata*, pp. 331-340.
- Ravichandran, S., Soundarapandian, P., & Kannupandi, T. (2001). Zonation and distribution of crabs in Pichavaram mangrove swamp, southeast coast of India. *Indian Journal of Fisheries*, 48(2), 221-226.
- Reaka-Kudla, M. L. (1997). The global biodiversity of coral reefs: A comparison with rainforests. In M. L. Reaka-Kudla, D. E. Wilson, & E. O. Wilson (Eds.), *Biodiversity II: Understanding and Protecting Our Natural Resources* (pp. 83–108). Joseph Henry / National Academy Press, Washington, D. C.

- Rebach, S. (1974). Burying Behavior in Relation to Substrate and Temperature in the Hermit Crab, *Pagurus Longicarpus*. *Ecology*, 55(1), 195–198.
<https://doi.org/10.2307/1934636>
- Reinsel, K. A., & Rittschof, D. (1995). Environmental regulation of foraging in the sand fiddler crab *Uca pugilator* (Bosc 1802). *Journal of Experimental Marine Biology and Ecology*, 187(2), 269-287.
- Reiss, H., Birchenough, S., Borja, A., Buhl-Mortensen, L., Craeymeersch, J., Dannheim, J., Darr, A., Galparsoro, I., Gogina, M., Neumann, H. and Populus, J. (2015). Benthos distribution modelling and its relevance for marine ecosystem management. *ICES Journal of Marine Science*, 72(2), 297-315.
- Resgalla Jr., C., Brasil, E. de S., & Salomão, L. C. (2007). The effect of temperature and salinity on the physiological rates of the mussel *Perna perna* (Linnaeus 1758). *Brazilian Archives of Biology and Technology*, 50(3), 543–556.
<https://doi.org/10.1590/s1516-89132007000300019>
- Ribeiro, P. D., Daleo, P., & Iribarne, O. O. (2010). Density affects mating mode and large male mating advantage in a fiddler crab. *Oecologia*, 164(4), 931–941.
<https://doi.org/10.1007/s00442-010-1801-3>
- Ribeiro, P. D., Iribarne, O., & Daleo, P. (2005). The relative importance of substratum characteristics and recruitment in determining the spatial distribution of the fiddler crab *Uca uruguayensis* Nobili. *Journal of Experimental Marine Biology and Ecology*, 314(1), 99–111.
<https://doi.org/10.1016/j.jembe.2004.09.014>
- Ringelberg, J. (1999). The photobehaviour of *Daphnia* spp. as a model to explain diel vertical migration in zooplankton. *Biological Reviews of the Cambridge Philosophical Society*, 74(4), 397–423.
<https://doi.org/10.1017/s0006323199005381>
- Ringold, P. (1979). Burrowing, root mat density, and the distribution of fiddler crabs in the eastern United States. *Journal of Experimental Marine Biology*

and *Ecology*, 36(1), 11–21. [https://doi.org/10.1016/0022-0981\(79\)90097-2](https://doi.org/10.1016/0022-0981(79)90097-2)

Robertson, J. R., Bancroft, K., Vermeer, G. K., & Plaisier, K. (1980). Experimental studies on the foraging behavior of the sand fiddler crab *Uca pugilator* (Bosc, 1802). *Journal of Experimental Marine Biology and Ecology*, 44(1), 67–83.

Robertson, W. D., & Kruger, A. (1994). Size at maturity, mating and spawning in the portunid crab *Scylla serrata* (Forskål) in Natal, South Africa. *Estuarine, Coastal and Shelf Science*, 39(2), 185–200.

Rodrigues, E., Freitas, R., Delgado, N. de C., & Soares-Gomes, A. (2016). Distribution patterns of the ghost crab *Ocypode cursor* on sandy beaches of a tropical island in the Cabo Verde archipelago, Eastern Central Atlantic. *African Journal of Marine Science*, 38(2), 181–188. <https://doi.org/10.2989/1814232x.2016.1176602>

Rosa, L. C. D., & Borzone, C. A. (2008). Spatial distribution of the *Ocypode quadrata* (Crustacea: Ocypodidae) along estuarine environments in the Paranaguá Bay Complex, southern Brazil. *Revista Brasileira de Zoologia*, 25, 383–388. [doi: <https://doi.org/10.1590/S0101-81752008000300001>]

Ruscoe, I. M., Shelley, C. C., & Williams, G. R. (2004). The combined effects of temperature and salinity on growth and survival of juvenile mud crabs (*Scylla serrata* Forskål). *Aquaculture*, 238(1-4), 239–247. <https://doi.org/10.1016/j.aquaculture.2004.05.030>

Saeedi, H., & Costello, M. J. (2019). A world dataset on the geographic distributions of Solenidae razor clams (Mollusca: Bivalvia). *Biodiversity Data Journal*, (7). [doi:<https://doi.org/10.3897/EBJ.7.e.31375>].

Saeedi, H., Costello, M. J., Warren, D., & Brandt, A. (2019). Latitudinal and bathymetrical species richness patterns in the NW Pacific and adjacent Arctic Ocean. *Scientific Reports*, 9(1), 9303. [doi: <https://doi.org/10.1038/s41598-019-45813-9>]

- Saeedi, H., Dennis, T. E., & Costello, M. J. (2017). Bimodal latitudinal species richness and high endemism of razor clams (Mollusca). *Journal of Biogeography*, 44(3), 592-604. [doi: <https://doi.org/10.1111/jbi.12903>]
- Saeedi, H., Kamrani, E., Nordhaus, I., & Diele, K. (2018). Sediment temperature impact on population structure and dynamics of the crab *Austruca iranica* Pretzmann, 1971 (Crustacea: Ocypodidae) in subtropical mangroves of the Persian Gulf. *Wetlands*, 38(3), 539-549. [doi:<https://doi.org/10.1007/s13157-018-0998-5>]
- Sahoo, D., Panda, S., Guru, B. C., & Bhatta, K. S. (2008). A new record of Indo-Pacific crab *Charybdis feriata* (Linn., Brachyura: Portunidae) from Chilika Lagoon, Orissa, India. *The Ecoscan, an International Quarterly Journal of Environmental Sciences*, 2(2), 177-179.
- Saint-Paul, U., & Schneider, H. (Eds.). (2010). *Mangrove dynamics and management in North Brazil* (Vol. 211). Springer Science & Business Media. [DOI: https://doi.org/10.1007/978-3-642-13457-9_1]
- Sakthivel, K., & Fernando, A. B. (2012). Brachyuran crabs diversity in Mudasal Odai and Nagapattinam coast of south east India. *Arthropods*, 1(4), 136-143.
- Sánchez, A. J., & Raz-Guzman, A. (1997). Distribution patterns of tropical estuarine brachyuran crabs in the Gulf of Mexico. *Journal of Crustacean Biology*, 17(4), 609-620.
- Sánchez, A. J., & Raz-Guzman, A. (1997). Distribution patterns of tropical estuarine brachyuran crabs in the Gulf of Mexico. *Journal of Crustacean Biology*, 17(4), 609-620.
- Sanchez-Contreras, M., & Vlisidou, I. (2008). The Diversity of Insect-bacteria Interactions and its Applications for Disease Control. *Biotechnology and Genetic Engineering Reviews*, 25(1), 203-244. <https://doi.org/10.5661/bger-25-203>

- Sanders, H. L. (1958). Benthic Studies in Buzzards Bay. I. Animal-Sediment Relationships 1. *Limnology and oceanography*, 3(3), 245-258.
- Sankarankutty, C. (1962) On decapoda brachyura from Andaman and Nicobar islands- 3. Families: Calappidae, Leucosiidae, Parthenopidae, Maiidae and Gecarcinidae. *Journal of the Marine Biological Associations of India*, 4(1), 151-164.
- Sara, L., Aguilar, R. O., Ingles, J. A., & Laureta, L. V. (2014). Habitat characteristics and relative abundance of the mud crab *Scylla serrata* (Forskål, 1775) in Lawele Bay, Southeast Sulawesi, Indonesia. *Su Ürünleri Dergisi*, 31(1), 11-18.
- Sasekumar, A. (1974). Distribution of macrofauna on a Malayan mangrove shore. *The Journal of Animal Ecology*, 51-69.
- Satheeshkumar, P., & Khan, A. B. (2010). An annotated checklist of brachyuran crabs (Crustacea: Decapoda) from Pondicherry Mangroves, southeast coast of India. *World Journal of Zoology*, 6(3), 312-317.
- Sathish, C., Rawat, S., Patel, K. J., Trivedi, J. K., Deshmukhe, G., & Jaiswar, A. K. (2023). First Confirmed Report on the Three Mangrove Associated Crab Species from Maharashtra Coast of India. *Thalassas: An International Journal of Marine Sciences*, 39(2), 1-8. <https://doi.org/10.1007/s41208-023-00589-8>
- Schindler, D. E., & Scheuerell, M. D. (2002). Habitat coupling in lake ecosystems. *Oikos*, 98(2), 177-189. <https://doi.org/10.1034/j.1600-0706.2002.980201.x>
- Schneider, K. R. (2008). Heat Stress in the Intertidal: Comparing Survival and Growth of an Invasive and Native Mussel Under a Variety of Thermal Conditions. *The Biological Bulletin*, 215(3), 253-264. <https://doi.org/10.2307/25470709>

- Schober, U. M., & Christy, J. H. (1993). Sand disposal of the painted ghost crab *Ocypode gaudichaudii* (Decapoda: Ocypodidae): A possible role in courtship. *Marine Biology*, 116(1), 53–60.
<https://doi.org/10.1007/bf00350731>
- Schram, F. R., & Castro, P. (2015). Introduction to Brachyura. In *Treatise on Zoology-Anatomy, Taxonomy, Biology. The Crustacea*, 9(2), 3–9.
- Schubart, C. D., & Diesel, R. (1998). Osmoregulatory capacities and penetration into terrestrial habitats: a comparative study of Jamaican crabs of the genus *Armases* Abele, 1992 (Brachyura: Grapsidae: Sesarminae). *Bulletin of Marine Science*, 62(3), 743–752.
- Schwaibold, U., & Pillay, N. (2005). Behavioral strategies of the African ice rat *Otomys sloggetti robertsi* in the cold. *Physiology & Behavior*, 88(4-5), 567–574.
- Seike, K., & Nara, M. (2008). Burrow morphologies of the ghost crabs *Ocypode ceratophthalma* and *O. sinensis* in foreshore, backshore, and dune sub environments of a sandy beach in Japan. *The Journal of the Geological Society of Japan*, 114(11), 591–596.
<https://doi.org/10.5575/geosoc.114.591>
- Seitz, R. D., Wennhage, H., Bergström, U., Lipcius, R. N., & Ysebaert, T. (2013). Ecological value of coastal habitats for commercially and ecologically important species. *ICES Journal of Marine Science*, 71(3), 648–665.
<https://doi.org/10.1093/icesjms/fst152>
- Sewell, R. B. S. (1934). A study of the fauna of the salt lakes, Calcutta. *Records of the Indian Museum*, 36, 45–121.
- Shen, H., Braband, A., & Scholtz, G. (2013). Mitogenomic analysis of decapod crustacean phylogeny corroborates traditional views on their relationships. *Molecular Phylogenetics and Evolution*, 66(3), 776–789.
<https://doi.org/10.1016/j.ympev.2012.11.002>

- Shet, G. N., Subash Chandran, M. D., & Ramachandra, T. V. (2016). Brachyuran crabs of Aghanashini Estuary, South Indian West Coast, Karnataka. *Proceedings of the Lake 2016: Conference on Conservation and Sustainable Management of Ecologically Sensitive Regions in Western Ghats*, 1–17.
- Shetye, S. R. (1999). Tides in the Gulf of Kutch, India. *Continental Shelf Research*, 19(14), 1771–1782. [https://doi.org/10.1016/s0278-4343\(99\)00038-2](https://doi.org/10.1016/s0278-4343(99)00038-2)
- Shih, H., Mok, H., & Chang, H. (2005). Chimney Building by Male *Uca formosensis* Rathbun, 1921 (Crustacea: Decapoda: Ocypodidae) after Pairing: A New Hypothesis for Chimney Function. *Zoological Studies*, 44(2), 242–251.
- Shinn, E. A. (1968). Burrowing in recent lime sediments of Florida and the Bahamas. *Journal of Paleontology*, 42(4), 879–894.
- Shirley, M. A., Hines, A. H., & Wolcott, T. G. (1990). Adaptive significance of habitat selection by molting adult blue crabs *Callinectes sapidus* (Rathbun) within a sub estuary of central Chesapeake Bay. *Journal of Experimental Marine Biology and Ecology*, 140(1-2), 107-119.
- Shukla, M., Patel, B., Trivedi, J., & Vachhrajani, K. (2013). Brachyuran crabs diversity of Mahi and Dhadhar estuaries, Gujarat, India. *Research Journal of Marine Sciences*, 2321, 1296.
- Silas, E. G., & Sankarankutty, C. (1965). Field investigations on the shore crabs of the Gulf of Mannar and Palk Bay, with special reference to the ecology and behaviour of the pellet crab *Scopimera proxima* Kemp. *Proceedings of Symposium on Crustacea*, 3, 1008–1025.
- Singh, H. S., Pandey, C. N., Yennawar, P., Asari, R. J., Patel, B. H., Tatu, K., & Raval, B. R. (2004). *The Marine National Park and Sanctuary in the Gulf of Kachchh-A comprehensive study on biodiversity and management issues*. (Vol. 46). GEER Foundation, Gandhinagar, Gujarat.
- Sinha, S. (2015). Behavioral Ecology of Crabs: A Brief Review. *IOSR Journal of Environmental Science, Toxicology and Food Technology*, 1(6), 50–54.

- Sinha, S., & Pati, A. K. (2008). Circannual rhythm in spatial distribution of burrows of freshwater crab, *Barytelphusa cunicularis* (Westwood, 1836). *Biological Rhythm Research*, 39(4), 359–368. <https://doi.org/10.1080/09291010701425074>
- Smetacek, V. S. (1985). Role of sinking in diatom life-history cycles: ecological, evolutionary and geological significance. *Marine Biology*, 84(3), 239–251. <https://doi.org/10.1007/bf00392493>
- Smith, J. R., Fong, P., & Ambrose, R. F. (2006). DRAMATIC DECLINES IN MUSSEL BED COMMUNITY DIVERSITY: RESPONSE TO CLIMATE CHANGE? *Ecology*, 87 (5), 1153–1161. [https://doi.org/10.1890/0012-9658\(2006\)87\[1153:ddimbc\]2.0.co;2](https://doi.org/10.1890/0012-9658(2006)87[1153:ddimbc]2.0.co;2)
- Smith, T. J., Boto, K. G., Frusher, S. D., & Giddins, R. L. (1991). Keystone species and mangrove forest dynamics: the influence of burrowing by crabs on soil nutrient status and forest productivity. *Estuarine, Coastal and Shelf Science*, 33(5), 419–432. [https://doi.org/10.1016/0272-7714\(91\)90081-1](https://doi.org/10.1016/0272-7714(91)90081-1)
- Snelgrove, P. V. (1999). Getting to the bottom of marine biodiversity: sedimentary habitats: ocean bottoms are the most widespread habitat on earth and support high biodiversity and key ecosystem services. *BioScience*, 49(2), 129-138.
- Snelgrove, P. V. (1999). Getting to the bottom of marine biodiversity: sedimentary habitats: ocean bottoms are the most widespread habitat on earth and support high biodiversity and key ecosystem services. *BioScience*, 49(2), 129-138.
- Snelgrove, P. V. R., & Butman, C. A., (1994). Animal-Sediment Relationships Revisited: Cause Versus Effect. In: *Oceanography and Marine Biology: Annual Review*, 32, 111-177.
- Snelgrove, P. V., Grassle, J. F., Grassle, J. P., Petrecca, R. F., & Stocks, K. I. (2001). The role of colonization in establishing patterns of community composition and

- diversity in shallow-water sedimentary communities. *Journal of Marine Research*, 59(5), 813-830.
- Snowden, R., Clayton, D., & Al-Taher, E. (1991). Population biology of *Ilyoplax stevensi* (Brachyura: Ocypodidae) on a Kuwait mudflat. *Marine Ecology Progress Series*, 71, 219–225. <https://doi.org/10.3354/meps071219>
- Somero, G. N. (2002). Thermal Physiology and Vertical Zonation of Intertidal Animals: Optima, Limits, and Costs of Living. *Integrative and Comparative Biology*, 42(4), 780–789. <https://doi.org/10.1093/icb/42.4.780>
- Souza, D. G. C., Petracco, M., Danin, A. P. F., & Pombo, M. (2021). Population structure and use of space by ghost crabs (Brachyura: Ocypodidae) on an equatorial, macrotidal sandy beach. *Estuarine, Coastal and Shelf Science*, 258, 107376. [doi: <https://doi.org/10.1016/j.ecss.2021.107376>]
- Sruthi, S., Nansimole, A., Gayathri Devi, T. V. & Radhakrishnan, T. (2014). Diversity of brachyuran crabs of Kanyakumari Area, South West Coasts of India. *International Journal of Scientific Research*, 3, 126-128.
- Steiner, A. J., & Leatherman, S. P. (1981). Recreational impacts on the distribution of ghost crabs *Ocypode quadrata* Fab. *Biological Conservation*, 20, 111–122.
- Stephenson, A. T., & Stephenson, A. (1949). The Universal Features of Zonation Between Tide Marks on Rocky Coasts. *The Journal of Ecology*, 37(2), 289–305. <https://doi.org/10.2307/2256610>
- Stevens, B. A. (1929). Ecological Observations on Callinassidae of Puget Sound. *Ecology*, 10(4), 399–405. <https://doi.org/10.2307/1931148>
- Stillman, J. H., & Somero, G. N. (2000). A Comparative Analysis of the Upper Thermal Tolerance Limits of Eastern Pacific Porcelain Crabs, Genus *Petrolisthes*: Influences of Latitude, Vertical Zonation, Acclimation, and Phylogeny. *Physiological and Biochemical Zoology*, 73(2), 200–208. <https://doi.org/10.1086/316738>

- Suchanek, T. H. (1985). Thalassinid shrimp burrows: ecological significance of species-specific architecture. In *Proceedings of the 5th International Coral Reef Congress, Tahiti, 27 May-1 June 1985* (pp. 205–210). Antenne Museum-EPHE, Moorea, French Polynesia.
- Suess, E. (1980). Particulate organic carbon flux in the oceans—surface productivity and oxygen utilization. *Nature*, 288(5788), 260–263. <https://doi.org/10.1038/288260a0>
- Suvarna Devi, S., Kumar, A. B., & Ravinesh, R. (2014). Diversity of brachyuran crabs of Kavaratti Island, Lakshadweep, with notes on two new reports. *Journal of Aquatic Biology and Fisheries*, 2, 792–794.
- Takahasi, S. (1935). Ecological notes on the Ocypodian crabs (Ocypodidae) in Formosa, Japan. *Annotationes Zoologicae Japonenses*, 15, 78–87.
- Takeda, S., & Kurihara, Y. (1987). The distribution and abundance of *Helice tridens* (De Haan) burrows and substratum conditions in a north eastern Japan salt marsh (Crustacea: Brachyura). *Journal of Experimental Marine Biology and Ecology*, 107(1), 9–19. [https://doi.org/10.1016/0022-0981\(87\)90119-5](https://doi.org/10.1016/0022-0981(87)90119-5)
- Takeda, S., Poovachiranon, S., & Murai, M. (2004). Adaptations for feeding on rock surfaces and sandy sediment by the fiddler crabs (Brachyura: Ocypodidae) *Uca tetragonon* (Herbst, 1790) and *Uca vocans* (Linnaeus, 1758). *Hydrobiologia*, 528, 87–97.
- Tan, C. G. S., & Ng, P. K. L. (1994). An annotated checklist of mangrove brachyuran crabs from Malaysia and Singapore. *Hydrobiologia*, 285(1), 75–84. <https://doi.org/10.1007/BF00005655>
- Tandel, L., Kardani, H., Tandel, B., Parmar, J., Vadher, P., & Kumar, V. (2022). Diversity of Brachyuran Crab Along Sikka Coast of Gujarat. *Journal of Experimental Zoology India*, 25(1), 761–773. <https://connectjournals.com/03895.2022.25.761>

- Taylor, P. R. (1981). Hermit crab fitness: The effect of shell condition and behavioral adaptations on environmental resistance. *Journal of Experimental Marine Biology and Ecology*, 52(2), 205–218. [https://doi.org/10.1016/0022-0981\(81\)90037-X](https://doi.org/10.1016/0022-0981(81)90037-X)
- Thomas, M. M. (1969). Catalogue of crabs and hermit-crabs in the reference collections of the central marine fisheries research institute. *Bulletin of the Central Marine Fisheries Research Institute*, 7, 41–50.
- Thongtham, N., & Kristensen, E. (2003). Physical and chemical characteristics of mangrove crab (*Neoepisesarma versicolor*) burrows in the Bangrong mangrove forest, Phuket, Thailand; with emphasis on behavioural response to changing environmental conditions. *Vie et Milieu*, 53, 141–151.
- Thrush, S. F., Hewitt, J. E., Gladstone-Gallagher, R. V., Savage, C., Lundquist, C. J., O'Meara, T., Vieillard, A. M., Hillman, J. R., Mangan, S., Douglas, E. J., Clark, D., Lohrer, A. M., & Pilditch, C. A. (2021). Cumulative stressors reduce the self-regulating capacity of coastal ecosystems. *Ecological Applications*, 31(1). <https://doi.org/10.1002/eap.2223>
- Thurman, C. L. (1998). Evaporative Water Loss, Corporal Temperature and the Distribution of Sympatric Fiddler Crabs (*Uca*) from South Texas. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 119(1), 279–286. [https://doi.org/10.1016/s1095-6433\(97\)00424-8](https://doi.org/10.1016/s1095-6433(97)00424-8)
- Timmermans, M. C. J. (2002). Hazira Harbour Siltation Study: A new Harbour in the Gulf of Khambhat, India. *TU Delft, Faculty of Civil Engineering and Geosciences, Hydraulic Engineering*.
- Tina, F. W., Jaroensutasinee, M., Jaroensutasinee, K., & Chumsri, A. (2018). Do mudballs around burrows affect burrow characteristics of the fiddler crab *Austruca annulipes* (H. Milne Edwards, 1837) (Brachyura, Ocypodidae)? *Crustaceana*, 91(4), 489–500. <https://doi.org/10.1163/15685403-00003779>

- Trivedi, D. J., Trivedi, J. N., Soni, G. M., Purohit, B. D., & Vachhrajani, K. D. (2015a). Crustacean fauna of Gujarat state of India: A review. *Electronic Journal of Environmental Sciences*, 8, 23-31.
- Trivedi, J. N. (2015). *Distribution, Diversity and Habitat Preference of Brachyuran Crabs along the Coastal Saurashtra, Gujarat*. (pp. 1-268) [Ph. D. Thesis]. <http://hdl.handle.net/10603/64294>
- Trivedi, J. N., & Vachhrajani, K. D. (2012). Distribution and diversity of brachyuran crabs along the coastal region of Junagadh district, Gujarat. *Proceeding of National Seminar on Biodiversity and Conservation of Coastal and Marine Ecosystems of India*, 6-12.
- Trivedi, J. N., & Vachhrajani, K. D. (2013a). First record of *Cryptopodia angulata* H. Milne Edwards and Lucas, 1841 from Saurashtra coast, Gujarat, India (Decapoda: Brachyura: Parthenopidae). *Check List*, 9(4), 897-897. <https://doi.org/10.15560/9.4.897>
- Trivedi, J. N., & Vachhrajani, K. D. (2016). On burrow morphology of the ghost crab, *Ocypode ceratophthalmus* (Decapoda: Brachyura: Ocypodidae) from sandy shore of Gujarat, India. *International Journal of Marine Science*, 6(11), 1-10. <https://doi.org/10.5376/ijms.2016.06.0015>
- Trivedi, J. N., Ahyong, S. T., Vachhrajani, K. D., & Kumar, A. B. (2020). An annotated checklist of the mantis shrimps of India (Crustacea: Stomatopoda). *Zootaxa*, 4768(2), 221-238. <https://doi.org/10.11646/zootaxa.4768.2.4>
- Trivedi, J. N., Bineesh, K. K., & Patel, H. (2023). First Record of *Arcotheres Alcocki* (Rathbun, 1909) (Crustacea: Brachyura: Pinnotheridae) from India. *Thalassas: An International Journal of Marine Sciences*, 39. <https://doi.org/10.1007/s41208-023-00523-y>
- Trivedi, J. N., Gadhavi, M. K., & Vachhrajani, K. D. (2012). Diversity and habitat preference of brachyuran crabs in Gulf of Kutch, Gujarat, India. *Arthropods*, 1(1), 13-23.

- Trivedi, J. N., Soni, G. M., & Vachhrajani, K. D. (2015). First record of brachyuran crab *Heteropanope glabra* Stimpson, 1858 (Crustacea, Decapoda, Pilumnidae) from India. *Marine Biodiversity Records*, 8, 1–5.
<https://doi.org/10.1017/s1755267215000883>
- Trivedi, J. N., Soni, G. M., & Vachhrajani, K. D. (2016). A new species of Lyphira Galil, 2009 (Crustacea, Decapoda, Leucosiidae) from Gujarat, India. *Tropical Zoology*, 29(3). <https://doi.org/10.1080/03946975.2016.1181954>
- Trivedi, J. N., Soni, G. M., Trivedi, D. J., & Vachhrajani, K. D. (2015c). A new species of *Ilyoplax* (Decapoda, Brachyura, Dotillidae) from Gujarat, India. *Journal of Asia-Pacific Biodiversity*, 8(2), 173-177.
<https://doi.org/10.1016/j.japb.2015.02.005>
- Trivedi, J. N., Trivedi, D. J., Vachhrajani, K. D., & Ng, P. K. L. (2018). An annotated checklist of the marine brachyuran crabs (Crustacea: Decapoda: Brachyura) of India. *Zootaxa*, 4502(1), 1–83.
<https://doi.org/10.11646/zootaxa.4502.1.1>
- Trivedi, J.N. & Vachhrajani, K.D. (2013b) Taxonomic account of genus *Scylla* (de Haan, 1833) from Gujarat State, India with two new records of species. *Arthropods*, 2(4), 159–171.
- Trivedi, J.N. & Vachhrajani, K.D. (2018) On distributional range extension of two species of brachyuran crabs *Perasesarma persicum* Naderloo & Schubart, 2010 and *Austruca iranica* (Pertzmann, 1971) in Indian waters. *Journal of Marine Biological Association of India*, 59(2), 20–24.
- Trivedi, J.N., Soni, G.M. & Vachhrajani, K.D. (2015b) First record of brachyuran crab *Heteropanope glabra* Stimpson, 1858 (Crustacea, Decapoda, Pilumnidae) from India. *Marine Biodiversity Records*, 8, 1–5.
<https://doi.org/10.1017/S1755267215000883>
- Trivedi, J.N., Soni, G.M., Arya, S. & Vachhrajani, K.D. (2014) First record of *Macrophthalmus laevis* A. Milne Edwards, 1867 (Decapoda: Brachyura:

- Macrophthalmidae) from India. *Journal of Marine Biological Association of India*, 56(2), 85–87. <https://doi.org/10.6024/jmbai.2014.56.2.01789-13>
- Trueman, E. R. (1970). The Mechanism of Burrowing of the Mole Crab, *Emerita*. *Journal of Experimental Biology*, 53(3), 701–710. <https://doi.org/10.1242/jeb.53.3.701>
- Turra, A., & Denadai, M. R. (2003). Daily activity of four tropical intertidal hermit crabs from Southeastern Brazil. *Brazilian Journal of Biology*, 63(3), 537–544. <https://doi.org/10.1590/s1519-69842003000300020>
- Tweedie, M. W. F. (1950). Notes on grapsoid crabs from the Raffles Museum. *Bulletin of the Raffles Museum*, 23, 310–324.
- Upadhyay, K. S., Patel, K. J., Prajapati, J. M., Rabari, V. M., Thacker, D. R., Patel, H. V., & Trivedi, J. N. (2022). Burrow Morphology of Brachyuran Crab *Dotilla blanfordi* Alcock, 1900 from Gulf of Khambhat, Gujarat, India. *International Journal of Zoological Investigations*, 08(02), 251–261. <https://doi.org/10.33745/ijzi.2022.v08i02.032>
- Vaidya, G. K. (2022). *Crab Ecology as Potential Bio-Monitoring Tool: Studies on Population and Behaviour Ecology of Ilyoplax sayajiraoi (Crustacea: Brachyura: Dotillidae)* (Doctoral dissertation, Maharaja Sayajirao University of Baroda (India)).
- Valero-Pacheco, E., Alvarez, F., Escobar, M., & Abarca-Arenas, L. (2007). Population density and activity pattern of the ghost crab, *Ocypode quadrata*, in Veracruz, Mexico. *Crustaceana*, 80(3), 313–325. <https://doi.org/10.1163/156854007780162479>
- Vannini, M. (1980). Researches on the Coast of Somalia. The Shore and the Dune of Sar Uanle. *Monitore Zoologico Italiano. Supplemento*, 13(1), 11–44. <https://doi.org/10.1080/00269786.1980.11758547>

- Varadharajan, D., & Soundarapandian, P. (2014). Crab biodiversity from Arukkattuthurai to Pasipattinam, southeast coast of India. *Indian Journal of Geo-Marine Sciences*, 43(4), 676–698.
- Varadharajan, D., Soundarapandian, P., & Pushparajan, N. (2013). The global science of crab biodiversity from Puducherry coast, south east coast of India. *Arthropods*, 2(1), 26–35.
- Vaugelas, J. D. (1984). Preliminary observations on two types of callianassid (Crustacea, Thalassinidea) burrows at Gulf of Aqaba (Red Sea). In M. A. H. Saad (Ed.), *Proceeding of the Symposium on Coral Reef Environments of the Red Sea*: (pp. 520–539). King Abdulaziz University Press, Jeddah.
- Velip, D. T., & Rivonker, C. U. (2015). *Hexapus bidentatus* sp. nov. (Crustacea: Decapoda: Brachyura: Hexapodidae), a new species from Goa, west coast of India. *Marine Biology Research*, 11(1), 97–105. <https://doi.org/10.1080/17451000.2014.889305>
- Venkataraman, K., & Raghunathan, C. (2015). Coastal and marine biodiversity of India. In K. Venkataraman & C. Sivaperuman (Eds.), *Marine Faunal Diversity in India* (pp. 303–348). Academic Press, India.
- Venkataraman, K., Jeyabaskaran, R., Raghuram, K. P., & Alfred, J. R. b. (2004). Bibliography and Checklist of Coral and Associated Organisms of India. *Records of Zoological Survey of India, Occasional Paper*, 226, 1–468.
- Venkataraman, K., Srinivasan, M., Satyanarayana, Ch. & Prabakar, D. (2002) Faunal diversity of Gulf of Mannar Biosphere Reserve. In: *Director ZSI, Kolkata (Ed.), Conservation Area Series, Zoological Survey of India, Kolkata, Vol. 15*, 1– 77.
- Vidhya, V., Jawahar, P., & Karuppasamy, K. (2017). Annotated check list of the brachyuran crabs (Crustacea: Decapoda) from Gulf of Mannar region, south east coast of India. *Journal of Entomology and Zoology Studies*, 5(6), 2331–2336.

- Violle, C., Reich, P. B., Pacala, S. W., Enquist, B. J., & Kattge, J. (2014). The emergence and promise of functional biogeography. *Proceedings of the National Academy of Sciences*, 111(38), 13690-13696. [doi:<https://doi.org/10.1073/pnas.1415442111>]
- Vogt, G. (2012). Ageing and longevity in the Decapoda (Crustacea): A review. *Zoologischer Anzeiger - a Journal of Comparative Zoology*, 251(1), 1-25. <https://doi.org/10.1016/j.jcz.2011.05.003>
- Wada, K. (1983). Spatial distributions and population structures in *Scopimera globosa* and *Ilyoplax pusillus* (Decapoda: Ocypodidae). *Publications of the Seto Marine Biological Laboratory*, 27(4-6), 281-291.
- Wada, K. (1984). Barricade building in *Ilyoplax pusilla* (De Haan) (Crustacea: Brachyura). *Journal of Experimental Marine Biology and Ecology*, 83, 73-88.
- Wada, K. (1994). Earthen structures built by *Ilyoplax pusilla* (Crustacea: Brachyura: Ocypodidae). *Ethology*, 96, 270-282.
- Wada, K., & Murata, I. (2000). Chimney Building in the Fiddler Crab *Uca Arcuata*. *Journal of Crustacean Biology*, 20(3), 505-509. <https://doi.org/10.1163/20021975-99990066>
- Wada, K., Yum, S. S., & Park, J. K. (1994). Mound building in *Ilyoplax pingi* (Crustacea: Brachyura: Ocypodidae). *Marine Biology*, 121(1), 61-65. <https://doi.org/10.1007/bf00349474>
- Wang, J. Q., Zhang, X. D., Jiang, L. F., Bertness, M. D., Fang, C. M., Chen, J. K., Hara, T., & Li, B. (2010). Bioturbation of Burrowing Crabs Promotes Sediment Turnover and Carbon and Nitrogen Movements in an Estuarine Salt Marsh. *Ecosystems*, 13(4), 586-599. <https://doi.org/10.1007/s10021-010-9342-5>
- Warren, J. H. (1990). The use of open burrows to estimate abundances of intertidal estuarine crabs. *Australian Journal of Ecology*, 15(3), 277-280.

- Weinstein, R. B. (1995). Locomotor behavior of nocturnal ghost crabs on the beach: focal animal sampling and instantaneous velocity from three-dimensional motion analysis. *Journal of Experimental Biology*, 198(4), 989–999. <https://doi.org/10.1242/jeb.198.4.989>
- Weinstein, R. B. (1998). Effects of Temperature and Water Loss on Terrestrial Locomotor Performance in Land Crabs: Integrating Laboratory and Field Studies. *American Zoologist*, 38(3), 518–527. <https://www.jstor.org/stable/4620172>
- Welsh, J. H. (1939). The action of eyestalk extracts on retinal pigment migration in the crayfish, *Cambarus bartoni*. *The Biological Bulletin*, 77, 119–125.
- Wethey, D. S. (1983). Geographic Limits And Local Zonation: The Barnacles *Semibalanus* (*Balanus*) and *Chthamalus* New England. *The Biological Bulletin*, 165(1), 330–341. <https://doi.org/10.2307/1541373>
- Whiting, N. H., & Moshiri, G. A. (1974). Certain organism-substrate relationships affecting the distribution of *Uca minax* (Crustacea: Decapoda). *Hydrobiologia*, 44(4), 481–493. [doi: <https://doi.org/10.1007/BF00036312>]
- Widdicombe, S., Austen, M. C., Kendall, M. A., Warwick, R. M., & Jones, M. B. (2000). Bioturbation as a mechanism for setting and maintaining levels of diversity in subtidal macrobenthic communities. *Springer EBooks*, 440((1-3)), 369–377. https://doi.org/10.1007/978-94-017-1982-7_34
- Wolff, G. A. (1989). Productivity of the oceans: Present and past. In W. H. Berger, V. S. Smetacek, & G. Wefer (Eds.), *Report of the Dahlem Workshop on Productivity of the Oceans* (p. 471). Wiley, Berlin.
- Wolfrath, B. (1992). Burrowing of the fiddler crab *Uca tangeri* in the Ria Formosa in Portugal and its influence on sediment structure. *Marine Ecology Progress Series*, 85, 237–243.

- Xin, P., Jin, G., Li, L., & David Andrew Barry, D. A. (2009). Effects of crab burrows on pore water flows in salt marshes. *Advances in Water Resource*, 32(3), 439–449. <https://doi.org/10.1016/j.advwatres.2008.12.008>
- Yamada, K., Hori, M., Tanaka, Y., Hasegawa, N., & Nakaoka, M. (2007). Temporal and spatial macrofaunal community changes along a salinity gradient in seagrass meadows of Akkeshi-ko estuary and Akkeshi Bay, northern Japan. *Hydrobiologia*, 592, 345-358.
- Zhang, X., Siveter, D. J., Waloszek, D., & Maas, A. (2007). An epipodite-bearing crown-group crustacean from the Lower Cambrian. *Nature*, 449(7162), 595–598. <https://doi.org/10.1038/nature06138>
- Zolkhiflee, N., Yahya, K., & Shuib, S. (2021). Intertidal zone preferences of fiddler crabs in tropical mangroves reflect species specific selection across multiple spatial and temporal scales. *Regional Studies in Marine Science*, 48, 101994. [doi: <https://doi.org/10.1016/j.rsma.2021.101994>]
- Zucker, N. (1981). The role of hood-building in defining territories and limiting combat in fiddler crabs. *Animal Behaviour*, 29(2), 387–395. [https://doi.org/10.1016/s0003-3472\(81\)80098-x](https://doi.org/10.1016/s0003-3472(81)80098-x)