***** Table of Contents:

Sr. No.	Content	Page No.
1	Introduction and Literature Review	1-24
2	Materials and Methodology	25-33
3	Results and Discussion	34-162
	Chapter 1: Brachyuran Crab Diversity on Kamboi Coast	34-50
	Chapter 2: Brachyuran Crab Distribution on Kamboi Coast	51-67
	Chapter 3: Burrow Architecture of Brachyuran Crabs	68-129
	Chapter 4: Effect of Lunar Cycle on Burrowing Behaviour	130-143
	Chapter 5: Foraging Behaviour and Bioturbation Activity	144-165
4	Conclusion	166-169
5	Bibliography	170-225
6	Publication and Presentations	

* Table of Figures:

Figure	Title	Page
No	Tiuc	No.
1.1	Classification of marine environment	8
2.1	Location of the study site: Kamboi Coast, Gulf of Khambhat,	25
	Gujarat, India	
2.2	Descriptive map of study area. A & B: Entire region of Gulf of	26
	Khambhat showing its peculiar structural features, C: Apical	
	most northern region of the gulf showing the mouth of Mahi and	
	Sabarmati Rivers, D: The entire survey area from coastal village	
	Dahegam (Left star) to Nahar (Right star), E: the area along	
	coast of Kamboi which was actually studied for the present	
	research.	
2.3	Scheme of sampling of counting and measuring of burrows of	31
	Austruca sindensis (Alcock, 1900), Ilyoplax sayajiraoi Trivedi,	
	Soni, Trivedi & Vachhrajani, 2015 and <i>Dotilla blanfordi</i> Alcock	
	1900 during duration of tidal exposure on various lunar phases	
	at Mud-flats of Gulf of Khambhat, Gujarat, India. Q: quadrat, TR:	
	transect	
3.1	Dotilla blanfordi Alcock, 1900 (A-Dorsal surface; B-Ventral	46
	surface); Macrophthalmus sulcatus H. Milne Edwards, 1852 (C-	
	Dorsal surface; D- Ventral surface)	
3.2	Ashtoret lunaris (Forskål, 1775) (A-Dorsal surface; B-Ventral	46
	surface); Austruca sindensis (Alcock, 1900) (C-Dorsal surface; D-	
	Ventral surface)	
3.3	Eurycarcinus orientalis A. Milne-Edwards, 1867 (A-Dorsal	47
	surface; B-Ventral surface); Ilyoplax sayajiraoi Trivedi, Soni,	
	Trivedi & Vachhrajani, 2015 (C-Dorsal surface; D- Ventral	
	surface)	
3.4	Parasesarma persicum Naderloo & Schubart, 2010 (A-Dorsal	47
	surface; B-Ventral surface); Scylla serrata (Forskål, 1775) (C-	
	Dorsal surface; D- Ventral surface)	

3.	5	Austruca iranica Alcock 1900; B. Heteropanope glabra Stimpson, 1958	48
4.	1	Different zones of intertidal area based on sediment	56
••	_	composition at Kamboi Coast, Gulf of Khambhat, Gujarat, India.	50
4.	2	Seasonal variation in the mean values of temperature at	58
		different intertidal region	
4.	3	Seasonal variation in burrow density of different burrowing	59
		crab species along the intertidal region of Kamboi coast.	
4.	4	Seasonal variation in burrow abundance of different burrowing	59
		crab species along the intertidal region of Kamboi coast.	
4.	5	Seasonal variation in frequency of burrow occurrence of	60
		different burrowing crab species along the intertidal region of	
		Kamboi coast.	
4.	6	Intertidal distribution of six burrowing crab species at Kamboi	61
		coast.	
4.	7	Intertidal distribution of three abundantly distributed	62
		brachyuran crab species at Kamboi coast in B. winter season; C.	
		Summer season and D. Monsoon season.	
5.	1	Burrow architecture of Austruca sindensis (Alcock, 1900) with	74
		various measurements of burrow cast. a. J-shaped burrow, b.	
		Spiral burrow, c. S-shaped burrow, d. Single tube burrow, e. U-	
		shaped burrow, f. Multi-branched burrow, g. J-shaped with	
		Branch burrow. (TDB - Total depth of Burrow; TLB - Total	
		length of burrow; IA- Burrow inclination angle; CL - Chamber	
		length; CW – Chamber width)	
5.	2	Indicates number of burrow count with respect to shape of the	75
		burrow in Austruca sindensis (Alcock, 1900).	
5.	3	Regression analysis for the relationship between crab carapace	80
		length and crab carapace width (CW) and Burrow opening	
		diameter (BOD) Austruca sindensis (Alcock, 1900) in winter	
		season.	

5.4	Regression analysis for the relationship between crab carapace	81
	length and Total length (TLB) and Total depth (TDB) of burrow	
	burrows of Austruca sindensis (Alcock, 1900) in winter season.	
5.5	Regression analysis for the relationship between crab carapace	82
	length and Burrow volume (BV) and Inclination angle (IA) of	
	burrows of Austruca sindensis (Alcock, 1900) in winter season.	
5.6	Vertical temperature profiles with the burrow depth in S-	83
	shaped and J-shaped burrows of Austruca sindensis (Alcock,	
	1900) during winter season.	
5.7	Vertical temperature profiles with the burrow depth in spiral	84
	and single tube (ST) burrows of Austruca sindensis (Alcock,	
	1900) during winter season.	
5.8	Scree plot indicating major 2 plots for winter season	85
5.9	Component plot in rotated space for winter season	86
5.10	Regression analysis for the relationship between crab carapace	88
	length and crab carapace width (CW) and Burrow opening	
	diameter (BOD) Austruca sindensis (Alcock, 1900) in summer	
	season.	
5.11	Regression analysis for the relationship between crab carapace	89
	length and Total length (TLB) and Total depth (TDB) of burrow	
	burrows of Austruca sindensis (Alcock, 1900) in summer season.	
5.12	Regression analysis for the relationship between crab carapace	90
	length and Burrow volume (BV) and Inclination angle (IA) of	
	burrows of Austruca sindensis (Alcock, 1900) in summer season.	
5.13	Vertical temperature profiles with the burrow depth in S-	91
	shaped and J-shaped burrows of Austruca sindensis (Alcock,	
	1900) during summer season.	
5.14	Vertical temperature profiles with the burrow depth in single	92
	tube (ST) burrows of Austruca sindensis (Alcock, 1900) during	
	summer season	
5.15	Scree plot indicating major 2 plots for summer season	93
5.16	Component plot in rotated space for summer season	93

5.17	Regression analysis for the relationship between crab carapace	96
	length and crab carapace width (CW) and Burrow opening	
	diameter (BOD) Austruca sindensis (Alcock, 1900) in monsoon	
	season.	
5.18	Regression analysis for the relationship between crab carapace	97
	length and Total length (TLB) and Total depth (TDB) of burrow	
	burrows of Austruca sindensis (Alcock, 1900) in monsoon	
	season.	
5.19	Regression analysis for the relationship between crab carapace	98
	length and Burrow volume (BV) and Inclination angle (IA) of	
	burrows of Austruca sindensis (Alcock, 1900) in monsoon	
	season.	
5.20	Vertical temperature profiles with the burrow depth in single	99
	tube (ST) burrows of Austruca sindensis (Alcock, 1900) during	
	monsoon season.	
5.21	Vertical temperature profiles with the burrow depth in I-shaped	100
	and S-shaped burrows of Austruca sindensis (Alcock, 1900)	
	during monsoon season.	
5.22	Scree plot indicating major 2 plots for monsoon season	101
5.23	Component plot in rotated space for monsoon season	102
5.24	Regression analysis for the relationship status between crab	108
	carapace length and burrow opening diameter of Ilyoplax	
	sayajiraoi JN Trivedi, Soni, DJ Trivedi & Vachhrajani, 2015	
	burrows during pre-monsoon season.	
5.25	Regression analysis for the relationship between crab carapace	109
	length and total length; total depth of burrow of Ilyoplax	
	sayajiraoi JN Trivedi, Soni, DJ Trivedi & Vachhrajani, 2015	
	during pre-monsoon season.	
5.26	Regression analysis for the relationship between crab carapace	110
	length and burrow volume; Inclination angle of burrow of	
	Ilyoplax sayajiraoi JN Trivedi, Soni, DJ Trivedi & Vachhrajani,	
	2015 during pre-monsoon season.	

5.27	Regression analysis for the relationship status between crab	111
	carapace length and burrow opening diameter; total length of	
	burrow (TLB) of <i>Ilyoplax sayajiraoi</i> JN Trivedi, Soni, DJ Trivedi &	
	Vachhrajani, 2015 during post-monsoon season.	
5.28	Regression analysis for the relationship status between crab	112
	carapace length and total depth of burrow (TDB); burrow	
	volume of <i>Ilyoplax sayajiraoi</i> JN Trivedi, Soni, DJ Trivedi &	
	Vachhrajani, 2015 during post-monsoon season.	
5.29	Regression analysis for the relationship status between crab	113
	carapace length and inclination angle of burrow of Ilyoplax	
	sayajiraoi JN Trivedi, Soni, DJ Trivedi & Vachhrajani, 2015	
	during post-monsoon season.	
5.30	Mean variation in Carapace length and burrow opening	114
	diameter of <i>Ilyoplax sayajiraoi</i> JN Trivedi, Soni, DJ Trivedi &	
	Vachhrajani, 2015 between pre-monsoon and post-monsoon.	
5.31	Mean variation in total length and total depth of burrow of	115
	Ilyoplax sayajiraoi JN Trivedi, Soni, DJ Trivedi & Vachhrajani,	
	2015 between pre-monsoon and post-monsoon.	
5.32	Mean variation in burrow volume and inclination angle of	116
	burrow of <i>Ilyoplax sayajiraoi</i> JN Trivedi, Soni, DJ Trivedi &	
	Vachhrajani, 2015 between pre-monsoon and post-monsoon.	
5.33	Mean variation in Carapace length and burrow opening	117
	diameter between male and female host crab individuals of	
	Ilyoplax sayajiraoi JN Trivedi, Soni, DJ Trivedi & Vachhrajani,	
	2015 during pre-monsoon.	
5.34	Mean variation in total length and total depth of burrow	118
	between male and female host crab individuals of Ilyoplax	
	sayajiraoi JN Trivedi, Soni, DJ Trivedi & Vachhrajani, 2015	
	during pre-monsoon.	
5.35	Mean variation in burrow volume and burrow inclination angle	119
	between male and female host crab individuals of Ilyoplax	
	sayajiraoi JN Trivedi, Soni, DJ Trivedi & Vachhrajani, 2015	
	during pre-monsoon.	

5.36	Mean variation in Carapace length and burrow opening	120
	diameter between male and female host crab individuals of	
	Ilyoplax sayajiraoi JN Trivedi, Soni, DJ Trivedi & Vachhrajani,	
	2015 during post-monsoon.	
5.37	Mean variation in total length and total depth of burrow	121
	between male and female host crab individuals of Ilyoplax	
	sayajiraoi JN Trivedi, Soni, DJ Trivedi & Vachhrajani, 2015	
	during post-monsoon.	
5.38	Mean variation in burrow volume and burrow inclination angle	122
	between male and female host crab individuals of Ilyoplax	
	sayajiraoi JN Trivedi, Soni, DJ Trivedi & Vachhrajani, 2015	
	during post-monsoon.	
5.39	Vertical temperature profiles with the depth of burrow pit	123
	during pre-monsoon	
5.40	Vertical temperature profiles with the depth of burrow pit	124
	during post-monsoon	
6.1	Representative image of eight phase of Lunar cycle.	132
6.2	Average densities of Austruca sindensis burrows during the	135
	lunar cycle on Kamboi coast, Gulf of Khambhat, Gujarat.	
6.3	Average densities of Ilyoplax sayajiraoi burrows during the	135
	lunar cycle on Kamboi coast, Gulf of Khambhat, Gujarat.	
6.4	Average densities of <i>Dotilla blanfordi</i> burrows during the lunar	136
	cycle on Kamboi coast, Gulf of Khambhat, Gujarat.	
6.5	Average diameter of Austruca sindensis burrows during the	136
	lunar cycle on Kamboi coast, Gulf of Khambhat, Gujarat.	
6.6		
	Average diameter of Ilyoplax sayajiraoi burrows during the	137
	Average diameter of <i>Ilyoplax sayajiraoi</i> burrows during the lunar cycle on Kamboi coast, Gulf of Khambhat, Gujarat.	137
6.7		137137
6.7	lunar cycle on Kamboi coast, Gulf of Khambhat, Gujarat.	
6.7	lunar cycle on Kamboi coast, Gulf of Khambhat, Gujarat. Average diameter of <i>Dotilla blanfordi</i> burrows during the lunar	
	lunar cycle on Kamboi coast, Gulf of Khambhat, Gujarat. Average diameter of <i>Dotilla blanfordi</i> burrows during the lunar cycle on Kamboi coast, Gulf of Khambhat, Gujarat.	137

6.9	Hour wised variation in densities of <i>Austruca sindensis</i> burrows	140
	during the lunar cycle on Kamboi coast, Gulf of Khambhat,	
	Gujarat: A- 0hr; B- 1.5hr; C-3hr; D-4.5hr	
6.10	Hour wised variation in densities of Austruca sindensis burrows	140
	during the lunar cycle on Kamboi coast, Gulf of Khambhat,	
	Gujarat: A- 3hr; B- 4.5hr	
7.1	Representative illustration of the role of burrowing crab in	145
	nutrient recycling	
7.2	Regression analysis for the relationship of crab Carapace Length	153
	(CL) with burrow opening diameter, total area of foraging, and	
	total number of mudballs of burrows of Austruca sindensis	
	(Alcock, 1900).	
7.3	Regression analysis for the relationship of crab Carapace Length	154
	(CL) with total weight of mudballs, average size of mudballs, and	
	distance between mudballs of burrows of Austruca sindensis	
	(Alcock, 1900).	
7.4	Regression analysis for the relationship of crab carapace length	155
	(CL) with burrow opening diameter, total area of foraging, and	
	total number of mudballs of burrows of Ilyoplax sayajiraoi	
	Trivedi, Soni, Trivedi & Vachhrajani, 2015.	
7.5	Regression analysis for the relationship of crab Carapace Length	156
	(CL) with total weight of mudballs, average size of mudballs, and	
	distance between mudballs of burrows of Ilyoplax sayajiraoi	
	Trivedi, Soni, Trivedi & Vachhrajani, 2015.	
7.6	Regression analysis for the relationship between Burrow	159
	opening diameter and Dry weight of Mudballs of the individual	
	burrow of Austruca sindensis (Alcock, 1900).	
7.7	Regression analysis for the relationship between Burrow	159
	opening diameter and Dry weight of Mudballs of the individual	
	burrow of <i>Ilyoplax sayajiraoi</i> Trivedi, Soni, Trivedi &	
	Vachhrajani, 2015.	
7.8	Average daily output of bioturbation potential of different crab	160
	species per square meter	

❖ *Table of Tables:*

Table No Title	Page No.
3.1 List of the brachyuran crab species recorded.	38
2.1 Represents four distinct zones at Kamboi coast.	55
3.1 Burrow characteristics of <i>Austruca sindensis</i> (Alcock, 1900)	77
3.2 Indicates Pearson correlation between various burrow	78
parameters of Austruca sindensis (Alcock, 1900) for Winter	
season in Zone I	
3.3 Indicates Pearson correlation between various burrow	79
parameters of Austruca sindensis (Alcock, 1900) for Winter	
season in Zone II	
3.4 Total Variance in Winter season in Austruca sindensis	85
(Alcock, 1900)	
3.5 Indicates Pearson correlation between various burrow	87
parameters of Austruca sindensis (Alcock, 1900) for Summer	
season in Zone I	
3.6 Indicates Pearson correlation between various burrow	87
parameters of Austruca sindensis (Alcock, 1900) for Summer	
season in Zone II	
3.7 Total Variance in Summer season in Austruca sindensis	92
(Alcock, 1900)	
3.8 Indicates Pearson correlation between various burrow	94
parameters of Austruca sindensis (Alcock, 1900) for	
Monsoon season in Zone I	
3.9 Indicates Pearson correlation between various burrow	95
parameters of Austruca sindensis (Alcock, 1900) for	
Monsoon season in Zone II	

3.10	Total Variance in Monsoon season in <i>Austruca sindensis</i> (Alcock, 1900)	101
3.11	Burrow parameters of <i>Ilyoplax sayajiraoi</i> JN Trivedi, Soni, DJ	106
0.11	Trivedi & Vachhrajani, 2015 during pre-monsoon. (CL:	200
	Carapace length; CW: Carapace width; BOD: Burrow opening	
	diameter; TLB: Total length of burrow; TDB: Total depth of	
	burrow; BV: Burrow volume; IA: Inclination angle; B1l:	
	Branch 1 length; ChL: Chamber length; ChW: Chamber	
	width)	
3.12	Burrow parameters of <i>Ilyoplax sayajiraoi</i> JN Trivedi, Soni, DJ	107
	Trivedi & Vachhrajani, 2015 during post-monsoon. (CL:	
	Carapace length; CW: Carapace width; BOD: Burrow opening	
	diameter; TLB: Total length of burrow; TDB: Total depth of	
	burrow; BV: Burrow volume; IA: Inclination angle; B1l:	
	Branch 1 length; ChL: Chamber length; ChW: Chamber	
	width)	
4.1	Temperature variation (maximum and minimum) observed	133
	on Kamboi coast, Gulf of Khambhat, Gujarat, during each	
	lunar phase	
5.1	Variation in foraging and mud balling pattern between	152
	different sexes of Austruca sindensis (Alcock, 1900). (BOD:	
	burrow opening diameter; TFA: total foraging area; TNM:	
	total number of mudballs; TWM: total weight of mudballs;	
	AMS: average mudball size; ADM: average distance between	
	mudballs; SW: seaward; LW: landward)	
5.2	Variation in foraging and mud balling pattern between	158
	different sexes of <i>Ilyoplax sayajiraoi</i> Trivedi, Soni, Trivedi &	
	Vachhrajani, 2015. (BOD: burrow opening diameter; TFA:	
	total foraging area; TNM: total number of mudballs; TWM:	
	total weight of mudballs; AMS: average mudball size; ADM:	
	average distance between mudballs; SW: seaward; LW:	
	landward)	