

Bibliography

- [1] R. Aaij, et al., Measurement of the shape of the $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$ differential decay rate, JHEP 12 (2020) 144. [arXiv:2003.08453](#), [doi:10.1007/JHEP12\(2020\)144](#).
- [2] W.-F. Wang, Z.-J. Xiao, The semileptonic decays $B/B_s \rightarrow (\pi, K)(\ell^+ \ell^-, \ell \nu, \nu \bar{\nu})$ in the perturbative QCD approach beyond the leading-order, Phys. Rev. D 86 (2012) 114025. [arXiv:1207.0265](#), [doi:10.1103/PhysRevD.86.114025](#).
- [3] R. N. Faustov, V. O. Galkin, Rare $B \rightarrow \pi l \bar{l}$ and $B \rightarrow \rho l \bar{l}$ decays in the relativistic quark model, Eur. Phys. J. C 74 (6) (2014) 2911. [arXiv:1403.4466](#), [doi:10.1140/epjc/s10052-014-2911-6](#).
- [4] R. N. Faustov, V. O. Galkin, Charmless weak B_s decays in the relativistic quark model, Phys. Rev. D 87 (9) (2013) 094028. [arXiv:1304.3255](#), [doi:10.1103/PhysRevD.87.094028](#).
- [5] J.-J. Wang, R.-M. Wang, Y.-G. Xu, Y.-D. Yang, The Rare decays $B+(u) \rightarrow \pi + l + l^-, \rho + l + l^-$, $B_0(d) \rightarrow l + l^-$ in the R-parity violating supersymmetry, Phys. Rev. D 77 (2008) 014017. [arXiv:0711.0321](#), [doi:10.1103/PhysRevD.77.014017](#).
- [6] D. Melikhov, B. Stech, Weak form-factors for heavy meson decays: An Update, Phys. Rev. D 62 (2000) 014006. [arXiv:hep-ph/0001113](#), [doi:10.1103/PhysRevD.62.014006](#).
- [7] Y.-L. Wu, M. Zhong, Y.-B. Zuo, $B(s), D(s) \rightarrow \pi, K, \eta, \rho, K^*, \omega, \phi$ Transition Form Factors and Decay Rates with Extraction of the CKM

- parameters — $V(ub)$ —, — $V(cs)$ —, — $V(cd)$ —, Int. J. Mod. Phys. A 21 (2006) 6125–6172. [arXiv:hep-ph/0604007](#), doi:10.1142/S0217751X06033209.
- [8] R. C. Verma, Decay constants and form factors of s-wave and p-wave mesons in the covariant light-front quark model, J. Phys. G 39 (2012) 025005. [arXiv:1103.2973](#), doi:10.1088/0954-3899/39/2/025005.
- [9] P. Ball, R. Zwicky, $B_{d,s} \rightarrow \rho, \omega, K^*, \phi$ decay form-factors from light-cone sum rules revisited, Phys. Rev. D 71 (2005) 014029. [arXiv:hep-ph/0412079](#), doi:10.1103/PhysRevD.71.014029.
- [10] Q. Chang, X.-N. Li, L.-T. Wang, Revisiting the form factors of $P \rightarrow V$ transition within the light-front quark models, Eur. Phys. J. C 79 (5) (2019) 422. [arXiv:1905.05098](#), doi:10.1140/epjc/s10052-019-6949-3.
- [11] J. M. Flynn, T. Izubuchi, T. Kawanai, C. Lehner, A. Soni, R. S. Van de Water, O. Witzel, $B \rightarrow \pi\ell\nu$ and $B_s \rightarrow K\ell\nu$ form factors and $|V_{ub}|$ from 2+1-flavor lattice QCD with domain-wall light quarks and relativistic heavy quarks, Phys. Rev. D 91 (7) (2015) 074510. [arXiv:1501.05373](#), doi:10.1103/PhysRevD.91.074510.
- [12] M. Tanabashi, et al., Review of Particle Physics, Phys. Rev. D 98 (3) (2018) 030001. doi:10.1103/PhysRevD.98.030001.
- [13] G. Buchalla, A. J. Buras, M. E. Lautenbacher, Weak decays beyond leading logarithms, Rev. Mod. Phys. 68 (1996) 1125–1144. [arXiv:hep-ph/9512380](#), doi:10.1103/RevModPhys.68.1125.
- [14] S.-P. Jin, X.-Q. Hu, Z.-J. Xiao, Study of $B_s \rightarrow K^{(*)}\ell^+\ell^-$ decays in the PQCD factorization approach with lattice QCD input, Phys. Rev. D 102 (1) (2020) 013001. [arXiv:2003.12226](#), doi:10.1103/PhysRevD.102.013001.
- [15] R. R. Horgan, Z. Liu, S. Meinel, M. Wingate, Lattice QCD calculation of form factors describing the rare decays $B \rightarrow K^*\ell^+\ell^-$ and $B_s \rightarrow \phi\ell^+\ell^-$, Phys. Rev. D 89 (9) (2014) 094501. [arXiv:1310.3722](#), doi:10.1103/PhysRevD.89.094501.
- [16] F. L. Wilson, Fermi’s Theory of Beta Decay, Am. J. Phys. 36 (12) (1968) 1150–1160. doi:10.1119/1.1974382.

- [17] S. L. Glashow, Partial Symmetries of Weak Interactions, *Nucl. Phys.* 22 (1961) 579–588. doi:10.1016/0029-5582(61)90469-2.
- [18] A. Salam, J. C. Ward, Electromagnetic and weak interactions, *Phys. Lett.* 13 (1964) 168–171. doi:10.1016/0031-9163(64)90711-5.
- [19] A. Salam, Weak and Electromagnetic Interactions, *Conf. Proc. C* 680519 (1968) 367–377. doi:10.1142/9789812795915_0034.
- [20] A. Astbury, FOUND: W AND Z. THE DISCOVERY OF THE W AND Z INTERMEDIATE VECTOR BOSONS AT CERN'S PROTON ANTI-PROTON COLLIDER, *Phys. Bull.* 34 (1983) 434–436.
- [21] M. Z. Akrawy, et al., Measurement of the Z^0 Mass and Width with the OPAL Detector at LEP, *Phys. Lett. B* 231 (1989) 530–538. doi:10.1016/0370-2693(89)90705-3.
- [22] V. M. Abazov, et al., Combination of CDF and D0 Results on W Boson Mass and Width, *Phys. Rev. D* 70 (2004) 092008. arXiv:hep-ex/0311039, doi:10.1103/PhysRevD.70.092008.
- [23] S. Schael, et al., Measurement of the W boson mass and width in e^+e^- collisions at LEP, *Eur. Phys. J. C* 47 (2006) 309–335. arXiv:hep-ex/0605011, doi:10.1140/epjc/s2006-02576-8.
- [24] P. Bagnaia, et al., Evidence for $Z^0 \rightarrow e^+e^-$ at the CERN $\bar{p}p$ Collider, *Phys. Lett. B* 129 (1983) 130–140. doi:10.1016/0370-2693(83)90744-X.
- [25] M. Kobayashi, T. Maskawa, CP-Violation in the Renormalizable Theory of Weak Interaction, *Progress of Theoretical Physics* 49 (2) (1973) 652–657. arXiv:<https://academic.oup.com/ptp/article-pdf/49/2/652/5257692/49-2-652.pdf>, doi:10.1143/PTP.49.652.
URL <https://doi.org/10.1143/PTP.49.652>
- [26] L.-L. Chau, W.-Y. Keung, Comments on the Parametrization of the Kobayashi-Maskawa Matrix, *Phys. Rev. Lett.* 53 (1984) 1802. doi:10.1103/PhysRevLett.53.1802.

- [27] R. Aaij, et al., Measurement of $|V_{cb}|$ with $B_s^0 \rightarrow D_s^{(*)-} \mu^+ \nu_\mu$ decays, Phys. Rev. D 101 (2020) 072004. doi:10.1103/PhysRevD.101.072004.
 URL <https://link.aps.org/doi/10.1103/PhysRevD.101.072004>
- [28] Y. Amhis, et al., Averages of b-hadron, c-hadron, and τ -lepton properties as of 2018, The European Physical Journal C 81 (3) (mar 2021). doi:10.1140/epjc/s10052-020-8156-7.
 URL <https://doi.org/10.1140/epjc/s10052-020-8156-7>
- [29] C. G. Boyd, B. Grinstein, R. F. Lebed, Model independent determinations of anti-B \rightarrow D (lepton), D^* (lepton) anti-neutrino form-factors, Nucl. Phys. B 461 (1996) 493–511. arXiv:hep-ph/9508211, doi:10.1016/0550-3213(95)00653-2.
- [30] F. U. Bernlochner, Z. Ligeti, M. Papucci, D. J. Robinson, Tensions and correlations in $|V_{cb}|$ determinations, Phys. Rev. D 96 (2017) 091503. doi:10.1103/PhysRevD.96.091503.
 URL <https://link.aps.org/doi/10.1103/PhysRevD.96.091503>
- [31] C. Bourrely, L. Lellouch, I. Caprini, Model-independent description of $b \rightarrow \pi l \nu$ decays and a determination of $|V_{ub}|$, Phys. Rev. D 79 (2009) 013008. doi:10.1103/PhysRevD.79.013008.
 URL <https://link.aps.org/doi/10.1103/PhysRevD.79.013008>
- [32] C. G. Boyd, B. Grinstein, R. F. Lebed, Precision corrections to dispersive bounds on form factors, Phys. Rev. D 56 (1997) 6895–6911. doi:10.1103/PhysRevD.56.6895.
 URL <https://link.aps.org/doi/10.1103/PhysRevD.56.6895>
- [33] C. G. Boyd, B. Grinstein, R. F. Lebed, Constraints on form factors for exclusive semileptonic heavy to light meson decays, Phys. Rev. Lett. 74 (1995) 4603–4606. doi:10.1103/PhysRevLett.74.4603.
 URL <https://link.aps.org/doi/10.1103/PhysRevLett.74.4603>
- [34] I. Caprini, L. Lellouch, M. Neubert, Dispersive bounds on the shape of anti-B \rightarrow $D^{(*)}$ lepton anti-neutrino form-factors, Nucl. Phys. B 530 (1998) 153–181. arXiv:hep-ph/9712417, doi:10.1016/S0550-3213(98)00350-2.

- [35] G. Ricciardi, Theory: Semileptonic B Decays and $|V_{xb}|$ update, PoS BEAUTY2020 (2021) 031. [arXiv:2103.06099](https://arxiv.org/abs/2103.06099), doi:10.22323/1.391.0031.
- [36] Y. Zhang, T. Zhong, H.-B. Fu, W. Cheng, L. Zeng, X.-G. Wu, Ratio R(Ds) for $B_s \rightarrow D_s \ell \nu \ell$ by using the QCD light-cone sum rules within the framework of heavy quark effective field theory, Phys. Rev. D 105 (9) (2022) 096013. [arXiv:2202.02730](https://arxiv.org/abs/2202.02730), doi:10.1103/PhysRevD.105.096013.
- [37] J. Harrison, C. T. H. Davies, M. Wingate, Lattice qcd calculation of the $B_{(s)} \rightarrow D_{(s)}^* \ell \nu$ form factors at zero recoil and implications for $|V_{cb}|$, Phys. Rev. D 97 (2018) 054502. doi:10.1103/PhysRevD.97.054502.
URL <https://link.aps.org/doi/10.1103/PhysRevD.97.054502>
- [38] C. J. Monahan, H. Na, C. M. Bouchard, G. P. Lepage, J. Shigemitsu, $B_s \rightarrow D_s \ell \nu$ form factors and the fragmentation fraction ratio f_s/f_d , Phys. Rev. D 95 (2017) 114506. doi:10.1103/PhysRevD.95.114506.
URL <https://link.aps.org/doi/10.1103/PhysRevD.95.114506>
- [39] A. G. Akeroyd, S. Recksiegel, Direct CP asymmetry of $B \rightarrow X(d,s)$ gamma in a model with vector quarks, Phys. Lett. B 525 (2002) 81–88. [arXiv:hep-ph/0109091](https://arxiv.org/abs/hep-ph/0109091), doi:10.1016/S0370-2693(01)01419-8.
- [40] A. Ali, C. Greub, Rare decays $B \rightarrow X(d)$ gamma in the standard model, Phys. Lett. B 287 (1992) 191–202. doi:10.1016/0370-2693(92)91899-K.
- [41] B.-Y. Cui, Y.-K. Huang, Y.-L. Shen, C. Wang, Y.-M. Wang, Precision calculations of $B_{d,s} \rightarrow \pi$, K decay form factors in soft-collinear effective theory, JHEP 03 (2023) 140. [arXiv:2212.11624](https://arxiv.org/abs/2212.11624), doi:10.1007/JHEP03(2023)140.
- [42] R. Aaij, et al., First observation of the decay $B^+ \rightarrow \pi^+ \mu^+ \mu^-$, JHEP 12 (2012) 125. [arXiv:1210.2645](https://arxiv.org/abs/1210.2645), doi:10.1007/JHEP12(2012)125.
- [43] J. T. Wei, et al., Search for $B \rightarrow \pi l^+ l^-$ Decays at Belle, Phys. Rev. D 78 (2008) 011101. [arXiv:0804.3656](https://arxiv.org/abs/0804.3656), doi:10.1103/PhysRevD.78.011101.
- [44] C. Belén, et al., Measurement of the branching fraction of the decay $B^+ \rightarrow \pi^+ \pi^- \ell^+ \nu_\ell$ in fully reconstructed events at Belle, Phys. Rev. D 103 (11) (2021) 112001. [arXiv:2005.07766](https://arxiv.org/abs/2005.07766), doi:10.1103/PhysRevD.103.112001.

- [45] I. Parnova, A. Ali, A. Y. Parkhomenko, Subleading contributions in rare semileptonic $B^+ \rightarrow \pi^+\ell^+\ell^-$ decay, PoS PANIC2021 (2022) 158. doi:10.22323/1.380.0158.
- [46] H.-M. Choi, Exclusive Rare $B_s \rightarrow (K, \eta, \eta')\ell^+\ell^-$ Decays in the Light-Front Quark Model, J. Phys. G 37 (2010) 085005. arXiv:1002.0721, doi:10.1088/0954-3899/37/8/085005.
- [47] C.-D. Lü, Y.-L. Shen, Y.-M. Wang, Y.-B. Wei, QCD calculations of $B \rightarrow \pi, K$ form factors with higher-twist corrections, JHEP 01 (2019) 024. arXiv:1810.00819, doi:10.1007/JHEP01(2019)024.
- [48] A. Khodjamirian, A. V. Rusov, $B_s \rightarrow K\ell\nu_\ell$ and $B_{(s)} \rightarrow \pi(K)\ell^+\ell^-$ decays at large recoil and CKM matrix elements, JHEP 08 (2017) 112. arXiv:1703.04765, doi:10.1007/JHEP08(2017)112.
- [49] N. Gubernari, A. Kokulu, D. van Dyk, $B \rightarrow P$ and $B \rightarrow V$ Form Factors from B -Meson Light-Cone Sum Rules beyond Leading Twist, JHEP 01 (2019) 150. arXiv:1811.00983, doi:10.1007/JHEP01(2019)150.
- [50] C. Hambrock, A. Khodjamirian, A. Rusov, Hadronic effects and observables in $B \rightarrow \pi\ell^+\ell^-$ decay at large recoil, Phys. Rev. D 92 (7) (2015) 074020. arXiv:1506.07760, doi:10.1103/PhysRevD.92.074020.
- [51] W. Cheng, X.-G. Wu, R.-Y. Zhou, H.-B. Fu, The $B \rightarrow \rho$ helicity form factors within the QCD light-cone sum rules, Phys. Rev. D 98 (9) (2018) 096013. arXiv:1808.10775, doi:10.1103/PhysRevD.98.096013.
- [52] M. Okamoto, et al., Semileptonic $D \rightarrow \pi/K$ and $B \rightarrow \pi/D$ decays in 2+1 flavor lattice QCD, Nucl. Phys. B Proc. Suppl. 140 (2005) 461–463. arXiv:hep-lat/0409116, doi:10.1016/j.nuclphysbps.2004.11.151.
- [53] E. Dalgic, A. Gray, M. Wingate, C. T. H. Davies, G. P. Lepage, J. Shigemitsu, B meson semileptonic form-factors from unquenched lattice QCD, Phys. Rev. D 73 (2006) 074502, [Erratum: Phys. Rev. D 75, 119906 (2007)]. arXiv:hep-lat/0601021, doi:10.1103/PhysRevD.75.119906.

- [54] J. A. Bailey, et al., The $B \rightarrow \pi \ell \nu$ semileptonic form factor from three-flavor lattice QCD: A Model-independent determination of $|V_{ub}|$, Phys. Rev. D 79 (2009) 054507. [arXiv:0811.3640](https://arxiv.org/abs/0811.3640), doi:10.1103/PhysRevD.79.054507.
- [55] A. Biswas, S. Nandi, S. K. Patra, I. Ray, Study of the $b \rightarrow d\ell\ell$ transitions in the Standard Model and test of New Physics sensitivities, JHEP 03 (2023) 247. [arXiv:2208.14463](https://arxiv.org/abs/2208.14463), doi:10.1007/JHEP03(2023)247.
- [56] K. Inaoka, H. Koyama, Cp-violation in eta, k-neutral decays and the delta-i=2 selection rule, Prog. Theor. Phys. 39 (1968) 751–767. doi:10.1143/PTP.39.751.
- [57] S. Y. Tsai, A phenomenological study of violation of pc invariance in decays of k-neutral and antikaon-neutral, Prog. Theor. Phys. 40 (1968) 1414–1423. doi:10.1143/PTP.40.1414.
- [58] S. K. Yun, Possible origin of the cp violation in the neutral k system, Phys. Rev. 178 (1969) 2439–2442. doi:10.1103/PhysRev.178.2439.
- [59] E. A. Paschos, QUARK MIXING AND CP VIOLATION, in: 21st Rencontres de Moriond: Perspectives in Electroweak Interactions and Unified Theories, 1986, pp. 347–360.
- [60] R. C. Kennicutt, Jr., A spectrophotometric atlas of galaxies, Astrophys. J. Suppl. 79 (1992) 255–284. doi:10.1086/191653.
- [61] S. Conetti, LHC-B: A dedicated beauty experiment for precision measurements of CP-violation, Nucl. Phys. B Proc. Suppl. 55 (1997) 286–293. doi:10.1016/S0920-5632(98)00107-8.
- [62] M. L. Perl, et al., Evidence for Anomalous Lepton Production in $e^+ - e^-$ Annihilation, Phys. Rev. Lett. 35 (1975) 1489–1492. doi:10.1103/PhysRevLett.35.1489.
- [63] P. Giubellino, The ALICE detector at LHC, Nucl. Instrum. Meth. A 344 (1994) 27–38. doi:10.1016/0168-9002(94)90647-5.

- [64] M. A. Moinester, V. Steiner, Primakoff physics for CERN COMPASS hadron beam: Hadron polarizabilities, hybrid mesons, chiral anomaly, meson radiative transitions, 1997. [arXiv:hep-ex/9801011](https://arxiv.org/abs/hep-ex/9801011).
- [65] M. Moinester, S. U. Chung, Hybrid meson structure at COMPASS, in: COMPASS Week, 1999. [arXiv:hep-ex/0003008](https://arxiv.org/abs/hep-ex/0003008).
- [66] C. P. Schuetz, The Dirac experiment at CERN, in: 38th Rencontres de Moriond on QCD and High-Energy Hadronic Interactions, 2003. [arXiv:hep-ph/0305121](https://arxiv.org/abs/hep-ph/0305121).
- [67] W. Bertsche, et al., The ALPHA experiment: A cold antihydrogen trap, AIP Conf. Proc. 796 (1) (2005) 301–308. doi:[10.1063/1.2130184](https://doi.org/10.1063/1.2130184).
- [68] N. Abgrall, Hadroproduction measurements: The NA61-SHINE experiment at CERN, AIP Conf. Proc. 981 (1) (2008) 157–159. doi:[10.1063/1.2898917](https://doi.org/10.1063/1.2898917).
- [69] S. Balev, Measurement of $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ at NA62 Experiment, eConf C080625 (2008) 0014.
- [70] Y. Gornushkin, Study of Tau Neutrino Production in NA65 Experiment at CERN SPS, Moscow Univ. Phys. Bull. 77 (2) (2022) 363–364. doi:[10.3103/S0027134922020370](https://doi.org/10.3103/S0027134922020370).
- [71] J. Morfin, RESULTS OF THE NEUTRAL CURRENT SEARCH IN GARGAMELLE., in: 2nd Summer Institute on Particle Interactions at Very High Energies: Duality in Elementary Particle Physics, 1973. doi:[10.1007/978-1-4684-2826-1_6](https://doi.org/10.1007/978-1-4684-2826-1_6).
- [72] G. b. Lubkin, P ANTI-P COLLISIONS YIELD INTERMEDIATE BOSON AT 80-GEV, AS PREDICTED, Phys. Today 36N4 (1983) 17–20.
- [73] A. L. Robinson, CERN VECTOR BOSON HUNT SUCCESSFUL, Science 221 (1983) 840–842. doi:[10.1126/science.221.4613.840](https://doi.org/10.1126/science.221.4613.840).
- [74] F. Abe, et al., Observation of top quark production in $\bar{p}p$ collisions, Phys. Rev. Lett. 74 (1995) 2626–2631. [arXiv:hep-ex/9503002](https://arxiv.org/abs/hep-ex/9503002), doi:[10.1103/PhysRevLett.74.2626](https://doi.org/10.1103/PhysRevLett.74.2626).

- [75] R. Y. Peters, Top Quark Mass Measurements at the Tevatron, PoS EPS-HEP2013 (2013) 243. [arXiv:1309.5783](https://arxiv.org/abs/1309.5783), doi:10.22323/1.180.0243.
- [76] J. S. Russ, SELEX - Hadroproduction of charm baryons out to large $x(F)$, Nucl. Phys. B Proc. Suppl. 50 (1996) 248–254. doi:10.1016/0920-5632(96)00399-4.
- [77] S. Carrillo, F. Vazquez, FOCUS: A charm photo-production experiment at Fermilab, AIP Conf. Proc. 422 (1) (1998) 377–381. doi:10.1063/1.55075.
- [78] P. Senger, Strangeness and charm of compressed baryonic matter: The CBM experiment at FAIR, J. Phys. G 31 (2005) S1111–S1114. doi:10.1088/0954-3899/31/6/069.
- [79] V. Friese, The CBM experiment at GSI/FAIR, Nucl. Phys. A 774 (2006) 377–386. doi:10.1016/j.nuclphysa.2006.06.018.
- [80] T. R. Saitoh, J. Pochodzalla, Hypernuclear physics projects with PANDA at GSI, Acta Phys. Polon. B 35 (2004) 1033–1041.
- [81] R. Krucken, The NuSTAR facility at FAIR, J. Phys. G 31 (2005) S1807–S1811. doi:10.1088/0954-3899/31/10/077.
- [82] V. Luth, BaBar: The Detector for the PEP-II B Factory at SLAC, in: 1994 Meeting of the American Physical Society, Division of Particles and Fields (DPF 94), 1994, pp. 1898–1903.
- [83] M. T. Cheng, et al., A Study of CP violation in B meson decays: Technical design report (3 1995).
- [84] L. Li, Open charm physics at BESIII, Nucl. Phys. B Proc. Suppl. 225-227 (2012) 107–111. doi:10.1016/j.nuclphysbps.2012.02.024.
- [85] Y. Huang, New observations on light hadron spectroscopy at BESIII, PoS ICHEP2010 (2010) 181. doi:10.22323/1.120.0181.
- [86] H. Jiang, Search for invisible decays at BESIII, PoS ICHEP2022 (2022) 156. doi:10.22323/1.414.0156.

- [87] V. Bertacchi, Recent results from Belle II, Nucl. Part. Phys. Proc. 324-329 (2023) 107–112. doi:10.1016/j.nuclphysbps.2023.01.022.
- [88] L. Novotny, Measurements of CP violation and lifetime properties of B mesons at ATLAS, PoS CKM2021 (2023) 104. doi:10.22323/1.411.0104.
- [89] R. Aaij, et al., Search for CP violation in $D_{(s)}^+ \rightarrow K^-K^+K^+$ decays, JHEP 07 (2023) 067. arXiv:2303.04062, doi:10.1007/JHEP07(2023)067.
- [90] W. Krupa, Recent Results of Measurement of CKM Angle γ and CPV in the Beauty Sector at the LHCb, Acta Phys. Polon. Supp. 16 (3) (2023) 6. doi:10.5506/APhysPolBSupp.16.3-A6.
- [91] G. Aad, et al., Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC, Phys. Lett. B 716 (2012) 1–29. arXiv:1207.7214, doi:10.1016/j.physletb.2012.08.020.
- [92] C. Aubin, et al., Charmed-meson decay constants in three-flavor lattice qcd, Phys. Rev. Lett. 95 (2005) 122002. doi:10.1103/PhysRevLett.95.122002.
URL <https://link.aps.org/doi/10.1103/PhysRevLett.95.122002>
- [93] E. Follana, C. T. H. Davies, G. P. Lepage, J. Shigemitsu, High-precision determination of the π , k , d , and D_s decay constants from lattice qcd, Phys. Rev. Lett. 100 (2008) 062002. doi:10.1103/PhysRevLett.100.062002.
URL <https://link.aps.org/doi/10.1103/PhysRevLett.100.062002>
- [94] C. T. H. Davies, C. McNeile, E. Follana, G. P. Lepage, H. Na, J. Shigemitsu, Update: Precision D_s decay constant from full lattice qcd using very fine lattices, Phys. Rev. D 82 (2010) 114504. doi:10.1103/PhysRevD.82.114504.
URL <https://link.aps.org/doi/10.1103/PhysRevD.82.114504>
- [95] A. Bazavov, et al., b - and d -meson decay constants from three-flavor lattice qcd, Phys. Rev. D 85 (2012) 114506. doi:10.1103/PhysRevD.85.114506.
URL <https://link.aps.org/doi/10.1103/PhysRevD.85.114506>
- [96] N. Carrasco, et al., B-physics from $nf = 2$ tmQCD: the standard model and beyond, Journal of High Energy Physics 2014 (3) (mar 2014).

doi:10.1007/jhep03(2014)016.

URL <https://doi.org/10.1007%2Fjhep03%282014%29016>

- [97] A. Bazavov, et al., *b*- and *d*-meson leptonic decay constants from four-flavor lattice qcd, Phys. Rev. D 98 (2018) 074512. doi:10.1103/PhysRevD.98.074512.
URL <https://link.aps.org/doi/10.1103/PhysRevD.98.074512>
- [98] E. Gámiz, C. T. H. Davies, G. P. Lepage, J. Shigemitsu, M. Wingate, Neutral *b* meson mixing in unquenched lattice qcd, Phys. Rev. D 80 (2009) 014503.
doi:10.1103/PhysRevD.80.014503.
URL <https://link.aps.org/doi/10.1103/PhysRevD.80.014503>
- [99] H. Na, C. J. Monahan, C. T. H. Davies, R. Horgan, G. P. Lepage, J. Shigemitsu, *b* and *B_s* meson decay constants from lattice qcd, Phys. Rev. D 86 (2012) 034506.
doi:10.1103/PhysRevD.86.034506.
URL <https://link.aps.org/doi/10.1103/PhysRevD.86.034506>
- [100] C. McNeile, et al., Heavy meson masses and decay constants from relativistic heavy quarks in full lattice qcd, Phys. Rev. D 86 (2012) 074503.
doi:10.1103/PhysRevD.86.074503.
URL <https://link.aps.org/doi/10.1103/PhysRevD.86.074503>
- [101] B. Colquhoun, C. T. H. Davies, J. Kettle, J. Koponen, A. T. Lytle, R. J. Dowdall, G. P. Lepage, *b*-meson decay constants: A more complete picture from full lattice qcd, Phys. Rev. D 91 (2015) 114509.
doi:10.1103/PhysRevD.91.114509.
URL <https://link.aps.org/doi/10.1103/PhysRevD.91.114509>
- [102] C. Bouchard, G. P. Lepage, C. Monahan, H. Na, J. Shigemitsu, Rare decay *b* → *k* $\ell^+\ell^-$ form factors from lattice qcd, Phys. Rev. D 88 (2013) 054509.
doi:10.1103/PhysRevD.88.054509.
URL <https://link.aps.org/doi/10.1103/PhysRevD.88.054509>
- [103] J. A. Bailey, et al., *b* → *kl⁺l⁻* decay form factors from three-flavor lattice qcd, Phys. Rev. D 93 (2016) 025026. doi:10.1103/PhysRevD.93.025026.
URL <https://link.aps.org/doi/10.1103/PhysRevD.93.025026>

- [104] R. J. Dowdall, C. T. H. Davies, R. R. Horgan, G. P. Lepage, C. J. Monahan, J. Shigemitsu, M. Wingate, Neutral B-meson mixing from full lattice QCD at the physical point, Phys. Rev. D 100 (9) (2019) 094508. [arXiv:1907.01025](https://arxiv.org/abs/1907.01025), doi:10.1103/PhysRevD.100.094508.
- [105] Y.-Q. Peng, M.-Z. Yang, Study of semileptonic decay of in qcd sum rule, Communications in Theoretical Physics 72 (9) (2020) 095201. doi:10.1088/1572-9494/aba244.
URL <https://dx.doi.org/10.1088/1572-9494/aba244>
- [106] V. V. Kiselev, A. K. Likhoded, A. I. Onishchenko, Semileptonic B_c meson decays in sum rules of QCD and NRQCD, Nucl. Phys. B 569 (2000) 473–504. [arXiv:hep-ph/9905359](https://arxiv.org/abs/hep-ph/9905359), doi:10.1016/S0550-3213(99)00505-2.
- [107] N. R. Soni, B. R. Joshi, R. P. Shah, H. R. Chauhan, J. N. Pandya, $Q\bar{Q}$ ($Q \in \{b, c\}$) spectroscopy using the Cornell potential, Eur. Phys. J. C 78 (7) (2018) 592. [arXiv:1707.07144](https://arxiv.org/abs/1707.07144), doi:10.1140/epjc/s10052-018-6068-6.
- [108] N. Soni, J. Patel, A. Gadaria, J. Pandya, Decay properties of heavy quarkonia employing Cornell potential, in: 65th DAE BRNS Symposium on nuclear physics, 2022.
- [109] E. Eichten, K. Gottfried, T. Kinoshita, K. D. Lane, T.-M. Yan, Charmonium: The Model, Phys. Rev. D 17 (1978) 3090, [Erratum: Phys.Rev.D 21, 313 (1980)]. doi:10.1103/PhysRevD.17.3090.
- [110] A. Martin, A FIT of Upsilon and Charmonium Spectra, Phys. Lett. B 93 (1980) 338–342. doi:10.1016/0370-2693(80)90527-4.
- [111] P. C. Vinodkumar, J. N. Pandya, V. M. Bannur, S. B. Khadkikar, A unified scheme for flavoured mesons and baryons, Eur. Phys. J. A 4 (1999) 83–90. doi:10.1007/s100500050206.
- [112] S. B. Khadkikar, RELATIVISTIC HARMONIC CONFINEMENT OF QUARKS AND GLUONS, Pramana 24 (1985) 63–68. doi:10.1007/BF02894818.

- [113] S. Scherer, M. R. Schindler, Chiral perturbation theory, 2022.
`arXiv:2211.17031`.
- [114] A. G. Grozin, Introduction to the heavy quark effective theory. part 1 (12 1992).
`arXiv:hep-ph/9908366`.
- [115] T. Branz, A. Faessler, T. Gutsche, M. A. Ivanov, J. G. Korner, V. E. Lyubovitskij, Relativistic constituent quark model with infrared confinement, Phys. Rev. D 81 (2010) 034010. `arXiv:0912.3710`, doi:10.1103/PhysRevD.81.034010.
- [116] G. V. Efimov, M. A. Ivanov, Confinement and quark structure of light hadrons, Int. J. Mod. Phys. A 4 (8) (1989) 2031–2060. doi:10.1142/S0217751X89000832.
- [117] G. V. Efimov, M. A. Ivanov, The Quark confinement model of hadrons, IOP, Bristol, 1993.
- [118] A. W. Thomas, W. Weise, The Structure of the Nucleon, Wiley, Germany, 2001. doi:10.1002/352760314X.
- [119] A. Khodjamirian, T. Mannel, N. Offen, Form-factors from light-cone sum rules with B-meson distribution amplitudes, Phys. Rev. D 75 (2007) 054013. `arXiv:hep-ph/0611193`, doi:10.1103/PhysRevD.75.054013.
- [120] B. Grinstein, D. Pirjol, Exclusive rare $B \rightarrow K^*\ell^+\ell^-$ decays at low recoil: Controlling the long-distance effects, Phys. Rev. D 70 (2004) 114005. `arXiv:hep-ph/0404250`, doi:10.1103/PhysRevD.70.114005.
- [121] C. D. Roberts, S. M. Schmidt, Dyson-Schwinger equations: Density, temperature and continuum strong QCD, Prog. Part. Nucl. Phys. 45 (2000) S1–S103. `arXiv:nucl-th/0005064`, doi:10.1016/S0146-6410(00)90011-5.
- [122] D. Melikhov, N. Nikitin, S. Simula, Rare exclusive semileptonic $b \rightarrow s$ transitions in the standard model, Phys. Rev. D 57 (1998) 6814–6828. `arXiv:hep-ph/9711362`, doi:10.1103/PhysRevD.57.6814.
- [123] D. Ebert, R. N. Faustov, V. O. Galkin, New analysis of semileptonic B decays in the relativistic quark model, Phys. Rev. D 75 (2007) 074008. `arXiv:hep-ph/0611307`, doi:10.1103/PhysRevD.75.074008.

- [124] M. Ladisa, G. Nardulli, P. Santorelli, Semileptonic and rare B meson decays into a light pseudoscalar meson, Phys. Lett. B 455 (1999) 283–290. [arXiv:hep-ph/9903206](https://arxiv.org/abs/hep-ph/9903206), doi:10.1016/S0370-2693(99)00386-X.
- [125] P. Colangelo, P. Santorelli, Dependence of the form-factors of $B \rightarrow \pi$ lepton neutrino on the heavy quark mass, Phys. Lett. B 327 (1994) 123–128. [arXiv:hep-ph/9312258](https://arxiv.org/abs/hep-ph/9312258), doi:10.1016/0370-2693(94)91539-3.
- [126] A. Faessler, T. Gutsche, M. A. Ivanov, J. G. Korner, V. E. Lyubovitskij, The Exclusive rare decays $B \rightarrow K(K^*) \bar{\ell}\ell$ and $B_c \rightarrow D(D^*) \bar{\ell}\ell$ in a relativistic quark model, Eur. Phys. J. direct 4 (1) (2002) 18. [arXiv:hep-ph/0205287](https://arxiv.org/abs/hep-ph/0205287), doi:10.1007/s1010502c0018.
- [127] M. A. Ivanov, J. G. Korner, P. Santorelli, Semileptonic decays of B_c mesons into charmonium states in a relativistic quark model, Phys. Rev. D 71 (2005) 094006, [Erratum: Phys.Rev.D 75, 019901 (2007)]. [arXiv:hep-ph/0501051](https://arxiv.org/abs/hep-ph/0501051), doi:10.1103/PhysRevD.75.019901.
- [128] M. A. Ivanov, J. G. Korner, O. N. Pakhomova, The Nonleptonic decays $B_c^+ \rightarrow D_s^+ \bar{D}^0$ and $B_c^+ \rightarrow D_s^+ D^0$ in a relativistic quark model, Phys. Lett. B 555 (2003) 189–196. [arXiv:hep-ph/0212291](https://arxiv.org/abs/hep-ph/0212291), doi:10.1016/S0370-2693(03)00052-2.
- [129] A. Salam, Lagrangian theory of composite particles, Nuovo Cim. 25 (1962) 224–227. doi:10.1007/BF02733330.
- [130] S. Weinberg, Elementary particle theory of composite particles, Phys. Rev. 130 (1963) 776–783. doi:10.1103/PhysRev.130.776.
- [131] M. A. Ivanov, P. Santorelli, N. Tancredi, The Semileptonic form-factors of B and D mesons in the quark confinement model, Eur. Phys. J. A 9 (2000) 109–114. [arXiv:hep-ph/9905209](https://arxiv.org/abs/hep-ph/9905209), doi:10.1007/s100500070060.
- [132] M. A. Ivanov, P. Santorelli, Leptonic and semileptonic decays of pseudoscalar mesons, Phys. Lett. B 456 (1999) 248–255. [arXiv:hep-ph/9903446](https://arxiv.org/abs/hep-ph/9903446), doi:10.1016/S0370-2693(99)00474-8.

- [133] M. A. Ivanov, M. P. Locher, V. E. Lyubovitskij, Electromagnetic form-factors of nucleons in a relativistic three quark model, *Few Body Syst.* 21 (1996) 131. [arXiv:hep-ph/9602372](https://arxiv.org/abs/hep-ph/9602372), doi:10.1007/s006010050044.
- [134] A. Faessler, T. Gutsche, M. A. Ivanov, J. G. Korner, V. E. Lyubovitskij, Semileptonic decays of double heavy baryons, *Phys. Lett. B* 518 (2001) 55–62. [arXiv:hep-ph/0107205](https://arxiv.org/abs/hep-ph/0107205), doi:10.1016/S0370-2693(01)01024-3.
- [135] C. Anastasiou, E. W. N. Glover, C. Oleari, M. E. Tejeda-Yeomans, Two-loop QCD corrections to the scattering of massless distinct quarks, *Nucl. Phys. B* 601 (2001) 318–340. [arXiv:hep-ph/0010212](https://arxiv.org/abs/hep-ph/0010212), doi:10.1016/S0550-3213(01)00079-7.
- [136] J. A. M. Vermaasen, The FORM project, *Nucl. Phys. B Proc. Suppl.* 183 (2008) 19–24. [arXiv:0806.4080](https://arxiv.org/abs/0806.4080), doi:10.1016/j.nuclphysbps.2008.09.076.
- [137] R. P. Feynman, Space - time approach to quantum electrodynamics, *Phys. Rev.* 76 (1949) 769–789. doi:10.1103/PhysRev.76.769.
- [138] J. Albrecht, D. van Dyk, C. Langenbruch, Flavour anomalies in heavy quark decays, *Prog. Part. Nucl. Phys.* 120 (2021) 103885. [arXiv:2107.04822](https://arxiv.org/abs/2107.04822), doi:10.1016/j.ppnp.2021.103885.
- [139] R. Aaij, et al., Measurement of $|V_{cb}|$ with $B_s^0 \rightarrow D_s^{(*)-} \mu^+ \nu_\mu$ decays, *Phys. Rev. D* 101 (7) (2020) 072004. [arXiv:2001.03225](https://arxiv.org/abs/2001.03225), doi:10.1103/PhysRevD.101.072004.
- [140] E. McLean, C. T. H. Davies, J. Koponen, A. T. Lytle, $B_s \rightarrow D_s \ell \nu$ Form Factors for the full q^2 range from Lattice QCD with non-perturbatively normalized currents, *Phys. Rev. D* 101 (7) (2020) 074513. [arXiv:1906.00701](https://arxiv.org/abs/1906.00701), doi:10.1103/PhysRevD.101.074513.
- [141] J. Harrison, C. T. H. Davies, $B_s \rightarrow D_s^*$ form factors for the full q^2 range from lattice QCD, *Phys. Rev. D* 105 (9) (2022) 094506. [arXiv:2105.11433](https://arxiv.org/abs/2105.11433), doi:10.1103/PhysRevD.105.094506.

- [142] X.-Q. Hu, S.-P. Jin, Z.-J. Xiao, Semileptonic decays $B/B_s \rightarrow (D^{(*)}, D_s^{(*)})l\nu_l$ in the PQCD approach with the lattice QCD input, Chin. Phys. C 44 (5) (2020) 053102. [arXiv:1912.03981](https://arxiv.org/abs/1912.03981), doi:10.1088/1674-1137/44/5/053102.
- [143] Y. Zhang, T. Zhong, H.-B. Fu, W. Cheng, X.-G. Wu, Ds-meson leading-twist distribution amplitude within the QCD sum rules and its application to the $B_s \rightarrow D_s$ transition form factor, Phys. Rev. D 103 (11) (2021) 114024. [arXiv:2104.00180](https://arxiv.org/abs/2104.00180), doi:10.1103/PhysRevD.103.114024.
- [144] M. Bordone, N. Gubernari, D. van Dyk, M. Jung, Heavy-Quark expansion for $\bar{B}_s \rightarrow D_s^{(*)}$ form factors and unitarity bounds beyond the $SU(3)_F$ limit, Eur. Phys. J. C 80 (4) (2020) 347. [arXiv:1912.09335](https://arxiv.org/abs/1912.09335), doi:10.1140/epjc/s10052-020-7850-9.
- [145] R.-H. Li, C.-D. Lu, Y.-M. Wang, Exclusive B_s decays to the charmed mesons $D_s^+(1968, 2317)$ in the standard model, Phys. Rev. D 80 (2009) 014005. [arXiv:0905.3259](https://arxiv.org/abs/0905.3259), doi:10.1103/PhysRevD.80.014005.
- [146] P. Blasi, P. Colangelo, G. Nardulli, N. Paver, Phenomenology of $B(s)$ decays, Phys. Rev. D 49 (1994) 238–246. [arXiv:hep-ph/9307290](https://arxiv.org/abs/hep-ph/9307290), doi:10.1103/PhysRevD.49.238.
- [147] X.-W. Kang, T. Luo, Y. Zhang, L.-Y. Dai, C. Wang, Semileptonic B and B_s decays involving scalar and axial-vector mesons, Eur. Phys. J. C 78 (11) (2018) 909. [arXiv:1808.02432](https://arxiv.org/abs/1808.02432), doi:10.1140/epjc/s10052-018-6385-9.
- [148] G. Li, F.-l. Shao, W. Wang, $B_s \rightarrow D_s(3040)$ form factors and B_s decays into $D_s(3040)$, Phys. Rev. D 82 (2010) 094031. [arXiv:1008.3696](https://arxiv.org/abs/1008.3696), doi:10.1103/PhysRevD.82.094031.
- [149] R. N. Faustov, V. O. Galkin, Weak decays of B_s mesons to D_s mesons in the relativistic quark model, Phys. Rev. D 87 (3) (2013) 034033. [arXiv:1212.3167](https://arxiv.org/abs/1212.3167), doi:10.1103/PhysRevD.87.034033.
- [150] S.-M. Zhao, X. Liu, S.-J. Li, Study on $B(s) \rightarrow D(sJ)$ (2317, 2460) 1 anti-nu Semileptonic Decays in the CQM Model, Eur. Phys. J. C 51 (2007) 601–606. [arXiv:hep-ph/0612008](https://arxiv.org/abs/hep-ph/0612008), doi:10.1140/epjc/s10052-007-0322-7.

- [151] K. Adamczyk, et al., Determination of $|V_{ub}|$ from untagged $B^0 \rightarrow \pi^- \ell^+ \nu_\ell$ decays using 2019-2021 Belle II data (10 2022). [arXiv:2210.04224](https://arxiv.org/abs/2210.04224).
- [152] Y. S. Amhis, et al., Averages of b-hadron, c-hadron, and τ -lepton properties as of 2021, Phys. Rev. D 107 (5) (2023) 052008. [arXiv:2206.07501](https://arxiv.org/abs/2206.07501), doi:10.1103/PhysRevD.107.052008.
- [153] S. B. Athar, et al., Study of the q^{**2} dependence of $B \rightarrow \pi l \nu$ and $B \rightarrow \rho \nu$ (ω) $l \nu$ decay and extraction of $-V(u b)$, Phys. Rev. D 68 (2003) 072003. [arXiv:hep-ex/0304019](https://arxiv.org/abs/hep-ex/0304019), doi:10.1103/PhysRevD.68.072003.
- [154] N. E. Adam, et al., A Study of Exclusive Charmless Semileptonic B Decay and $-V(ub)$, Phys. Rev. Lett. 99 (2007) 041802. [arXiv:hep-ex/0703041](https://arxiv.org/abs/hep-ex/0703041), doi:10.1103/PhysRevLett.99.041802.
- [155] R. Gray, et al., A Study of Exclusive Charmless Semileptonic B Decays and Extraction of $|V_{ub}|$ at CLEO, Phys. Rev. D 76 (2007) 012007, [Addendum: Phys.Rev.D 76, 039901 (2007)]. [arXiv:hep-ex/0703042](https://arxiv.org/abs/hep-ex/0703042), doi:10.1103/PhysRevD.76.012007.
- [156] B. Aubert, et al., Study of $B \rightarrow \pi \ell \nu$ and $B \rightarrow \rho \ell \nu$ decays and determination of $|V_{ub}|$, Phys. Rev. D 72 (2005) 051102. [arXiv:hep-ex/0507003](https://arxiv.org/abs/hep-ex/0507003), doi:10.1103/PhysRevD.72.051102.
- [157] P. del Amo Sanchez, et al., Study of $B \rightarrow \pi \ell \nu$ and $B \rightarrow \rho \ell \nu$ Decays and Determination of $|V_{ub}|$, Phys. Rev. D 83 (2011) 032007. [arXiv:1005.3288](https://arxiv.org/abs/1005.3288), doi:10.1103/PhysRevD.83.032007.
- [158] P. del Amo Sanchez, et al., Measurement of the $B^0 \rightarrow \pi^\ell \ell^+ \nu$ and $B^+ \rightarrow \eta^{(\prime)} \ell^+ \nu$ Branching Fractions, the $B^0 \rightarrow \pi^- \ell^+ \nu$ and $B^+ \rightarrow \eta \ell^+ \nu$ Form-Factor Shapes, and Determination of $|V_{ub}|$, Phys. Rev. D 83 (2011) 052011. [arXiv:1010.0987](https://arxiv.org/abs/1010.0987), doi:10.1103/PhysRevD.83.052011.
- [159] J. P. Lees, et al., Branching fraction and form-factor shape measurements of exclusive charmless semileptonic B decays, and determination of $|V_{ub}|$, Phys. Rev. D 86 (2012) 092004. [arXiv:1208.1253](https://arxiv.org/abs/1208.1253), doi:10.1103/PhysRevD.86.092004.

- [160] P. Ball, R. Zwicky, New results on $B \rightarrow \pi, K, \eta$ decay formfactors from light-cone sum rules, Phys. Rev. D 71 (2005) 014015. [arXiv:hep-ph/0406232](https://arxiv.org/abs/hep-ph/0406232), doi:10.1103/PhysRevD.71.014015.
- [161] E. Bagan, P. Ball, V. M. Braun, Radiative corrections to the decay $B \rightarrow \pi e \nu$ and the heavy quark limit, Phys. Lett. B 417 (1998) 154–162. [arXiv:hep-ph/9709243](https://arxiv.org/abs/hep-ph/9709243), doi:10.1016/S0370-2693(97)01371-3.
- [162] A. Khodjamirian, R. Rückl, S. Weinzierl, O. I. Yakovlev, Perturbative QCD correction to the $B \rightarrow \pi$ transition form-factor, Phys. Lett. B 410 (1997) 275–284. [arXiv:hep-ph/9706303](https://arxiv.org/abs/hep-ph/9706303), doi:10.1016/S0370-2693(97)00936-2.
- [163] H.-Y. Cheng, C.-Y. Cheung, C.-W. Hwang, Mesonic form-factors and the Isgur-Wise function on the light front, Phys. Rev. D 55 (1997) 1559–1577. [arXiv:hep-ph/9607332](https://arxiv.org/abs/hep-ph/9607332), doi:10.1103/PhysRevD.55.1559.
- [164] F. Bahr, D. Banerjee, F. Bernardoni, A. Joseph, M. Koren, H. Simma, R. Sommer, Continuum limit of the leading-order HQET form factor in $B_s \rightarrow K \ell \nu$ decays, Phys. Lett. B 757 (2016) 473–479. [arXiv:1601.04277](https://arxiv.org/abs/1601.04277), doi:10.1016/j.physletb.2016.03.088.
- [165] C. M. Bouchard, G. P. Lepage, C. Monahan, H. Na, J. Shigemitsu, $B_s \rightarrow K \ell \nu$ form factors from lattice QCD, Phys. Rev. D 90 (2014) 054506. [arXiv:1406.2279](https://arxiv.org/abs/1406.2279), doi:10.1103/PhysRevD.90.054506.
- [166] A. Bazavov, et al., $B_s \rightarrow K \ell \nu$ decay from lattice QCD, Phys. Rev. D 100 (3) (2019) 034501. [arXiv:1901.02561](https://arxiv.org/abs/1901.02561), doi:10.1103/PhysRevD.100.034501.
- [167] C.-D. Lu, W. Wang, Z.-T. Wei, Heavy-to-light form factors on the light cone, Phys. Rev. D 76 (2007) 014013. [arXiv:hep-ph/0701265](https://arxiv.org/abs/hep-ph/0701265), doi:10.1103/PhysRevD.76.014013.
- [168] A. Ali, G. Kramer, Y. Li, C.-D. Lu, Y.-L. Shen, W. Wang, Y.-M. Wang, Charmless non-leptonic B_s decays to PP , PV and VV final states in the pQCD approach, Phys. Rev. D 76 (2007) 074018. [arXiv:hep-ph/0703162](https://arxiv.org/abs/hep-ph/0703162), doi:10.1103/PhysRevD.76.074018.

- [169] R.-H. Li, C.-D. Lu, W. Wang, Transition form factors of B decays into p-wave axial-vector mesons in the perturbative QCD approach, Phys. Rev. D 79 (2009) 034014. [arXiv:0901.0307](#), doi:10.1103/PhysRevD.79.034014.
- [170] R. Aaij, et al., First observation of the decay $B_s^0 \rightarrow K^- \mu^+ \nu_\mu$ and Measurement of $|V_{ub}|/|V_{cb}|$, Phys. Rev. Lett. 126 (8) (2021) 081804. [arXiv:2012.05143](#), doi:10.1103/PhysRevLett.126.081804.
- [171] M. Wirbel, B. Stech, M. Bauer, Exclusive Semileptonic Decays of Heavy Mesons, Z. Phys. C 29 (1985) 637. doi:10.1007/BF01560299.
- [172] R. L. Workman, et al., Review of Particle Physics, PTEP 2022 (2022) 083C01. doi:10.1093/ptep/ptac097.
- [173] M. A. Ivanov, J. G. Körner, C.-T. Tran, Analyzing new physics in the decays $\bar{B}^0 \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ with form factors obtained from the covariant quark model, Phys. Rev. D 94 (9) (2016) 094028. [arXiv:1607.02932](#), doi:10.1103/PhysRevD.94.094028.
- [174] M. A. Ivanov, J. G. Körner, C.-T. Tran, Probing new physics in $\bar{B}^0 \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ using the longitudinal, transverse, and normal polarization components of the tau lepton, Phys. Rev. D 95 (3) (2017) 036021. [arXiv:1701.02937](#), doi:10.1103/PhysRevD.95.036021.
- [175] A. Bharucha, D. M. Straub, R. Zwicky, $B \rightarrow V\ell^+\ell^-$ in the Standard Model from light-cone sum rules, JHEP 08 (2016) 098. [arXiv:1503.05534](#), doi:10.1007/JHEP08(2016)098.
- [176] M. A. Ivanov, J. G. Körner, C. T. Tran, Exclusive decays $B \rightarrow \ell^-\bar{\nu}$ and $B \rightarrow D^{(*)}\ell^-\bar{\nu}$ in the covariant quark model, Phys. Rev. D 92 (11) (2015) 114022. [arXiv:1508.02678](#), doi:10.1103/PhysRevD.92.114022.
- [177] A. Issadykov, M. A. Ivanov, The decays $B_c \rightarrow J/\psi + \bar{\ell}\nu_\ell$ and $B_c \rightarrow J/\psi + \pi(K)$ in covariant confined quark model, Phys. Lett. B 783 (2018) 178–182. [arXiv:1804.00472](#), doi:10.1016/j.physletb.2018.06.056.

- [178] G. Caria, et al., Measurement of $\mathcal{R}(D)$ and $\mathcal{R}(D^*)$ with a semileptonic tagging method, Phys. Rev. Lett. 124 (16) (2020) 161803. [arXiv:1910.05864](https://arxiv.org/abs/1910.05864), doi:10.1103/PhysRevLett.124.161803.
- [179] N. R. Soni, A. Issadykov, A. N. Gadaria, Z. Tyulemissov, J. J. Patel, J. N. Pandya, Form factors and branching fraction calculations for $B_s \rightarrow D_s^{(*)}\ell^+\nu_\ell$ in view of LHCb observation, Eur. Phys. J. Plus 138 (2) (2023) 163. [arXiv:2110.12740](https://arxiv.org/abs/2110.12740), doi:10.1140/epjp/s13360-023-03779-8.
- [180] R. Aaij, et al., Measurement of Form-Factor-Independent Observables in the Decay $B^0 \rightarrow K^{*0}\mu^+\mu^-$, Phys. Rev. Lett. 111 (2013) 191801. [arXiv:1308.1707](https://arxiv.org/abs/1308.1707), doi:10.1103/PhysRevLett.111.191801.
- [181] R. Aaij, et al., Test of lepton universality with $B^0 \rightarrow K^{*0}\ell^+\ell^-$ decays, JHEP 08 (2017) 055. [arXiv:1705.05802](https://arxiv.org/abs/1705.05802), doi:10.1007/JHEP08(2017)055.
- [182] R. Aaij, et al., Test of lepton universality using $B^+ \rightarrow K^+\ell^+\ell^-$ decays, Phys. Rev. Lett. 113 (2014) 151601. [arXiv:1406.6482](https://arxiv.org/abs/1406.6482), doi:10.1103/PhysRevLett.113.151601.
- [183] M. Huschle, et al., Measurement of the branching ratio of $\bar{B} \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ relative to $\bar{B} \rightarrow D^{(*)}\ell^-\bar{\nu}_\ell$ decays with hadronic tagging at Belle, Phys. Rev. D 92 (7) (2015) 072014. [arXiv:1507.03233](https://arxiv.org/abs/1507.03233), doi:10.1103/PhysRevD.92.072014.
- [184] Y. Sato, et al., Measurement of the branching ratio of $\bar{B}^0 \rightarrow D^{*+}\tau^-\bar{\nu}_\tau$ relative to $\bar{B}^0 \rightarrow D^{*+}\ell^-\bar{\nu}_\ell$ decays with a semileptonic tagging method, Phys. Rev. D 94 (7) (2016) 072007. [arXiv:1607.07923](https://arxiv.org/abs/1607.07923), doi:10.1103/PhysRevD.94.072007.
- [185] R. Aaij, et al., Measurement of the ratio of branching fractions $\mathcal{B}(\bar{B}^0 \rightarrow D^{*+}\tau^-\bar{\nu}_\tau)/\mathcal{B}(\bar{B}^0 \rightarrow D^{*+}\mu^-\bar{\nu}_\mu)$, Phys. Rev. Lett. 115 (11) (2015) 111803, [Erratum: Phys. Rev. Lett. 115, 159901 (2015)]. [arXiv:1506.08614](https://arxiv.org/abs/1506.08614), doi:10.1103/PhysRevLett.115.111803.
- [186] A. Ali, P. Ball, L. T. Handoko, G. Hiller, A Comparative study of the decays $B \rightarrow (K, K^*)\ell^+\ell^-$ in standard model and supersymmetric theories, Phys. Rev. D 61 (2000) 074024. [arXiv:hep-ph/9910221](https://arxiv.org/abs/hep-ph/9910221), doi:10.1103/PhysRevD.61.074024.

- [187] M. Beneke, T. Feldmann, D. Seidel, Systematic approach to exclusive $B \rightarrow Vl^+l^-$, $V\gamma$ decays, Nucl. Phys. B 612 (2001) 25–58. [arXiv:hep-ph/0106067](https://arxiv.org/abs/hep-ph/0106067), doi:10.1016/S0550-3213(01)00366-2.
- [188] C.-H. Chen, C.-Q. Geng, Exclusive decays of $B \rightarrow K^{(*)}$ lepton+ lepton- in the PQCD, Phys. Rev. D 63 (2001) 114025. [arXiv:hep-ph/0103133](https://arxiv.org/abs/hep-ph/0103133), doi:10.1103/PhysRevD.63.114025.
- [189] A. Ali, G. Kramer, G.-h. Zhu, $B \rightarrow K^+l^+l^-$ decay in soft-collinear effective theory, Eur. Phys. J. C 47 (2006) 625–641. [arXiv:hep-ph/0601034](https://arxiv.org/abs/hep-ph/0601034), doi:10.1140/epjc/s2006-02596-4.
- [190] U. Egede, T. Hurth, J. Matias, M. Ramon, W. Reece, New observables in the decay mode $\bar{B}_d \rightarrow \bar{K}^{*0}l^+l^-$, JHEP 11 (2008) 032. [arXiv:0807.2589](https://arxiv.org/abs/0807.2589), doi:10.1088/1126-6708/2008/11/032.
- [191] C. Bobeth, G. Hiller, G. Piranishvili, CP Asymmetries in bar $B \rightarrow \bar{K}^*(\rightarrow \bar{K}\pi)\bar{\ell}\ell$ and Untagged \bar{B}_s , $B_s \rightarrow \phi(\rightarrow K^+K^-)\bar{\ell}\ell$ Decays at NLO, JHEP 07 (2008) 106. [arXiv:0805.2525](https://arxiv.org/abs/0805.2525), doi:10.1088/1126-6708/2008/07/106.
- [192] W. Altmannshofer, P. Ball, A. Bharucha, A. J. Buras, D. M. Straub, M. Wick, Symmetries and Asymmetries of $B \rightarrow K^*\mu^+\mu^-$ Decays in the Standard Model and Beyond, JHEP 01 (2009) 019. [arXiv:0811.1214](https://arxiv.org/abs/0811.1214), doi:10.1088/1126-6708/2009/01/019.
- [193] T. Hurth, M. Nakao, Radiative and Electroweak Penguin Decays of B Mesons, Ann. Rev. Nucl. Part. Sci. 60 (2010) 645–677. [arXiv:1005.1224](https://arxiv.org/abs/1005.1224), doi:10.1146/annurev.nucl.012809.104424.
- [194] T. Blake, G. Lanfranchi, D. M. Straub, Rare B Decays as Tests of the Standard Model, Prog. Part. Nucl. Phys. 92 (2017) 50–91. [arXiv:1606.00916](https://arxiv.org/abs/1606.00916), doi:10.1016/j.ppnp.2016.10.001.
- [195] R. Aaij, et al., Observation of the suppressed decay $\Lambda_b^0 \rightarrow p\pi^-\mu^+\mu^-$, JHEP 04 (2017) 029. [arXiv:1701.08705](https://arxiv.org/abs/1701.08705), doi:10.1007/JHEP04(2017)029.

- [196] G. D'Ambrosio, G. F. Giudice, G. Isidori, A. Strumia, Minimal flavor violation: An Effective field theory approach, Nucl. Phys. B 645 (2002) 155–187. [arXiv:hep-ph/0207036](https://arxiv.org/abs/hep-ph/0207036), doi:10.1016/S0550-3213(02)00836-2.
- [197] J. Albrecht, S. Reichert, D. van Dyk, Status of rare exclusive B meson decays in 2018, Int. J. Mod. Phys. A 33 (18n19) (2018) 1830016. [arXiv:1806.05010](https://arxiv.org/abs/1806.05010), doi:10.1142/S0217751X18300168.
- [198] R. Aaij, et al., Measurement of CP violation parameters and polarisation fractions in $B_s^0 \rightarrow J/\psi \bar{K}^{*0}$ decays, JHEP 11 (2015) 082. [arXiv:1509.00400](https://arxiv.org/abs/1509.00400), doi:10.1007/JHEP11(2015)082.
- [199] A. J. Buras, M. Munz, Effective Hamiltonian for $B \rightarrow X(s)$ e+ e- beyond leading logarithms in the NDR and HV schemes, Phys. Rev. D 52 (1995) 186–195. [arXiv:hep-ph/9501281](https://arxiv.org/abs/hep-ph/9501281), doi:10.1103/PhysRevD.52.186.
- [200] F. Kruger, L. M. Sehgal, CP violation in the decay $B \rightarrow X(d)$ e+ e-, Phys. Rev. D 55 (1997) 2799–2805. [arXiv:hep-ph/9608361](https://arxiv.org/abs/hep-ph/9608361), doi:10.1103/PhysRevD.55.2799.
- [201] C.-H. Chen, C. Q. Geng, Baryonic rare decays of $\Lambda(b) \rightarrow \Lambda$ lepton+ lepton-, Phys. Rev. D 64 (2001) 074001. [arXiv:hep-ph/0106193](https://arxiv.org/abs/hep-ph/0106193), doi:10.1103/PhysRevD.64.074001.
- [202] N. G. Deshpande, J. Trampetic, K. Panose, Resonance Background to the Decays $b \rightarrow s$ Lepton+ Lepton-, $B \rightarrow K^*$ Lepton+ Lepton- and $B \rightarrow K$ Lepton+ Lepton-, Phys. Rev. D 39 (1989) 1461. doi:10.1103/PhysRevD.39.1461.
- [203] M. Jezabek, J. H. Kuhn, Lepton Spectra from Heavy Quark Decay, Nucl. Phys. B 320 (1989) 20–44. doi:10.1016/0550-3213(89)90209-5.
- [204] C. S. Lim, T. Morozumi, A. I. Sanda, A Prediction for d gamma ($b \rightarrow s$ Lepton anti-Lepton) / $d q^{**2}$ Including the Long Distance Effects, Phys. Lett. B 218 (1989) 343–347. doi:10.1016/0370-2693(89)91593-1.
- [205] M. Misiak, The $b \rightarrow se^+e^-$ and $b \rightarrow s\gamma$ decays with next-to-leading logarithmic QCD corrections, Nucl. Phys. B 393 (1993) 23–45, [Erratum: Nucl.Phys.B 439, 461–465 (1995)]. doi:10.1016/0550-3213(93)90235-H.

- [206] P. J. O'Donnell, H. K. K. Tung, Resonance contributions to the decay $b \rightarrow s(l^+ l^-)$, Phys. Rev. D 43 (1991) 2067–2069. doi:10.1103/PhysRevD.43.R2067.
- [207] A. Ali, T. Mannel, T. Morozumi, Forward backward asymmetry of dilepton angular distribution in the decay $b \rightarrow s l^+ l^-$, Phys. Lett. B 273 (1991) 505–512. doi:10.1016/0370-2693(91)90306-B.
- [208] C. Bobeth, M. Misiak, J. Urban, Photonic penguins at two loops and m_t dependence of $BR[B \rightarrow X_s l^+ l^-]$, Nucl. Phys. B 574 (2000) 291–330. arXiv:hep-ph/9910220, doi:10.1016/S0550-3213(00)00007-9.
- [209] G. Ganbold, T. Gutsche, M. A. Ivanov, V. E. Lyubovitskij, On the meson mass spectrum in the covariant confined quark model, J. Phys. G 42 (7) (2015) 075002. arXiv:1410.3741, doi:10.1088/0954-3899/42/7/075002.
- [210] S. Dubnička, A. Z. Dubničková, A. Issadykov, M. A. Ivanov, A. Liptaj, S. K. Sakhiyev, Decay $B_s \rightarrow \phi \ell^+ \ell^-$ in covariant quark model, Phys. Rev. D 93 (9) (2016) 094022. arXiv:1602.07864, doi:10.1103/PhysRevD.93.094022.
- [211] R. N. Faustov, V. O. Galkin, Rare B_s decays in the relativistic quark model, Eur. Phys. J. C 73 (10) (2013) 2593. arXiv:1309.2160, doi:10.1140/epjc/s10052-013-2593-5.
- [212] R. Aaij, et al., First measurement of the differential branching fraction and CP asymmetry of the $B^\pm \rightarrow \pi^\pm \mu^+ \mu^-$ decay, JHEP 10 (2015) 034. arXiv:1509.00414, doi:10.1007/JHEP10(2015)034.
- [213] R. Aaij, et al., Evidence for the decay $B_S^0 \rightarrow \overline{K}^{*0} \mu^+ \mu^-$, JHEP 07 (2018) 020. arXiv:1804.07167, doi:10.1007/JHEP07(2018)020.
- [214] G. Buchalla, A. J. Buras, The rare decays $K \rightarrow \pi \nu \bar{\nu}$, $B \rightarrow X \nu \bar{\nu}$ and $B \rightarrow l^+ l^-$: An Update, Nucl. Phys. B 548 (1999) 309–327. arXiv:hep-ph/9901288, doi:10.1016/S0550-3213(99)00149-2.
- [215] J. A. Bailey, et al., $B \rightarrow \pi \ell \ell$ form factors for new-physics searches from lattice QCD, Phys. Rev. Lett. 115 (15) (2015) 152002. arXiv:1507.01618, doi:10.1103/PhysRevLett.115.152002.

- [216] N. R. Soni, J. N. Pandya, Decay $D \rightarrow K^{(*)}\ell^+\nu_\ell$ in covariant quark model, Phys. Rev. D 96 (1) (2017) 016017, [Erratum: Phys.Rev.D 99, 059901 (2019)]. arXiv:1706.01190, doi:10.1103/PhysRevD.96.016017.
- [217] N. R. Soni, M. A. Ivanov, J. G. Körner, J. N. Pandya, P. Santorelli, C. T. Tran, Semileptonic $D_{(s)}$ -meson decays in the light of recent data, Phys. Rev. D 98 (11) (2018) 114031. arXiv:1810.11907, doi:10.1103/PhysRevD.98.114031.
- [218] M. A. Ivanov, J. G. Körner, J. N. Pandya, P. Santorelli, N. R. Soni, C.-T. Tran, Exclusive semileptonic decays of D and D_s mesons in the covariant confining quark model, Front. Phys. (Beijing) 14 (6) (2019) 64401. arXiv:1904.07740, doi:10.1007/s11467-019-0908-1.
- [219] A. Ali, A. Y. Parkhomenko, A. V. Rusov, Precise Calculation of the Dilepton Invariant-Mass Spectrum and the Decay Rate in $B^\pm \rightarrow \pi^\pm \mu^+ \mu^-$ in the SM, Phys. Rev. D 89 (9) (2014) 094021. arXiv:1312.2523, doi:10.1103/PhysRevD.89.094021.
- [220] D. Seidel, Analytic two loop virtual corrections to $b \rightarrow d l + l -$, Phys. Rev. D 70 (2004) 094038. arXiv:hep-ph/0403185, doi:10.1103/PhysRevD.70.094038.
- [221] B. Aubert, et al., Search for the rare decay $B \rightarrow \pi l^+ l^-$, Phys. Rev. Lett. 99 (2007) 051801. arXiv:hep-ex/0703018, doi:10.1103/PhysRevLett.99.051801.
- [222] J. P. Lees, et al., Search for the rare decays $B \rightarrow \pi l^+ l^-$ and $B^0 \rightarrow \eta l^+ l^-$, Phys. Rev. D 88 (3) (2013) 032012. arXiv:1303.6010, doi:10.1103/PhysRevD.88.032012.
- [223] S. Dubnička, A. Z. Dubničková, N. Habyl, M. A. Ivanov, A. Liptaj, G. S. Nurbakova, Decay $B \rightarrow K^*(\rightarrow K\pi)\ell^+\ell^-$ in covariant quark model, Few Body Syst. 57 (2) (2016) 121–143. arXiv:1511.04887, doi:10.1007/s00601-015-1034-4.
- [224] N. R. Soni, A. Issadykov, A. N. Gadaria, J. J. Patel, J. N. Pandya, Rare $b \rightarrow d$ decays in covariant confined quark model, Eur. Phys. J. A 58 (3) (2022) 39. arXiv:2008.07202, doi:10.1140/epja/s10050-022-00685-y.

- [225] L. Zhang, X.-W. Kang, X.-H. Guo, L.-Y. Dai, T. Luo, C. Wang, A comprehensive study on the semileptonic decay of heavy flavor mesons, JHEP 02 (2021) 179. [arXiv:2012.04417](https://arxiv.org/abs/2012.04417), doi:10.1007/JHEP02(2021)179.
- [226] R. N. Faustov, V. O. Galkin, X.-W. Kang, Relativistic description of the semileptonic decays of bottom mesons, Phys. Rev. D 106 (1) (2022) 013004. [arXiv:2206.10277](https://arxiv.org/abs/2206.10277), doi:10.1103/PhysRevD.106.013004.
- [227] T. Gutsche, M. A. Ivanov, J. G. Körner, V. E. Lyubovitskij, P. Santorelli, N. Habyl, Semileptonic decay $\Lambda_b \rightarrow \Lambda_c + \tau^- + \bar{\nu}_\tau$ in the covariant confined quark model, Phys. Rev. D 91 (7) (2015) 074001, [Erratum: Phys.Rev.D 91, 119907 (2015)]. [arXiv:1502.04864](https://arxiv.org/abs/1502.04864), doi:10.1103/PhysRevD.91.074001.
- [228] J. Matias, F. Mescia, M. Ramon, J. Virto, Complete Anatomy of $\bar{B}_d -> \bar{K}^{*0}(-> K\pi)l^+l^-$ and its angular distribution, JHEP 04 (2012) 104. [arXiv:1202.4266](https://arxiv.org/abs/1202.4266), doi:10.1007/JHEP04(2012)104.
- [229] R. Aaij, et al., Angular analysis of the $B^0 \rightarrow K^{*0}\mu^+\mu^-$ decay using 3 fb^{-1} of integrated luminosity, JHEP 02 (2016) 104. [arXiv:1512.04442](https://arxiv.org/abs/1512.04442), doi:10.1007/JHEP02(2016)104.
- [230] R. Aaij, et al., Measurement of CP -Averaged Observables in the $B^0 \rightarrow K^{*0}\mu^+\mu^-$ Decay, Phys. Rev. Lett. 125 (1) (2020) 011802. [arXiv:2003.04831](https://arxiv.org/abs/2003.04831), doi:10.1103/PhysRevLett.125.011802.
- [231] S. Wehle, et al., Lepton-Flavor-Dependent Angular Analysis of $B \rightarrow K^*\ell^+\ell^-$, Phys. Rev. Lett. 118 (11) (2017) 111801. [arXiv:1612.05014](https://arxiv.org/abs/1612.05014), doi:10.1103/PhysRevLett.118.111801.
- [232] S. Descotes-Genon, J. Matias, M. Ramon, J. Virto, Implications from clean observables for the binned analysis of $B -> K^*\mu^+\mu^-$ at large recoil, JHEP 01 (2013) 048. [arXiv:1207.2753](https://arxiv.org/abs/1207.2753), doi:10.1007/JHEP01(2013)048.
- [233] J. Matias, On the S-wave pollution of $B -> K^* l^+l^-$ observables, Phys. Rev. D 86 (2012) 094024. [arXiv:1209.1525](https://arxiv.org/abs/1209.1525), doi:10.1103/PhysRevD.86.094024.

- [234] B. Kindra, N. Mahajan, Predictions of angular observables for $\bar{B}_s \rightarrow K^* \ell \ell$ and $\bar{B} \rightarrow \rho \ell \ell$ in the standard model, Phys. Rev. D 98 (9) (2018) 094012. [arXiv:1803.05876](#), doi:10.1103/PhysRevD.98.094012.
- [235] B. Aubert, et al., Search for flavor-changing neutral current and lepton flavor violating decays of $D^0 \rightarrow \ell^+ \ell^-$, Phys. Rev. Lett. 93 (2004) 191801. [arXiv:hep-ex/0408023](#), doi:10.1103/PhysRevLett.93.191801.
- [236] M. Petric, et al., Search for leptonic decays of D^0 mesons, Phys. Rev. D 81 (2010) 091102. [arXiv:1003.2345](#), doi:10.1103/PhysRevD.81.091102.
- [237] J. P. Lees, et al., Search for the decay modes $D^0 \rightarrow e^+ e^-$, $D^0 \rightarrow \mu^+ \mu^-$, and $D^0 \rightarrow e \mu$, Phys. Rev. D 86 (2012) 032001. [arXiv:1206.5419](#), doi:10.1103/PhysRevD.86.032001.
- [238] R. Aaij, et al., Search for the rare decay $D^0 \rightarrow \mu^+ \mu^-$, Phys. Lett. B 725 (2013) 15–24. [arXiv:1305.5059](#), doi:10.1016/j.physletb.2013.06.037.
- [239] M. Ablikim, et al., First observation of the semileptonic decay $\Lambda c \rightarrow p K^- e + \nu e$, Phys. Rev. D 106 (11) (2022) 112010. [arXiv:2207.11483](#), doi:10.1103/PhysRevD.106.112010.
- [240] J. P. Alexander, et al., First observation of $\Xi_c(c) \rightarrow \Xi_c 0$ electron-neutrino and a measurement of the $\Xi_c(c) \rightarrow \Xi_c 0$ lifetime ratio, Phys. Rev. Lett. 74 (1995) 3113–3117, [Erratum: Phys. Rev. Lett. 75, 4155 (1995)]. doi:10.1103/PhysRevLett.74.3113.
- [241] Q.-A. Zhang, et al., First lattice QCD calculation of semileptonic decays of charmed-strange baryons Ξ_c^* , Chin. Phys. C 46 (1) (2022) 011002. [arXiv:2103.07064](#), doi:10.1088/1674-1137/ac2b12.
- [242] Y. B. Li, et al., Measurements of the branching fractions of the semileptonic decays $\Xi_c^0 \rightarrow \Xi^- \ell^+ \nu_\ell$ and the asymmetry parameter of $\Xi_c^0 \rightarrow \Xi^- \pi^+$, Phys. Rev. Lett. 127 (12) (2021) 121803. [arXiv:2103.06496](#), doi:10.1103/PhysRevLett.127.121803.
- [243] J. Zhu, Hadronization studies at the LHC with ALICE, PoS ICHEP2020 (2021) 524. [arXiv:2104.11309](#), doi:10.22323/1.390.0524.

- [244] R. N. Faustov, V. O. Galkin, Semileptonic Ξ_c baryon decays in the relativistic quark model, *Eur. Phys. J. C* 79 (8) (2019) 695. [arXiv:1905.08652](https://arxiv.org/abs/1905.08652), doi:10.1140/epjc/s10052-019-7214-5.
- [245] Z.-X. Zhao, Weak decays of heavy baryons in the light-front approach, *Chin. Phys. C* 42 (9) (2018) 093101. [arXiv:1803.02292](https://arxiv.org/abs/1803.02292), doi:10.1088/1674-1137/42/9/093101.
- [246] K. Azizi, Y. Sarac, H. Sundu, Light cone QCD sum rules study of the semileptonic heavy Ξ_Q and Ξ'_Q transitions to Ξ and Σ baryons, *Eur. Phys. J. A* 48 (2012) 2. [arXiv:1107.5925](https://arxiv.org/abs/1107.5925), doi:10.1140/epja/i2012-12002-1.
- [247] C. Q. Geng, Y. K. Hsiao, C.-W. Liu, T.-H. Tsai, Antitriplet charmed baryon decays with SU(3) flavor symmetry, *Phys. Rev. D* 97 (7) (2018) 073006. [arXiv:1801.03276](https://arxiv.org/abs/1801.03276), doi:10.1103/PhysRevD.97.073006.
- [248] C.-Q. Geng, C.-W. Liu, T.-H. Tsai, S.-W. Yeh, Semileptonic decays of anti-triplet charmed baryons, *Phys. Lett. B* 792 (2019) 214–218. [arXiv:1901.05610](https://arxiv.org/abs/1901.05610), doi:10.1016/j.physletb.2019.03.056.
- [249] Y.-K. Hsiao, L. Yang, C.-C. Lih, S.-Y. Tsai, Charmed Ω_c weak decays into Ω in the light-front quark model, *Eur. Phys. J. C* 80 (11) (2020) 1066. [arXiv:2009.12752](https://arxiv.org/abs/2009.12752), doi:10.1140/epjc/s10052-020-08619-y.
- [250] A. Faessler, T. Gutsche, M. A. Ivanov, J. G. Korner, V. E. Lyubovitskij, Semileptonic decays of double heavy baryons in a relativistic constituent three-quark model, *Phys. Rev. D* 80 (2009) 034025. [arXiv:0907.0563](https://arxiv.org/abs/0907.0563), doi:10.1103/PhysRevD.80.034025.
- [251] Y. B. Li, et al., First test of lepton flavor universality in the charmed baryon decays $\Omega_c \rightarrow \Omega^- \ell^+ \nu \ell$ using data of the Belle experiment, *Phys. Rev. D* 105 (9) (2022) L091101. [arXiv:2112.10367](https://arxiv.org/abs/2112.10367), doi:10.1103/PhysRevD.105.L091101.