

List of Tables

2.1	Leptonic decay constant f_H (in MeV)	25
2.2	CCQM model parameters: quark masses, meson size parameters and infrared cut-off parameter (all in GeV)	25
3.1	Form factors and double pole parameters for $B_s \rightarrow K^{(*)-}$ transitions . .	33
3.2	Form factors for $B_s \rightarrow K$ transitions at maximum recoil	34
3.3	Form factors for $B_s \rightarrow K^*(892)$ transition at maximum recoil	35
3.4	Branching fraction for $B_s \rightarrow K^{(*)-}$ transitions (in the unit of 10^{-4}) . .	36
3.5	Form factors and double pole parameters for $B_s \rightarrow D_s^{(*)-}$ transitions . .	37
3.6	Comparison of the form factors at maximum recoil $F(0)$ with other theoretical approaches	37
3.7	Normalized decay rates in terms of w bins in comparison with LQCD and LHCb data.	39
3.8	Branching fraction for $B_s \rightarrow D_s^{(*)-}$ transitions (in %)	40
3.9	Ratio of semileptonic decay widths with heavy quark expansion (HQE), LHCb and Lattice QCD data	41
3.10	Masses, total widths and dilepton decay widths of vector resonance states [12]	43
3.11	Values of the input parameters [13] and SM Wilson coefficients [14]. Values of masses are in GeV	44
3.12	Form factors and double pole parameters for $B \rightarrow (\pi, \rho, \omega)\ell^+\ell^-$ transitions	47

3.13	Branching fractions of $B \rightarrow (\pi, \rho, \omega)$ transitions	52
3.14	Branching fractions of $B_s \rightarrow K/K^*$ transitions	53
3.15	Form factor for $B \rightarrow \pi$ transitions at maximum recoil	55
3.16	Form factors for $B \rightarrow \rho$ transitions at maximum recoil	58
3.17	Form factors for $B \rightarrow \omega$ transitions at maximum recoil	60
3.18	Comparison of form factors at higher q^2 values with LQCD (data from Table VI of Ref. [11]) for $B_s \rightarrow K$ and $B \rightarrow \pi$ channels	61
3.19	Comparison of form factors at higher q^2 (in GeV^2) values with LQCD [15] for $B_s \rightarrow K^*(892)$ channel	62
3.20	Partial branching fractions for $B^+ \rightarrow \pi^+ \mu^+ \mu^-$ in Unit of 10^{-9}	63
3.21	Ratios of the Branching fractions	64
4.1	Other Observable Parameters for $B_s \rightarrow (K^{(*)-}, D_s^{(*)-})$ transitions . . .	70
4.2	q^2 - averages of polarization observables over the whole allowed kine- matic region for $B^+ \rightarrow \rho^+ \ell^+ \ell^-$, $B^0 \rightarrow \rho^0 \ell^+ \ell^-$ channels	80
4.3	q^2 - averages of polarization observables over the whole allowed kine- matic region for $B^0 \rightarrow \omega \ell^+ \ell^-$ and $B_s^0 \rightarrow \bar{K}^*(892)^0 \ell^+ \ell^-$ channels	81
4.4	Value of other observables at different bin values for $B \rightarrow \rho \ell^+ \ell^-$ channel where, $\ell = e$ and μ	82
4.5	Value of other observables at different bin values for $B^0 \rightarrow \omega \ell^+ \ell^-$ channel where, $\ell = e$ and μ	83
4.6	Value of other observables at different bin values for $B_s^0 \rightarrow \bar{K}^*(892)^0 \ell^+ \ell^-$ channel where, $\ell = e$ and μ	84
5.1	Overview of Branching fraction for the semileptonic decay of baryons .	88