

## LIST OF FIGURES

<b>Studies on development of Saline Tolerance in plants</b>		
<b>Figure No</b>	<b>Title</b>	<b>Page No</b>
<b>Chapter -1</b>	<b>REVIEW OF LITERATURE</b>	
Fig. 1.1	Regulation of ion (e.g., Na <sup>+</sup> and K <sup>+</sup> ) homeostasis by the SOS pathway	4
Fig.1.2	Biosynthesis of betaine	20
<b>Chapter -2</b>	<b>TISSUE CULTURE STUDIES FOR DEVELOPMENT OF REGENERATION PROTOCOL FOR GROUNDNUT</b>	
Fig.-2.1A	Seed coat peeled off from pre-soaked sterilized seed	32
Fig.-2.1B	Embryonated cotyledon with immature leaflets intact in pre-soaked sterilized seed	32
Fig.2.2A	Young green folded leaflets developed during seed germination within 7 day incubation	32
Fig.2.2B	Young green folded leaflet lamina separated from seeds and used as explants	32
Fig. 2.3A	Shoot bud initiation after 15 days of inoculation	34
Fig.2.3B	Shoot elongation 30 days of inoculation	34
Fig. 2.4A	Initiation and development of shoot buds at the proximal end of the de-embryonated CN explants	39
Fig. 2.4B	Elongation of shoots after two sub culture	39

Fig. 2.5	Rooting of the elongated shoots	40
Fig. 2.6A	Initiation of callus at the cut edge of the leaf margin	43
Fig. 2.6B	White-greenish friable callus growth after 3 sub culture on MS medium with BAP & 2,4,D	43
Fig.2.7A	Brownish-hard compact callus with development of root initial after 3 sub culture on MS medium with kinetin & NAA.	43
Fig.2.7B	Development of shoot buds and plantlets from the brownish hard callus on MS medium with reduced kinetin concentration after 5 subcultures	43
<b>Chapter 3</b>	<b>BIOCHEMICAL CHANGES SEEN IN GROUND NUT DUE TO SALT STRESS</b>	
Fig. 3.1	Reactions producing ROS and organic free radicals	47
Fig. 3.2	Defence systems (enzymes, antioxidants) against ROS	49
Fig. 3.3	ROS and its harmful effect.	51
Fig. 3.4	Ground nut plants under different level of NaCl stress.	60
Fig. 3.5	Ground nut plants under NaCl stress Control and 10mM on 22nd day	60
Fig. 3.6	Ground nut plants under NaCL stress 25mM and 50mM on 22nd day.	62
Fig. 3.7	Leaves size of plants under different level of NaCl stress.	62
Fig. 3.8	Ground nut plants under NaCl stress 100mM on 22nd day.	63

Fig. 3.9	<i>In vitro</i> germinating seeds on MS medium on 3 <sup>rd</sup> day of inoculation with different level of NaCl stress.	64
Fig. 3.10	Release of H <sub>2</sub> O <sub>2</sub> in different concentration of NaCl stress.	65
Fig. 3.11	Catalase activity at different concentration of NaCl stress.	66
Fig. 3.12	Peroxidase activity at different concentration of NaCl stress.	67
<b>Chapter 4</b>	<b>TRANSFORMATION OF AGROBACTERIUM</b>	
Fig. 4.1	Confirmation of the plasmid by agarose gel electrophoresis.	76
Fig. 4.2	Triparental mating (A: <i>Agrobacterium tumefaciens</i> (Rif <sup>R</sup> Tet <sup>R</sup> ); D: Donor pHS724 (Kan <sup>R</sup> ); H: Helper strain (Kan <sup>R</sup> ))	77
Fig. 4.3	<i>Agrobacterium tumefaciens</i> on LA plate	79
Fig. 4.4	<i>Agrobacterium</i> and <i>Agrobacterium</i> transformants on LA Plate	79
Fig. 4.5	Ketolactose test	81
Fig. 4.6	<i>Agrobacterium</i> and <i>Agrobacterium</i> transformants on a medium containing 0.1M NaCl	81
Fig. 4.7	<i>Agrobacterium</i> and <i>Agrobacterium</i> transformants on a medium containing 0.2M NaCl	82
Fig. 4.8	<i>Agrobacterium</i> and <i>Agrobacterium</i> transformants on a medium containing 0.3M NaCl	82

Fig. 4.9	<i>Agrobacterium</i> and <i>Agrobacterium</i> transformants on a medium containing 0.4M NaCl	83
<b>Chapter 5</b>	<b>TRANSFORMATION OF GROUNDNUT PLANTS FOR SALINE TOLERANCE</b>	
Fig. 5.1	Ti Plasmid	88
Fig. 5.2	Transformation of plant cells by <i>Agrobacterium</i> (Gelvin, 2005)	89
Fig. 5.3	Binary Vector System	92
Fig. 5.4	pHS724 Vector map	94
Fig. 5.5A	Infection of leaf explants after 5 day co-cultivation with <i>Agrobacterium</i> .	98
Fig. 5.5B	Infection of cotyledon explants after 5 day co-cultivation with <i>Agrobacterium</i>	98
Fig. 5.6A	Swelling of explants with growing buds and callus at the proximal end of cotyledon explants.	102
Fig. 5.6B	Shoot bud producing plants from the proximal end of cotyledon explants.	102
Fig. 5.7A	Untransformed plants in MS medium without kanamycin	104
Fig. 5.7B	Untransformed plants in MS medium with 60mM kanamycin	104
Fig. 5.7C	Transformed plants in MS medium with 60mM kanamycin	106
Fig. 5.8	Transformed multiple shoot from embryo axis explants.	106
Fig. 5.9	Comparison of transformed plant with the control plants growing on MS medium without stress.	109

Fig. 5.10	Transformed plants which survived 2 cycle selection under kanamycin rich medium growing on MS medium with 100 mM NaCl	109
Fig. 5.11	Comparison of performance of transformed plants with control plants.	110
Fig. 5.12A	Comparison of X – gluc staining in the leaves of control & transformed plants	111
Fig. 5.12B	Blue colored X – gluc staining in the leaves of transformed plants.	111
Fig. 5.13	Comparison of X – gluc staining in the cotyledon tissues of control & transformed plants	112
Fig. 5.14	Blue spots of X – gluc staining in the cells of stems	112