

## *Table of Contents*

Content	Page No.
1. Introduction.....	1-43
2. Materials and Methods.....	44-58
3. Results and Discussion.....	59-129
<b>Chapter 1.</b> To study population structure of <i>Clibanarius</i> <i>rhabdodactylus</i> .....	59-88
<b>Chapter 2.</b> To study the seasonal variation in intertidal distribution pattern of <i>Clibanarius rhabdodactylus</i> . ....	89-99
<b>Chapter 3.</b> To study the shell utilization pattern of <i>Clibanarius</i> <i>rhabdodactylus</i> .....	100-129
4. Summary, Conclusion and Recommendations.....	130-138
5. References.....	139-184
6. Publications and Presentations.....	185-190

## *List of Figures*

Figure 1.	Classification of marine environment.	4
Figure 2.	General habit of marine organisms.	5
Figure 3.	Intertidal zonation pattern showing intertidal distribution of marine organisms.	10
Figure 4.	External morphology of hermit crab.	16
Figure 5.	Internal arrangement of hermit crab in an empty gastropod shell	30
Figure 6.	Hermit crabs occupying variety of pseudoshells.	34
Figure 7.	Dorsal habitus of <i>Clibanarius rhabdodactylus</i> Forest, 1953.	42
Figure 8.	Location of study area 1. Veraval; 2. Sutrapada and 3. Dhamlej.	44
Figure 9.	Abundance of <i>Clibanarius rhabdodactylus</i> in different microhabitats of study sites.	45
Figure 10.	Habitat of study area (A) Veraval (B) Sutrapada (C) Dhamlej; (D) Quadrature sampling method for collection of <i>Clibanarius rhabdodactylus</i> individuals.	48
Figure 11.	Ventral view of female <i>Clibanarius rhabdodactylus</i> showing gonopores on the first segment of the second pair of walking legs (third periopod).	52
Figure 12.	Eggs of <i>Clibanarius rhabdodactylus</i> measured under ocular micrometre.	52
Figure 13.	External morphology of a gastropod shell.	58
Figure 14.	Association between monthly temperature and <i>Clibanarius rhabdodactylus</i> juveniles (of both the sexes) occurrence at Veraval coast.	64

Figure 15.	Overall size–frequency distributions of all individuals of <i>Clibanarius rhabdodactylus</i> collected at Veraval coast.	67
Figure 16.	Size frequency distribution <i>Clibanarius rhabdodactylus</i> in the months of winter season.	71
Figure 17.	Size frequency distribution <i>Clibanarius rhabdodactylus</i> in the months of summer season.	72
Figure 18.	Size frequency distribution <i>Clibanarius rhabdodactylus</i> in the months of monsoon season.	75
Figure 19.	Relationship between <i>Clibanarius rhabdodactylus</i> shield length and A. total number of eggs; B. Egg size and C. weight of egg mass.	77
Figure 20.	Relationship between <i>Clibanarius rhabdodactylus</i> wet weight and A. total number of eggs; B. Egg size and C. weight of egg mass.	78
Figure 21.	Seasonal variation in the mean values of ambient temperature and tidepool temperature at different intertidal region.	92
Figure 22.	Seasonal variation in the mean values of abundance of different sexes of <i>Clibanarius rhabdodactylus</i> .	93
Figure 23.	A. Intertidal microhabitat coverage of the study area; Intertidal distribution of <i>Clibanarius rhabdodactylus</i> at Veraval coast in B. winter season; C. Summer season and D. Monsoon season.	95
Figure 24.	Clustering observed in <i>Clibanarius rhabdodactylus</i> at the rocky intertidal region of Veraval coast, Saurashtra.	98
Figure 25.	Size frequency (SL) distribution of male and female individuals of (A) <i>Clibanarius rhabdodactylus</i> (n= 1000) and (B) <i>Clibanarius ransonii</i> (n= 1000).	105

Figure 26.	Five most frequently occupied gastropod shells by the hermit crab species <i>Clibanarius rhabdodactylus</i> and <i>Clibanarius ransoni</i> .	106
Figure 27.	Abundance of commonly occurring gastropod species in the intertidal zone of Veraval, Saurashtra coast, Gujarat, India. (CC: <i>C. caeruleum</i> , LC: <i>L. coronata</i> , TG: <i>T. granulata</i> , TB: <i>T. bruneus</i> , PU: <i>P. undosa</i> ).	113
Figure 28.	Box and whisker plots of (A) shell length, (B) shell aperture length, (C) shell aperture width, (D) shell volume and (E) shell dry weight of five frequently occupied shells by <i>Clibanarius rhabdodactylus</i> and <i>Clibanarius ransoni</i> . CC: <i>C. caeruleum</i> , LC: <i>L. coronata</i> , TG: <i>T. granulata</i> , TB: <i>T. bruneus</i> , PU: <i>P. undosa</i> ; Box = 25 <sup>th</sup> and 75 <sup>th</sup> percentiles, midline = mean, whiskers = minimum and maximum values.	123
Figure 29.	Canonical correspondence analysis (CCA) of shell use by <i>Clibanarius rhabdodactylus</i> and <i>Clibanarius ransoni</i> of different size classes as influenced by shell parameters. CC: <i>C. caeruleum</i> , LC: <i>L. coronata</i> , TG: <i>T. granulata</i> , TB: <i>T. bruneus</i> , PU: <i>P. undosa</i> . Lines indicate shell attributes in a direction of increasing magnitude. Filled circles indicate hermit crab species (CRH = <i>C. rhabdodactylus</i> , CRS= <i>C. ransoni</i> ) by sex (M = male, F = female) and size class (please refer to Table 11 for the explanation).	124

## *List of Tables*

Table 1.	Carapace shield length values of different sexes of <i>Clibanarius rhabdodactylus</i> (ANOVA; ***p<0.001; N = total individuals; SL= Shield length).	63
Table 2.	Total number of <i>Clibanarius rhabdodactylus</i> specimens collected at rocky intertidal region of Veraval. (M = Male; NOF = Non-ovigerous female; OF= Ovigerous female).	66
Table 3.	Summary statistics of hermit crab weight, shield length, shield width, weight of egg mass, egg number and egg size in ovigerous females of <i>Clibanarius rhabdodactylus</i> collected from Veraval coast. (N = total individuals).	76
Table 4.	Carapace shield length values of <i>Clibanarius rhabdodactylus</i> and <i>Clibanarius ransoni</i> . (ANOVA; ***p<0.001; N = total individuals; SL = Shield length; M = Male; NOF = Non-ovigerous female; OF = Ovigerous female).	103
Table 5.	Gastropod shell utilization by <i>Clibanarius rhabdodactylus</i> . (N = total individuals; M = Male; NOF = Non-ovigerous female; OF = Ovigerous female).	108
Table 6.	Gastropod shell utilization by <i>Clibanarius ransoni</i> . (N = total individuals; M = Male; NOF = Non-ovigerous female; OF = Ovigerous female).	110
Table 7.	Regression equation in relation to the morphological parameters of different sexes of <i>Clibanarius rhabdodactylus</i> and <i>Clibanarius ransoni</i> and that of gastropod shells measures (*P < 0.05; **P<0.01; ***P<0.001; Shield length= SL; Hermit crab wet weight= HW; Shell length= SHL; Shell aperture length= SHAL; Shell aperture width= SHAW; Shell dry weight= SHW; Shell volume= SHV).	114

Table 8.	Mean values of different morphological parameters of five highly occupied gastropod shells by different sexes of <i>Clibanarius rhabdodactylus</i> and <i>Clibanarius ransonii</i> (Male= M; Ovigerous female= OF; Female= F; Shield length= SL; Shell length= SHL; Shell aperture length= SHAL; Shell aperture width= SHAW; Shell dry weight= SHW; Shell volume= SHV; NO= Not occupied).	117
Table 9.	Mean values of different morphological parameters of five highly occupied gastropod shells by different reproductive stages (juveniles=1–3 mm; adults= 3–7 mm) of <i>Clibanarius rhabdodactylus</i> Male= M; Ovigerous female= OF; Female= F; Shield length= SL; Shell length= SHL; Shell aperture length= SHAL; Shell aperture width= SHAW; Shell dry weight= SHW; Shell volume= SHV; NO= Not occupied).	119
Table 10.	Mean values of different morphological parameters of five highly occupied gastropod shells by different reproductive stages (juveniles=1–3 mm; adults= 3–7 mm) of <i>Clibanarius ransonii</i> (N = total individuals; M = Male; OF = Ovigerous female; F = Female; SL = Shield length; SHL = Shell length; SHAL = Shell aperture length; SHAW = Shell aperture width; SHW = Shell dry weight; SHV = Shell volume; NO = Not occupied).	120
Table 11.	Groupings of hermit crab individuals on the basis of species, sex and size classes (shield length) with their annotated codes for canonical correspondence analysis (CCA). (N = total individuals).	125